


4: Discrete probability



Readings: BT 1.1-1.2, Rosen 6.1-6.2

Sample space: S is the set of all possible outcomes of an experiment (Ω in your text book—Greek uppercase omega)

Coin flip: $S = \{\text{Heads, Tails}\}$

Flipping two coins: $S = \{(H,H), (H,T), (T,H), (T,T)\}$

Roll of one 6-sided die: $S = \{1, 2, 3, 4, 5, 6\}$

emails in a day: $S = \{x : x \in \mathbb{Z}, x \geq 0\}$

YouTube hrs. in a day: $S = \{x : x \in \mathbb{R}, 0 \leq x \leq 24\}$

Events: $E \subseteq S$ is some subset of the sample space

Coin flip is heads: $E = \{\text{Head}\}$

At least one head in 2 flips: $E = \{(H,H), (H,T), (T,H)\}$


Roll of die is 3 or less: $E = \{1, 2, 3\}$

emails in a day < 20 : $E = \{x : x \in \mathbb{Z}, 0 \leq x < 20\}$

Wasted day (>5 YT hrs): $E = \{x : x \in \mathbb{R}, x > 5\}$

set operations on events


E and F are events in the sample space S



set operations on events

E and F are events in the sample space S

Event “ E OR F ”, written $E \cup F$




$S = \{1, 2, 3, 4, 5, 6\}$
outcome of one die roll

$E = \{1, 2\}$, $F = \{2, 3\}$
 $E \cup F = \{1, 2, 3\}$

set operations on events

E and F are events in the sample space S

Event “ E AND F ”, written $E \cap F$ or EF




$S = \{1,2,3,4,5,6\}$
outcome of one die roll

$E = \{1,2\}$, $F = \{2,3\}$
 $E \cap F = \{2\}$

set operations on events

E and F are events in the sample space S

$EF = \emptyset \Leftrightarrow E, F$ are “mutually exclusive”




$S = \{1, 2, 3, 4, 5, 6\}$
outcome of one die roll

$E = \{1, 2\}$, $F = \{2, 3\}$, $G = \{5, 6\}$
 $EF = \{2\}$, *not* mutually
exclusive, but E, G and F, G are

set operations on events

E and F are events in the sample space S

Event “not E ,” written \bar{E} or $\neg E$




$S = \{1, 2, 3, 4, 5, 6\}$
outcome of one die roll


$E = \{1, 2\}$ $\neg E = \{3, 4, 5, 6\}$

DeMorgan's Laws

$$\overline{E \cup F} = \bar{E} \cap \bar{F}$$



$$\overline{E \cap F} = \bar{E} \cup \bar{F}$$



axioms of probability

Intuition: Probability as the relative frequency of an event

$$\Pr(E) = \lim_{n \rightarrow \infty} (\# \text{ of occurrences of } E \text{ in } n \text{ trials})/n$$

Axiom 1: $0 \leq \Pr(E) \leq 1$

Axiom 2: $\Pr(S) = 1$

Axiom 3: If E and F are mutually exclusive ($EF = \emptyset$), then

$$\Pr(E \cup F) = \Pr(E) + \Pr(F)$$

For any sequence E_1, E_2, \dots, E_n of mutually exclusive events,

$$\Pr\left(\bigcup_{i=1}^n E_i\right) = \Pr(E_1) + \dots + \Pr(E_n)$$

implications of axioms

- $\Pr(\bar{E}) = 1 - \Pr(E)$

$$\Pr(\bar{E}) = \Pr(S) - \Pr(E) \text{ because } S = E \cup \bar{E}$$


- If $E \subseteq F$, then $\Pr(E) \leq \Pr(F)$

$$\Pr(F) = \Pr(E) + \Pr(F - E) \geq \Pr(E)$$

- $\Pr(E \cup F) = \Pr(E) + \Pr(F) - \Pr(EF)$

inclusion-exclusion formula

- And many others



equally likely outcomes

Simplest case: sample spaces with equally likely outcomes.

Coin flips: $S = \{\text{Heads, Tails}\}$

Flipping two coins: $S = \{(H,H), (H,T), (T,H), (T,T)\}$

Roll of 6-sided die: $S = \{1, 2, 3, 4, 5, 6\}$

$$\Pr(\text{each outcome}) = \frac{1}{|S|}$$

uniform distribution

In that case,

$$\Pr(E) = \frac{\text{number of outcomes in } E}{\text{number of outcomes in } S} = \frac{|E|}{|S|}$$

rolling two dice

Roll two 6-sided dice. What is $\Pr(\text{sum of dice} = 7)$?

$$S = \{ (1,1), (1,2), (1,3), (1,4), (1,5), (1,6), \\ (2,1), (2,2), (2,3), (2,4), (2,5), (2,6), \\ (3,1), (3,2), (3,3), (3,4), (3,5), (3,6), \\ (4,1), (4,2), (4,3), (4,4), (4,5), (4,6), \\ (5,1), (5,2), (5,3), (5,4), (5,5), (5,6), \\ (6,1), (6,2), (6,3), (6,4), (6,5), (6,6) \}$$

$$E = \{ (6,1), (5,2), (4,3), (3,4), (2,5), (1,6) \}$$

$$\Pr(\text{sum} = 7) = |E|/|S| = 6/36 = 1/6.$$

birthdays




What is the probability that, of n people, none share the same birthday?

$$|S| = (365)^n$$

$$|E| = (365)(364)(363)\cdots(365-n+1)$$

$$\begin{aligned}\text{Pr}(\text{no matching birthdays}) &= |E|/|S| \\ &= (365)(364)\cdots(365-n+1)/(365)^n\end{aligned}$$



Some values of n ...

$$n = 23: \text{Pr}(\text{no matching birthdays}) < 0.5$$

$$n = 77: \text{Pr}(\text{no matching birthdays}) < 1/5000$$

$$n = 100: \text{Pr}(\text{no matching birthdays}) < 1/3,000,000$$

$$n = 150: \text{Pr}(\dots) < 1/3,000,000,000,000,000$$

$n = 366?$

$Pr = 0$

Above formula gives this, since

$$(365)(364)\dots(365-n+1)/(365)^n == 0$$

when $n = 366$ (or greater).

Even easier to see via pigeon hole principle.

What is the probability that, of n people, none share the same birthday as you?

$$|S| = (365)^n$$

$$|E| = (364)^n$$

$$\begin{aligned} \text{Pr}(\text{no birthdays matches yours}) &= |E|/|S| \\ &= (364)^n/(365)^n \end{aligned}$$

Some values of n ...

$$n = 23: \quad \text{Pr}(\text{no matching birthdays}) \approx 0.9388$$

$$n = 77: \quad \text{Pr}(\text{no matching birthdays}) \approx 0.8096$$

$$n = 253: \quad \text{Pr}(\text{no matching birthdays}) \approx 0.4995$$

poker hands



any straight in poker

Consider 5 card poker hands.


A “straight” is 5 consecutive rank cards of any suit

What is $\Pr(\text{straight})$?


$$|S| = \binom{52}{5}$$

$$|E| = 10 \cdot \binom{4}{1}^5$$

$$\Pr(\text{straight}) = \frac{10 \binom{4}{1}^5}{\binom{52}{5}} \approx 0.00394$$



card flipping



card flipping

52 card deck. Cards flipped one at a time.

After first ace (of any suit) appears, consider next card

$\Pr(\text{next card} = \text{ace of spades}) < \Pr(\text{next card} = \text{2 of clubs}) ?$

Case 1: Take Ace of Spades out of deck

Shuffle remaining 51 cards, add ace of spades after first ace

$|S| = 52!$ (all cards shuffled)

$|E| = 51!$ (only 1 place ace of spades can be added)

Case 2: Do the same thing with the 2 of clubs







$|S|$ and $|E|$ have same size

So,

$\Pr(\text{next} = \text{Ace of spades}) = \Pr(\text{next} = \text{2 of clubs}) = 1/52$

Ace of Spades: 2/6

2 of Clubs: 2/6



Theory is the same for a 3-card deck; $Pr = 2!/3! = 1/3_{22}$

hats



i don't belong to a magician, i just really like hats.

n persons at a party throw hats in middle, select at random. What is $\Pr(\text{no one gets own hat})$?



$\Pr(\text{no one gets own hat}) =$
 $1 - \Pr(\text{someone gets own hat})$

$\Pr(\text{someone gets own hat}) = \Pr(\bigcup_{i=1}^n E_i)$, where
 E_i = event that person i gets own hat

$$\Pr(\bigcup_{i=1}^n E_i) = \sum_i P(E_i) - \sum_{i < j} \Pr(E_i E_j) + \sum_{i < j < k} \Pr(E_i E_j E_k) \dots$$

Visualizing the sample space S :

People:

P_1	P_2	P_3	P_4	P_5
-------	-------	-------	-------	-------

Hats:

H_4	H_2	H_5	H_1	H_3
-------	-------	-------	-------	-------




I.e., a sample point is a *permutation* π of $1, \dots, n$

4	2	5	1	3
---	---	---	---	---


$$|S| = n!$$

hats: events

E_i = event that person i gets own hat: $\pi(i) = i$




Counting single events:



$$|E_i| = (n-1)! \text{ for all } i$$

Counting pairs:

$$E_i E_j : \pi(i) = i \ \& \ \pi(j) = j$$



$$|E_i E_j| = (n-2)! \text{ for all } i, j$$

All points in $E_2 \cap E_5$

n persons at a party throw hats in middle, select at random. What is $\Pr(\text{no one gets own hat})$?



E_i = event that person i gets own hat

$$\Pr(\bigcup_{i=1}^n E_i) = \sum_i P(E_i) - \sum_{i < j} \Pr(E_i E_j) + \sum_{i < j < k} \Pr(E_i E_j E_k) \dots$$

$$\Pr(k \text{ fixed people get own back}) = (n-k)!/n!$$

$$\binom{n}{k} \text{ times that} = \frac{n!}{k!(n-k)!} \frac{(n-k)!}{n!} = 1/k!$$

$$\Pr(\text{none get own}) = 1 - \Pr(\text{some do}) =$$

$$1 - 1/1! + 1/2! - 1/3! + 1/4! \dots + (-1)^n/n! \approx 1/e \approx .37$$

$$\Pr(\text{none get own}) = 1 - \Pr(\text{some do}) =$$
$$1 - \left(1 + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} \dots + \frac{(-1)^n}{n!}\right) \approx e^{-1} \approx .37$$

