## **CSE 312 Practice Midterm**

1. True/False, Short Answer. Provide a short justification for your answer.

a) True or False. For any random variables X and Y,  $Var(aX + bY + c) = a^2 Var(X) + b^2 Var(Y)$ .

b) Let X be a discrete random variable with pmf  $p(x) = a^x$ , x = 1, 2, ... Find a, or say if there is not enough information.

c) True or False. Suppose  $F_1, F_2, ..., F_n$  partition the sample space, and let E be any event. Then,  $P(E) = P(E|F_1) + P(E|F_2) + \cdots + P(E|F_n)$ 

d) True or False. For any random variable X,  $\log E[X] = E[\log X]$ .

e) If  $Var(2X) = E[3X^2] = 12$ , find E[X], or say if there is not enough information.

f) True or False. If *E* and *F* are independent with  $P(E) \neq 0$ ,  $P(F) \neq 0$ , they are mutually exclusive.

g) True or False. If *E* and *F* are mutually exclusive with  $P(E) \neq 0$ ,  $P(F) \neq 0$ , they are independent.

h) True or False. If P(A) = 0.4, P(B) = 0.3, and  $P(A \cup B) = 0.58$ , then A and B are independent.

2. Annie, Bill, Charlie, Danielle, Ethan, Felicia, and George are sitting in a row of <u>nine</u> seats (Note: there are only seven people). Annie and Bill are exes, so they cannot sit next to each other. Charlie and Danielle are dating, so they must sit next to each other. Ethan, Felicia, and George are best friends, so they also want to sit next to each other, but Felicia must be in the middle of Ethan and George (with no spaces between the three). In how many ways can they sit in the row of nine seats? (Hint: Start by grouping Ethan, Felicia, and George, as well as Charlie and Danielle. Then work on placing Annie and Bill).

3. The hypergeometric distribution represents the following scenario: Suppose there are N items in a bag, with K of them marked as successes in total (and the rest are marked as failures). We draw n of them, <u>without</u> replacement. Each item is equally likely to be drawn. Then, if X is the random variable that denotes the number of successes we get in the n draws, we say  $X \sim HypGeo(N, K, n)$ .

a) Based on the situation, specify restrictions on the relationships among N, K, and n.

b) Find the probability mass function for the hypergeometric random variable X, p(k). Be sure to specify the domain for the pmf.

 4. Patients arriving at a hospital emergency room are categorized as either being in critical, serious, or stable condition. In the last year, we know

- 60% of the patients were stable
- 10% of the patients were critical
- 40% of critical patients died
- 90% of serious patients survived
- 1% of stable patients died

Given that a patient survived, what is the probability that the patient was categorized as serious?

5. Let  $X_1, X_2, \dots, X_n$  be iid (independent and identically distributed) random variables each with pmf

$$p(x) = \begin{cases} 1/3, & x \in \{2,3\} \\ p, & x \in \{0,1\} \\ 0, & otherwise \end{cases}$$

Furthermore, let  $X = X_1 + X_2 + \dots + X_n$ 

a) Find p.

b) Find E[-3X - 4].

c) Find Var(-3X - 4), and Var(3X + 4).