

# CSE 332: Data Structures and Parallelism




## Section 2: Heaps, Asymptotics, & Recurrences Solutions

### 0. Heaps

Insert 10, 7, 15, 17, 12, 20, 6, 32 into a *min heap*.

Now, insert the same values into a *max heap*.

Now, insert the same values into a *min heap*, but use Floyd's buildHeap algorithm. **Solution:**



## 1. Big-Oh Proofs

For each of the following, prove that  $f \in \mathcal{O}(g)$ .

(a)  $f(n) = 7n$   $g(n) = \frac{n}{10}$

**Solution:** Choose  $c = 70$ ,  $n_0 = 1$ . Then, note that  $7n = \frac{70n}{10} \leq 70 \left(\frac{n}{10}\right)$  for all  $n \geq 1$ . So,  $f(n) \in \mathcal{O}(g(n))$ .

(b)  $f(n) = 1000$   $g(n) = 3n^3$

**Solution:** Choose  $c = 1$ ,  $n_0 = 1000$ . Then, note that  $1000 \leq n \leq n^3 \leq 3n^3$  for all  $n \geq 1000$ . So,  $f(n) \in \mathcal{O}(g(n))$ .

(c)  $f(n) = 7n^2 + 3n$   $g(n) = n^4$

**Solution:** Choose  $c = 14$ ,  $n_0 = 1$ . Then, note that  $7n^2 + 3n \leq 7(n^4 + n^4) \leq 14n^4$  for all  $n \geq 1$ . So,  $f(n) \in \mathcal{O}(g(n))$ .

(d)  $f(n) = n + 2n \lg n$   $g(n) = n \lg n$

**Solution:** Choose  $c = 3$ ,  $n_0 = 2$ . Then, note that  $n + 2n \lg n \leq n \lg n + 2n \lg n = 3n \lg n$  for all  $n \geq 2$ . So,  $f(n) \in \mathcal{O}(g(n))$ .

## 2. Is Your Program Running? Better Catch It!

For each of the following, determine the asymptotic worst-case runtime in terms of  $n$ .

(a)

```
1 int x = 0;
2 for (int i = n; i >= 0; i--) {
3     if ((i % 3) == 0) {
4         break;
5     }
6     else {
7         x += n;
8     }
9 }
```

**Solution:** This is  $\Theta(1)$ , because  $n$ ,  $n - 1$ , or  $n - 2$  will be divisible by three. So, the loop runs at most 3 times.

(b)

```
1 int x = 0;
2 for (int i = 0; i < n; i++) {
3     for (int j = 0; j < (n * n / 3); j++) {
4         x += j;
5     }
6 }
```

**Solution:**

$$\sum_{i=0}^{n-1} \sum_{j=0}^{n^2/3-1} 1 = \sum_{i=0}^{n-1} \frac{n^2}{3} = n \left( \frac{n^2}{3} \right) \in \Theta(n^3)$$

(c)

```
1 int x = 0;
2 for (int i = 0; i <= n; i++) {
3     for (int j = 0; j < (i * i); j++) {
4         x += j;
5     }
6 }
```

**Solution:**

$$\sum_{i=0}^n \sum_{j=0}^{i^2-1} 1 = \sum_{i=0}^n i^2 = \left( \frac{n(n+1)(2n+1)}{6} \right) \in \Theta(n^3)$$

### 3. Induction Shmduction

Prove  $\sum_{i=0}^n 2^i = 2^{n+1} - 1$  by induction on  $n$ .

**Solution:**

Let  $P(n)$  be the statement " $\sum_{i=0}^n 2^i = 2^{n+1} - 1$ " for all  $n \in \mathbb{N}$ . We prove  $P(n)$  by induction on  $n$ .

**Base Case.** Note that  $\sum_{i=0}^0 2^i = 0 = 2^0 - 1$ . So,  $P(0)$  is true.

**Induction Hypothesis.** Suppose  $P(k)$  is true for some  $k \in \mathbb{N}$ .

**Induction Step.** Note that

$$\begin{aligned} \sum_{i=0}^{k+1} 2^i &= \sum_{i=0}^k 2^i + 2^{k+1} \\ &= 2^{k+1} - 1 + 2^{k+1} && \text{[By IH]} \\ &= 2^{k+2} - 1 \end{aligned}$$

Note that this is exactly  $P(k+1)$ .

So, the claim is true by induction on  $n$ .

### 4. The Implications of Asymptotics

For each of the following, determine if the statement is true or false.

(a)  $f(n) \in \Theta(g(n)) \rightarrow f(n) \in \mathcal{O}(g(n))$

**Solution:**

This is true. By definition of  $f(n) \in \Theta(g(n))$ , we have  $f(n) \in \mathcal{O}(g(n))$ .

(b)  $f(n) \in \Theta(g(n)) \rightarrow g(n) \in \Theta(f(n))$

**Solution:**

This is true. By definition of  $f(n) \in \Theta(g(n))$ , we have  $f(n) \in \mathcal{O}(g(n))$  and  $f(n) \in \Omega(g(n))$ . So, there exist  $n_0, n_1, c_0, c_1 > 0$  such that  $f(n) \leq c_0 g(n)$  for all  $n \geq n_0$  and  $f(n) \geq c_1 g(n)$  for all  $n \geq n_1$ . Define  $n_2 = \max(n_0, n_1)$  and note that both inequalities hold for all  $n \geq n_2$ . Then, dividing both sides by their constants, we have:

$$\begin{aligned} g(n) &\geq \frac{f(n)}{c_0} \\ g(n) &\leq \frac{f(n)}{c_1} \end{aligned}$$

So, we've found constants  $\left(\frac{1}{c_0}, \frac{1}{c_1}\right)$  and a minimum  $n$  ( $n_2$ ) that satisfy the definitions of Omega and Oh. It follows that  $g(n)$  is  $\Theta(f(n))$ .

(c)  $f(n) \in \Omega(g(n)) \rightarrow g(n) \in \mathcal{O}(f(n))$

**Solution:**

This is true. This is basically identical to the previous part (except we only have to do half the work).

## 5. Asymptotic Analysis

For each of the following, determine if  $f \in \mathcal{O}(g)$ ,  $f \in \Omega(g)$ ,  $f \in \Theta(g)$ , several of these, or none of these.

(a)  $f(n) = \log n$   $g(n) = \log \log n$

**Solution:**  $f(n) \in \Omega(g(n))$

(b)  $f(n) = 2^n$   $g(n) = 3^n$

**Solution:**  $f(n) \in \mathcal{O}(g(n))$

(c)  $f(n) = 2^{2n}$   $g(n) = 2^n$

**Solution:**  $f(n) \in \Omega(g(n))$