

CSE 417

Algorithms and Complexity

Autumn 2023
Lecture 15
Divide and Conquer and Recurrences

11/1/2023

CSE 417

1

Announcements

11/1/2023

CSE 417

2

Divide and Conquer

- Recurrences, Sections 5.1 and 5.2
- Algorithms
 - Median (Selection)
 - Fast Matrix Multiplication
 - Counting Inversions (5.3)
 - Multiplication (5.5)

11/1/2023

CSE 417

3

Divide and Conquer : Merge Sort

```
Array Mergesort(Array a){  
    n = a.Length;  
    if (n <= 1)  
        return a;  
    b = Mergesort(a[0 .. n/2]);  
    c = Mergesort(a[n/2+1 .. n-1]);  
    return Merge(b, c);  
}
```

11/1/2023

CSE 417

4

Algorithm Analysis

- Cost of Merge
- Cost of Mergesort

11/1/2023

CSE 417

5

$$T(n) = 2T(n/2) + cn; T(1) = c;$$

11/1/2023

CSE 417

6

Recurrence Analysis


- Solution methods
 - Unrolling recurrence
 - Guess and verify
 - Plugging in to a “Master Theorem”

11/1/2023

CSE 417

7

Unrolling the recurrence



11/1/2023

CSE 417

8

$$T(n) = 2T(n/2) + n; T(1) = 1;$$

Substitution

Prove $T(n) \leq n(\log_2 n + 1)$ for $n \geq 1$

Induction:

Base Case:

Induction Hypothesis:

11/1/2023

CSE 417

9

A better mergesort (?)

- Divide into 3 subarrays and recursively sort
- Apply 3-way merge

What is the recurrence?

11/1/2023

CSE 417

10

Unroll recurrence for $T(n) = 3T(n/3) + n$

11/1/2023

CSE 417

11

$$T(n) = aT(n/b) + f(n)$$

11/1/2023

CSE 417

12

$$T(n) = T(n/2) + cn$$

Where does this recurrence arise?


Quicksort

QuickSort(S):

1. Pick an element v in S . This is the **pivot** value.
2. Partition $S - \{v\}$ into two disjoint subsets, S_1 and S_2 such that:
 - elements in S_1 are all $< v$
 - elements in S_2 are all $> v$
3. Return concatenation of QuickSort(S_1), v , QuickSort(S_2)

Recursion ends if QuickSort() receives an array of length 0 or 1.

The steps of Quicksort



Picking the pivot

- Choose the first element in the subarray
- Choose a value that might be close to the middle
 - Median of three
- Choose a random element

Recurrence for Quicksort

$$QS(n) = \sum_{i=1}^n \frac{1}{n} \{QS(i-1) + QS(n-i)\}$$

Computing the Median

- Given n numbers, find the number of rank $n/2$
- One approach is sorting
 - Sort the elements, and choose the middle one
 - Can you do better?

Problem generalization

- *Selection*, given n numbers and an integer k , find the k -th largest

11/1/2023

CSE 417

19

Select(A, k)

```

Select(A, k){
  Choose element x from A
  S1 = {y in A | y < x}
  S2 = {y in A | y > x}
  S3 = {y in A | y = x}
  if (|S2| >= k)
    return Select(S2, k)
  else if (|S2| + |S3| >= k)
    return x
  else
    return Select(S1, k - |S2| - |S3|)
}
    
```



11/1/2023

CSE 417

20

Randomized Selection

- Choose the element at random
- Analysis can show that the algorithm has expected run time $O(n)$

11/1/2023

CSE 417

21

$$T(n) = T(n/2) + cn$$

Where does this recurrence arise?

11/1/2023

CSE 417

22

Solving the recurrence exactly

11/1/2023


CSE 417

23

Total Work

$$\sum_{k=0}^{\log n} 2^k n = (2n - 1)n$$

$$T(n) = 4T(n/2) + n$$




11/1/2023

CSE 417

24

$$T(n) = 2T(n/2) + n^2$$




11/1/2023

CSE 417

25

$$T(n) = 2T(n/2) + n^{1/2}$$



11/1/2023

CSE 417

26

Recurrences

- Three basic behaviors
 - Dominated by initial case
 - Dominated by base case
 - All cases equal – we care about the depth

11/1/2023

CSE 417

27

What you really need to know about recurrences

- Work per level changes geometrically with the level
 - The bottom level wins
- Geometrically increasing ($x > 1$)
 - The bottom level wins
- Geometrically decreasing ($x < 1$)
 - The top level wins
- Balanced ($x = 1$)
 - Equal contribution

11/1/2023

CSE 417

28

Classify the following recurrences (Increasing, Decreasing, Balanced)

- $T(n) = n + 5T(n/8)$
- $T(n) = n + 9T(n/8)$
- $T(n) = n^2 + 4T(n/2)$
- $T(n) = n^3 + 7T(n/2)$
- $T(n) = n^{1/2} + 3T(n/4)$

11/1/2023

CSE 417

29