

Sample midterm questions

**Instructions:**

- Exam will consist of 5 to 8 questions.
- Closed book, closed notes, no cell phones, no calculators.
- Time limit: 50 minutes.
- Answer the problems on the exam paper.
- If you need extra space use the back of a page.
- Lists of equivalences and inference rules for your use are given on the final two pages.

**Problem 1:**

- Show that the expression  $(p \rightarrow q) \rightarrow (p \rightarrow r)$  is a contingency.
- Give an expression that is logically equivalent to  $(p \rightarrow q) \rightarrow (p \rightarrow r)$  using the logical operators  $\neg$ ,  $\vee$ , and  $\wedge$  (but not  $\rightarrow$ ).

**Problem 2:**

Using the predicates:

$Likes(p, f)$ : "Person  $p$  likes to eat the food  $f$ ."

$Serves(r, f)$ : "Restaurant  $r$  serves the food  $f$ ."

translate the following statements into logical expressions.

- Every restaurant serves a food that no one likes.
- Every restaurant that serves TOFU also serves a food which RANDY does not like.

**Problem 3:**

Use rules of inference to show that if the premises  $\forall x(P(x) \rightarrow Q(x))$ ,  $\forall x(Q(x) \rightarrow R(x))$ , and  $\neg R(a)$ , where  $a$  is in the domain, are true, then the conclusion  $\neg P(a)$  is true. (Note: You do not need to give the names for the rules of inference.)

**Problem 4:**

Prove that if  $n$  is even and  $m$  is odd, then  $(n + 1)(m + 1)$  is even.

**Problem 5:**

Prove or disprove:

- a) For positive integers  $x$ ,  $p$ , and  $q$ ,  $(x \bmod p) \bmod q = x \bmod pq$ .
- b) For positive integers  $x$ ,  $p$ , and  $q$ ,  $(x \bmod p) \bmod q = (x \bmod q) \bmod p$ .

**Problem 6:**

- a) Find the multiplicative inverse of 2 modulo 9 (in other words, find a solution to the equation  $2x \bmod 9 = 1$ .)
- b) Which integers in  $\{1, 2, \dots, 8\}$  have multiplicative inverses modulo 9?

**Problem 7:**

Let  $T(n)$  be defined by:  $T(0) = 1$ ,  $T(n) = 2nT(n - 1)$  for  $n \geq 1$ . Prove that for all  $n \geq 0$ ,  $T(n) = 2^n n!$ .

**Problem 8:**

Let  $x_1, x_2, \dots, x_n$  be odd integers. Prove by induction that  $x_1 x_2 \cdots x_n$  is also an odd integer.

**Problem 9:**

Determine whether the following compound proposition is a tautology, a contradiction, or a contingency:  $((s \vee p) \wedge (s \vee \neg p)) \rightarrow ((p \rightarrow q) \rightarrow r)$ .

**Problem 10:**

Find predicates  $P(x)$  and  $Q(x)$  such that  $\forall x(P(x) \oplus Q(x))$  is true, but  $\forall x P(x) \oplus \forall x Q(x)$  is false.

**Problem 11:**

Show that the following is a tautology:  $((\neg p \vee q) \wedge (p \vee r)) \rightarrow (q \vee r)$ .

**Problem 12:**

Prove that the sum of an odd number and an even number is an odd number.

**Problem 13:**

Use mathematical induction to show that 3 divides  $n^3 - n$  whenever  $n$  is a non-negative integer.

**Problem 14:**

Let the predicates  $D(x, y)$  mean “team  $x$  defeated team  $y$ ” and  $P(x, y)$  mean “team  $x$  has played team  $y$ .” Give quantified formulas with the following meanings:

- a) Every team has lost at least one game.
- b) There is a team that has beaten every team it has played.

**Problem 15:**

Let  $P(x, y)$  be the predicate “ $x < y$ ” and let the universe for all variables be the real numbers. Express each of the following statements as predicate logic formulas using  $P$ :

- a) For every number there is a smaller one.
- b) 7 is smaller than any other number.
- c) 7 is between  $a$  and  $b$ . (Don't forget to handle both the possibility that  $b$  is smaller than  $a$  as well as the possibility that  $a$  is smaller than  $b$ .)
- d) Between any two different numbers there is another number.
- e) For any two numbers, if they are different then one is less than the other.

**Problem 16:**

Let  $V(x, y)$  be the predicate “ $x$  voted for  $y$ ”, let  $M(x, y)$  be the predicate “ $x$  received more votes than  $y$ ”, and let the universe for all variables be the set of all people. Express each of the following statements as predicate logic formulas using  $V$  and  $M$ :

- a) Everybody received at least one vote.
- b) Jane and John voted for the same person.
- c) Ross won the election. (The winner is the person who received the most votes.)
- d) Nobody who votes for him/herself can win the election.
- e) Everybody can vote for at most one person.

**Problem 17:**

Prove the following for all natural numbers  $n$  by induction,  $\sum_{i=0}^n \frac{i}{2^i} = 2 - \frac{n+2}{2^n}$ .

**Problem 18:**

Use Euclid's algorithm to help you solve  $11x \equiv 4 \pmod{27}$  for  $x$ .

**Problem 19:**

Write an expression equivalent to  $(p \rightarrow q) \rightarrow r$  that is:

- a) A sum of products
- b) A product of sums

Equivalences	
Identity Laws	$p \wedge \mathbf{T} \equiv p$ $p \vee \mathbf{F} \equiv p$
Domination Laws	$p \vee \mathbf{T} \equiv \mathbf{T}$ $p \wedge \mathbf{F} \equiv \mathbf{F}$
Idempotent Laws	$p \vee p \equiv p$ $p \wedge p \equiv p$
Commutative Laws	$p \vee q \equiv q \vee p$ $p \wedge q \equiv q \wedge p$
Associative Laws	$(p \vee q) \vee r \equiv p \vee (q \vee r)$ $(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$
Distributive Laws	$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$ $p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$
De Morgan's Laws	$\neg(p \wedge q) \equiv \neg p \vee \neg q$ $\neg(p \vee q) \equiv \neg p \wedge \neg q$
Negation Laws	$p \vee \neg p \equiv \mathbf{T}$ $p \wedge \neg p \equiv \mathbf{F}$
Double Negation Law	$\neg\neg p \equiv p$
Contrapositive Law	$p \rightarrow q \equiv \neg q \rightarrow \neg p$
Implication Law	$p \rightarrow q \equiv \neg p \vee q$
Quantifier Negation Laws	$\neg\exists xP(x) \equiv \forall x\neg P(x)$ $\neg\forall xP(x) \equiv \exists x\neg P(x)$

Propositional and Predicate Equivalences

Inferences	
Modus Ponens	$\frac{p, p \rightarrow q}{\therefore q}$
Direct Proof	$\frac{p \Rightarrow q}{\therefore p \rightarrow q}$
Elim $\wedge$	$\frac{p \wedge q}{\therefore p, q}$
Intro $\wedge$	$\frac{p, q}{\therefore p \wedge q}$
Elim $\vee$	$\frac{p \vee q, \neg p}{\therefore q}$
Intro $\vee$	$\frac{p}{\therefore p \vee q, q \vee p}$
Excluded Middle	$\frac{}{\therefore p \vee \neg p}$
Elim $\forall$	$\frac{\forall x P(x)}{\therefore P(a) \text{ for any } a}$
Intro $\forall$	$\frac{\text{Let } a \text{ be anything...} P(a)}{\therefore \forall x P(x)}$
Elim $\exists$	$\frac{\exists x P(x)}{\therefore P(c) \text{ for some special } c}$
Intro $\exists$	$\frac{P(c) \text{ for some } c}{\therefore \exists x P(x)}$

Propositional and Predicate Inferences