


# cse 311: foundations of computing

---

Fall 2015

## Lecture 7: Proofs



## an inference rule: *Modus Ponens*

---

- If  $p$  and  $p \rightarrow q$  are both true then  $q$  must be true
- Write this rule as 
$$\frac{p, p \rightarrow q}{\therefore q}$$
- Given:
  - If it is Monday then you have a 311 class today.
  - It is Monday.
- Therefore, by modus ponens:
  - You have a 311 class today.

---

Show that  $r$  follows from  $p$ ,  $p \rightarrow q$ , and  $q \rightarrow r$

1.  $p$  given
2.  $p \rightarrow q$  given
3.  $q \rightarrow r$  given
4.  $q$  modus ponens from 1 and 2
5.  $r$  modus ponens from 3 and 4

- Each **inference rule** is written as: 
$$\frac{A, B}{\therefore C, D}$$

...which means that if both A and B are true then you can infer C and you can infer D.

- For rule to be correct  $(A \wedge B) \rightarrow C$  and  $(A \wedge B) \rightarrow D$  must be a tautologies
- Sometimes rules don't need anything to start with. These rules are called **axioms**:
  - e.g. *Excluded Middle Axiom*

$$\frac{}{\therefore p \vee \neg p}$$

# proofs can use equivalences too

---

Show that  $\neg p$  follows from  $p \rightarrow q$  and  $\neg q$

1.  $p \rightarrow q$  given
2.  $\neg q$  given
3.  $\neg q \rightarrow \neg p$  contrapositive of 1
4.  $\neg p$  modus ponens from 2 and 3

# important: applications of inference rules

---

- You can use equivalences to make substitutions of any sub-formula.
- Inference rules only can be applied to whole formulas (not correct otherwise)

e.g. ~~1.  $p \rightarrow q$  given~~  
~~2.  $(p \vee r) \rightarrow q$  intro  $\vee$  from 1.~~

**Does not follow!** e.g .  $p=F, q=F, r=T$

# simple propositional inference rules

---

Excluded middle plus two inference rules per binary connective, one to eliminate it and one to introduce it:

Elim- $\wedge$

$$\frac{p \wedge q}{\therefore p, q}$$

$$\frac{p, q}{\therefore p \wedge q}$$

Intro- $\wedge$

Elim- $\vee$

$$\frac{p \vee q, \neg p}{\therefore q}$$

$$\frac{p}{\therefore p \vee q, q \vee p}$$

Intro- $\vee$

MP

$$\frac{p, p \rightarrow q}{\therefore q}$$

$$\frac{p \Rightarrow q}{\therefore p \rightarrow q}$$

Direct Proof Rule  
Not like other rules

# direct proof of an implication

---

- $p \Rightarrow q$  denotes a proof of  $q$  given  $p$  as an assumption
- **The direct proof rule:**  
If you have such a proof then you can conclude that  $p \rightarrow q$  is true

Example:

		proof subroutine
	1. $p$	<b>assumption</b>
2.	$p \vee q$	intro for $\vee$ from 1
3.	$p \rightarrow (p \vee q)$	direct proof rule



# proofs using the direct proof rule

---

Show that  $p \rightarrow r$  follows from  $q$  and  $(p \wedge q) \rightarrow r$

1.  $q$  given
2.  $(p \wedge q) \rightarrow r$  given
3.  $p$  assumption
4.  $p \wedge q$  from 1 and 3 via Intro  $\wedge$  rule
5.  $r$  modus ponens from 2 and 4
6.  $p \rightarrow r$  direct proof rule

Prove:  $(p \wedge q) \rightarrow (p \vee q)$

Prove:  $((p \rightarrow q) \wedge (q \rightarrow r)) \rightarrow (p \rightarrow r)$

# one general proof strategy

---

1. Look at the rules for introducing connectives to see how you would build up the formula you want to prove from pieces of what is given
2. Use the rules for eliminating connectives to break down the given formulas so that you get the pieces you need to do (1).
3. Write the proof beginning with what you figured out for (2) followed by (1).

# inference rules for quantifiers

---

Intro- $\exists$

$P(c)$  for some  $c$

---

$\therefore \exists x P(x)$

Elim- $\forall$

$\forall x P(x)$

---

$\therefore P(a)$  for any  $a$

Intro- $\forall$

“Let  $a$  be anything\*” ... $P(a)$

---

$\therefore \forall x P(x)$

Elim- $\exists$

$\exists x P(x)$

---

$\therefore P(c)$  for some *special\*\**  $c$

\* in the domain of  $P$

\*\* By special, we mean that  $c$  is a name for a value where  $P(c)$  is true. We can't use anything else about that value, so  $c$  has to be a NEW variable!

# proofs using quantifiers

---

“There exists an even prime number.”

Prime( $x$ ):  $x$  is an integer  $> 1$  and  $x$  is not a multiple of any integer strictly between 1 and  $x$

# proofs using quantifiers

---

- |    |   |                       |
|----|---|-----------------------|
| 1. | $\text{Even}(2)$                                    | Fact* (math)          |
| 2. | $\text{Prime}(2)$                                   | Fact* (math)          |
| 3. | $\text{Even}(2) \wedge \text{Prime}(2)$             | Intro $\wedge$ : 1, 2 |
| 4. | $\exists x (\text{Even}(x) \wedge \text{Prime}(x))$ | Intro $\exists$ : 3   |

Those first two lines are sort of cheating; we should prove those “facts”.

- |    |                                       |                              |
|----|---------------------------------------|------------------------------|
| 1. | $2 = 2*1$                             | Definition of Multiplication |
| 2. | $\text{Even}(2)$                      | Intro $\exists$ : 1          |
| 3. | There are no integers between 1 and 2 | Definition of Integers       |
| 4. | 2 is an integer                       | Definition of 2              |
| 5. | $\text{Prime}(2)$                     | Intro $\wedge$ : 3, 4        |

**Prime(x):** x is an integer  $> 1$  and x is not a multiple of any integer strictly between 1 and x

**Even(x)**  $\equiv \exists y (x=2y)$

# proofs using quantifiers

---

1.	$2 = 2*1$	Definition of Multiplication
2.	$\text{Even}(2)$	Intro $\exists$ : 1
3.	There are no integers between 1 and 2	Definition of Integers
4.	2 is an integer	Definition of 2
5.	$\text{Prime}(2)$	Intro $\wedge$ : 3, 4
6.	$\text{Even}(2) \wedge \text{Prime}(2)$	Intro $\wedge$ : 2, 5
7.	$\exists x (\text{Even}(x) \wedge \text{Prime}(x))$	Intro $\exists$ : 7

English version:

“Note that  $2 = 2*1$  by definition of multiplication. It follows that there is a  $y$  such that  $2 = 2y$ ; so, 2 is even. Furthermore, 2 is an integer, and there are no integers between 1 and 2; so, by definition of a prime number, 2 is prime. Since 2 is both even and prime,  $\exists x (\text{Even}(x) \wedge \text{Prime}(x))$ .”

**Prime(x):**  $x$  is an integer  $> 1$  and  $x$  is not a multiple of any integer strictly between 1 and  $x$

**Even(x)**  $\equiv \exists y (x=2y)$



Prove: “The square of every even number is even.”

Formal proof of:  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

Even(x)  $\equiv \exists y (x=2y)$   
Odd(x)  $\equiv \exists y (x=2y+1)$   
Domain: Integers

Prove: “The square of every even number is even.”

Formal proof of:  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

- |  |  |
|--|--|
| 1. $\text{Even}(a)$  | Assumption: $a$ arbitrary integer              |
| 2. $\exists y (a = 2y)$                                      | Definition of Even                             |
| 3. $a = 2c$  | By elim $\exists$ : $c$ special depends on $a$ |
| 4. $a^2 = 4c^2 = 2(2c^2)$                                    | Algebra  |
| 5. $\exists y (a^2 = 2y)$                                    | By intro $\exists$ rule                        |
| 6. $\text{Even}(a^2)$  | Definition of Even                             |
| 7. $\text{Even}(a) \rightarrow \text{Even}(a^2)$             | Direct proof rule                              |
| 8. $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$ | By intro $\forall$ rule                        |

$\text{Even}(x) \equiv \exists y (x=2y)$   
 $\text{Odd}(x) \equiv \exists y (x=2y+1)$   
Domain: Integers

Prove: “The square of every odd number is odd.”

English proof of:  $\forall x (\text{Odd}(x) \rightarrow \text{Odd}(x^2))$

Even(x)  $\equiv \exists y (x=2y)$   
Odd(x)  $\equiv \exists y (x=2y+1)$   
Domain: Integers

Prove: “The square of every odd number is odd.”

English proof of:  $\forall x (\text{Odd}(x) \rightarrow \text{Odd}(x^2))$

Let  $x$  be an odd number.

Then  $x = 2k + 1$  for some integer  $k$  (depending on  $x$ )

Therefore  $x^2 = (2k+1)^2 = 4k^2 + 4k + 1 = 2(2k^2+2k) + 1$ .

Since  $2k^2 + 2k$  is an integer,  $x^2$  is odd.  $\square$

Even( $x$ )  $\equiv \exists y (x=2y)$   
Odd( $x$ )  $\equiv \exists y (x=2y+1)$   
Domain: Integers

## proof by contradiction: one way to prove $\neg p$

If we assume  $p$  and derive False (a contradiction), then we have proved  $\neg p$ .

1.  $p$       assumption

...

3. **F**

4.  $p \rightarrow \mathbf{F}$       direct Proof rule

5.  $\neg p \vee \mathbf{F}$       equivalence from 4

6.  $\neg p$       equivalence from 5