

Fall 2015

Lecture 27: Infinities and diagonalization

*54.43. $\vdash : \alpha, \beta \in 1. \supset : \alpha \cap \beta = \Lambda. \equiv . \alpha \cup \beta \in 2$

Dem.

$\vdash . *54.26. \supset \vdash : \alpha = \iota'x. \beta = \iota'y. \supset : \alpha \cup \beta \in 2. \equiv . x \neq y.$

[*51.231] $\equiv . \iota'x \cap \iota'y = \Lambda.$

[*13.12] $\equiv . \alpha \cap \beta = \Lambda$ (1)

$\vdash . (1). *11.11.35. \supset$

$\vdash : (\exists x, y). \alpha = \iota'x. \beta = \iota'y. \supset : \alpha \cup \beta \in 2. \equiv . \alpha \cap \beta = \Lambda$ (2)

$\vdash . (2). *11.54. *52.1. \supset \vdash . \text{Prop}$

From this proposition it will follow, when arithmetical addition has been defined, that $1 + 1 = 2$.


[proved on page 86 of Volume II:
"The above proposition is
occasionally useful."]

Russell's paradox: The set S of all
sets that do not contain themselves.

$$S \in S \Rightarrow S \notin S \Rightarrow S \in S$$

AUTHOR KATHARINE GATES RECENTLY ATTEMPTED
TO MAKE A CHART OF ALL SEXUAL FETISHES.

LITTLE DID SHE KNOW THAT RUSSELL AND WHITEHEAD
HAD ALREADY FAILED AT THIS SAME TASK.



computers from thought

Computers as we know them grew out of a desire to avoid bugs in mathematical reasoning.

Hilbert gave a famous speech at the International Congress of Mathematicians in 1900.

His goal was to **mechanize all of mathematics**.

In the 1930s, work of Gödel and Turing showed that Hilbert's program is **impossible**.

Gödel's incompleteness theorem
Undecidability of the Halting Problem

Both of these employ an idea we will see today called **diagonalization**.

The ideas are simple but so revolutionary that the inventor Georg Cantor was shunned by the mathematical leaders of the time:

Poincaré referred to them as a "grave disease infecting mathematics."

Kronecker fought to keep Cantor's papers out of his journals.

Cantor spent the last 30 years of his life battling depression, living often in "sanatoriums" (psychiatric hospitals).



cardinality

What does it mean that two sets have the same size?




cardinality

What does it mean that two sets have the same size?



Definition: Two sets A and B have the same **cardinality** if there is a one-to-one correspondence between the elements of A and those of B . More precisely, if there is a **1-1 and onto** function $f : A \rightarrow B$.



The definition also makes sense for infinite sets!

Do the natural numbers and the even natural numbers have the same cardinality?

Yes!

0	1	2	3	4	5	6	7	8	9	10	11	12	13	14
0	2	4	6	8	10	12	14	16	18	20	22	24	26	28

What's the map $f : \mathbb{N} \rightarrow 2\mathbb{N}$?

$$f(n) = 2n$$

countable sets

Definition: A set is **countable** iff it has the same cardinality as \mathbb{N} .

Equivalent: A set S is countable iff there is an 1-1 and onto function

$$g : \mathbb{N} \rightarrow S$$

Equivalent: A set S is countable iff we can order the elements

$$S = \{x_1, x_2, x_3, \dots\}$$

Question:

If $g : \mathbb{N} \rightarrow S$ is just **onto**, do we still know that S is countable?

$$g(n) = n \bmod 2 \quad S = \{0, 1\}$$

Definition: A set S is “at most countable” if it is finite or countable.

the set \mathbb{Z} of all integers

$$\mathbb{Z} = \{ \dots, -3, -2, -1, 0, 1, 2, 3, \dots \}$$

0	1	2	3	4	5	6
0	-1	1	-2	2	-3	3

$$f(n) = (-1)^n \left\lfloor \frac{n}{2} \right\rfloor$$

0	-1	1	-2	2	-3	3	...
---	----	---	----	---	----	---	-----

$$= \begin{cases} \frac{n}{2} & \text{if } n \text{ even} \\ -\frac{n+1}{2} & \text{if } n \text{ odd} \end{cases}$$

the set \mathbb{Q} of rational numbers

$$\mathbb{Q} = \left\{ \frac{p}{q} : \begin{array}{l} p \in \mathbb{Z} \\ q \in \mathbb{N}, q \neq 0 \end{array} \right\}$$



We can't do the same thing we did for the integers.

Between any two rational numbers there are an infinite number of others.

the set of positive rational numbers

A grid of positive rational numbers is shown, with rows labeled 0 through 7 on the left and columns labeled 1 through 8 on top. The numbers are arranged in a grid, with some cells containing ellipses. Handwritten annotations include red and blue numbers and green diagonal lines.

0	1/1	1/2	1/3	1/4	1/5	1/6	1/7	1/8	...
2	2/1	2/2	2/3	2/4	2/5	2/6	2/7	2/8	...
3	3/1	3/2	3/3	3/4	3/5	3/6	3/7	3/8	...
4	4/1	4/2	4/3	4/4	4/5	4/6	4/7	4/8	...
5	5/1	5/2	5/3	5/4	5/5	5/6	5/7
6	6/1	6/2	6/3	6/4	6/5	6/6
7	7/1	7/2	7/3	7/4	7/5
...

Handwritten annotations include red numbers (0, 2, 3, 8, 10, 9) and blue numbers (0, 0, 1, 2, 3, 4, 5, 6, 7, 8, 9) placed above or below the grid. Green diagonal lines connect the top-left to the bottom-right, passing through the main diagonal.

the set of positive rational numbers

The set of all positive rational numbers is **countable**.

$$\mathbb{Q}^+ = \{1/1, 2/1, 1/2, 3/1, \del{2/2}, \del{1/3}, 4/1, 2/3, 3/2, 1/4, 5/1, \del{4/2}, \del{3/3}, 2/4, 1/5, \dots\}$$

List elements in order of numerator+denominator, breaking ties according to denominator.

Only k numbers have total of sum of $k + 1$, so every positive rational number comes up some point.

Technique is called “dovetailing.”

Notice that repeats are OK because we can skip over them.

Formal statement about “skipping”:

A set S is **countable** iff S is infinite and there is an onto map $g : \mathbb{N} \rightarrow S$.

the set \mathbb{Q} of rational numbers

Claim: Σ^* is ^{at most} countable for every finite Σ

$$\Sigma = \{a, b, c\}$$

$$\Sigma^* = \{\epsilon, a, b, c, \\ aa, ab, ac, \\ \dots\}$$

ϵ

X_1

$$X_1 \in \Sigma$$

$X_1 X_2$

$$X_1, X_2 \in \Sigma$$

$X_1 X_2 X_3$

$$X_1, X_2, X_3 \in \Sigma$$

?

0 1 2 3 4 5 ...
 $\epsilon, a, b, c, aa, ab, ac, ba, bb, bc, ca, cb, cc, \\ aaa, aab, \dots$

the set of all Java programs is countable

$$\text{Java} \subseteq \Sigma^*$$

for some Σ

$$g: \mathbb{N} \rightarrow \Sigma^* \quad \text{1-1, onto}$$

exists by prev. slide

want: $f: \mathbb{N} \rightarrow \text{Java}$ onto

$$f(n) = \begin{cases} g(n) & \text{if } g(n) \in \text{Java} \\ P_0 & \text{otherwise} \end{cases}$$

$P_0 =$ fixed Java program

ok ok, everything is countable except your mom

“Your mamma so fat she couldn’t be put into one to one correspondence with the natural numbers.”

Burn.

are the real numbers countable?

Theorem [Cantor]:

The set of real numbers between 0 and 1 is **not** countable.

Proof will be by contradiction. Uses a new method called diagonalization.

proof that $[0,1)$ is uncountable

Suppose, for the sake of contradiction, that there is a list of them:

		1	2	3	4	5	6	7	8	9	...
r_1	0.	5	0	0	0	0	0	0	0
r_2	0.	3	3	3	3	3	3	3	3
r_3	0.	1	4	2	8	5	7	1	4
r_4	0.	1	4	1	5	9	2	6	5
r_5	0.	1	2	1	2	2	1	2	2
r_6	0.	2	5	0	0	0	0	0	0
r_7	0.	7	1	8	2	8	1	8	2
r_8	0.	6	1	8	0	3	3	9	4
...

proof that $[0,1)$ is uncountable

Suppose, for the sake of contradiction, that there is a list of them:

		1	2	3	4	5	6	7	8	9	...
r_1	0.	5	0	0	0	0	0	0	0
r_2	0.	3	3	3	3	3	3	3	3
r_3	0.	1	4	2	8	5	7	1	4
r_4	0.	1	4	1	5	9	2	6	5
r_5	0.	1	2	1	2	2	1	2	2
r_6	0.	2	5	0	0	0	0	0	0
r_7	0.	7	1	8	2	8	1	8	2
r_8	0.	6	1	8	0	3	3	9	4
...

proof that $[0,1)$ is uncountable

Suppose, for the sake of contradiction, that there is a list of them:

		1	2	3	4						
r_1	0.	5	0	0	0						
r_2	0.	3	3	3	3						
r_3	0.	1	4	2	8	5	7	1	4
r_4	0.	1	4	1	5	9	2	6	5
r_5	0.	1	2	1	2	2	1	2	2
r_6	0.	2	5	0	0	0	0	0	0
r_7	0.	7	1	8	2	8	1	8	2
r_8	0.	6	1	8	0	3	3	9	4
...

Flipping rule:
Only if the other driver deserves it.

proof that $[0,1)$ is uncountable

Suppose, for the sake of contradiction, that there is a list of them:

		1	2	3	4						
r_1	0.	5 ¹	0	0	0						
r_2	0.	3	3 ⁵	3	3						
r_3	0.	1	4	2 ⁵	8	5	7	1	4
r_4	0.	1	4	1	5 ¹	9	2	6	5
r_5	0.	1	2	1	2	2 ⁵	1	2	2
r_6	0.	2	5	0	0	0	0 ⁵	0	0
r_7	0.	7	1	8	2	8	1	8 ⁵	2
r_8	0.	6	1	8	0	3	3	9	4 ⁵
...

Flipping rule:

If digit is **5**, make it **1**.

If digit is not **5**, make it **5**.

proof that $[0,1)$ is uncountable

Suppose, for the sake of contradiction, that there is a list of them:

		1	2	3	4						
r_1	0.	5 ¹	0	0	0						
r_2	0.	3	3 ⁵	3	3						
r_3	0.	1	4	2 ⁵	8	5	7	1	4
r_4	0.	1	4	1	5 ¹	9	2	6	5
r_5	0.	1	2	1	2	2 ⁵	1	2	2
r_6	0.	2	5	0	0	0	0 ⁵	0	0
r_7	0.	7	1	8	2	8	1	8 ⁵	2

Flipping rule:

If digit is **5**, make it **1**.

If digit is not **5**, make it **5**.

If diagonal element is $0.x_{11}x_{22}x_{33}x_{44}x_{55}\dots$ then let's call the flipped number $0.\hat{x}_{11}\hat{x}_{22}\hat{x}_{33}\hat{x}_{44}\hat{x}_{55}\dots$ **It cannot appear anywhere on the list!**

proof that $[0,1)$ is uncountable

Suppose, for the sake of contradiction, that there is a list of them:

		1	2	3	4
r_1	0.	5 ¹	0	0	0
r_2	0.	3	3 ⁵	3	3
r_3	0.	1	4	2 ⁵	8
r_4	0.	1	4	1	5 ¹

Flipping rule:

If digit is **5**, make it **1**.

If digit is not **5**, make it **5**.

5	7	1	4
9	2	6	5
2 ⁵	1	2	2
0	0 ⁵	0	0
8	1	8 ⁵	2

For every $n \geq 1$:

$$r_n \neq 0.\hat{x}_{11}\hat{x}_{22}\hat{x}_{33}\hat{x}_{44}\hat{x}_{55} \dots$$

because the numbers differ on the n th digit!

If diagonal element is $0.x_{11}x_{22}x_{33}x_{44}x_{55} \dots$ then let's call it the flipped number $0.\hat{x}_{11}\hat{x}_{22}\hat{x}_{33}\hat{x}_{44}\hat{x}_{55} \dots$ **It cannot appear anywhere on the list!**

proof that $[0,1)$ is uncountable

Suppose, for the sake of contradiction, that there is a list of them:

		1	2	3	4
r_1	0.	5 ¹	0	0	0
r_2	0.	3	3 ⁵	3	3
r_3	0.	1	4	2 ⁵	8
r_4	0.	1	4	1	5 ¹

Flipping rule:

If digit is **5**, make it **1**.

If digit is not **5**, make it **5**.

5	7	1	4
9	2	6	5
2 ⁵	1	2	2
0	0 ⁵	0	0
8	1	8 ⁵	2

For every $n \geq 1$:

$$r_n \neq 0.\hat{x}_{11}\hat{x}_{22}\hat{x}_{33}\hat{x}_{44}\hat{x}_{55} \dots$$

because the numbers differ on the n th digit!

So the list is incomplete, which is a contradiction.

Thus the real numbers between 0 and 1 are **uncountable**.

the set of all functions $f : \mathbb{N} \rightarrow \{0, \dots, 9\}$ is uncountable

uncomputable functions

We have seen that:

- The set of all (Java) programs is countable
- The set of all functions $f : \mathbb{N} \rightarrow \{0, \dots, 9\}$ is not countable

So: There must be some function $f : \mathbb{N} \rightarrow \{0, \dots, 9\}$ that is not computable by any program!

Interesting... maybe.

Can we come up with an explicit function that is uncomputable?

(Next time)