




Spring 2015

Lecture 26: Relations and directed graphs



Let A and B be sets.

A **binary relation** from A to B is a subset of  $A \times B$ 

Let A be a set.

A **binary relation on A** is a subset of  $A \times A$ 

## relations you already know!

 $\geq$  on  $\mathbb{N}$ That is:  $\{(x,y) : x \geq y \text{ and } x, y \in \mathbb{N}\}$  $<$  on  $\mathbb{R}$ That is:  $\{(x,y) : x < y \text{ and } x, y \in \mathbb{R}\}$  $=$  on  $\Sigma^*$ That is:  $\{(x,y) : x = y \text{ and } x, y \in \Sigma^*\}$  $\subseteq$  on  $P(U)$  for universe UThat is:  $\{(A,B) : A \subseteq B \text{ and } A, B \in P(U)\}$ 

## properties of relations

Let R be a relation on A.

R is **reflexive** iff  $(a,a) \in R$  for every  $a \in A$ R is **symmetric** iff  $(a,b) \in R$  implies  $(b,a) \in R$ R is **antisymmetric** iff  $(a,b) \in R$  and  $a \neq b$  implies  $(b,a) \notin R$ R is **transitive** iff  $(a,b) \in R$  and  $(b,c) \in R$  implies  $(a,c) \in R$ 

## examples

 $R_1 = \{(a, 1), (a, 2), (b, 1), (b, 3), (c, 3)\}$  $R_2 = \{(x, y) \mid x \equiv y \pmod{5}\}$  $R_3 = \{(c_1, c_2) \mid c_1 \text{ is a prerequisite of } c_2\}$  $R_4 = \{(s, c) \mid \text{student } s \text{ had taken course } c\}$ 

## combining relations

Let R be a relation from A to B.

Let S be a relation from B to C.

The **composition** of R and S,  $S \circ R$  is the relation from A to C defined by: $S \circ R = \{(a, c) \mid \exists b \text{ such that } (a,b) \in R \text{ and } (b,c) \in S\}$ 

Intuitively, a pair is in the composition if there is a "connection" from the first to the second.

## examples

$(a,b) \in \text{Parent}$  iff  $b$  is a parent of  $a$

$(a,b) \in \text{Sister}$  iff  $b$  is a sister of  $a$

When is  $(x,y) \in \text{Sister} \circ \text{Parent}$ ?

When is  $(x,y) \in \text{Parent} \circ \text{Sister}$ ?

$$S \circ R = \{(a, c) \mid \exists b \text{ such that } (a,b) \in R \text{ and } (b,c) \in S\}$$

## examples

Using the relations: Parent, Child, Brother, Sister, Sibling, Father, Mother, Husband, Wife express:

Uncle:  $b$  is an uncle of  $a$

Uncle = Brother  $\circ$  Parent

Cousin:  $b$  is a cousin of  $a$

Cousin = Child  $\circ$  Sibling  $\circ$  Parent

## powers of a relation

$$R^2 = R \circ R \\ = \{(a, c) \mid \exists b \text{ such that } (a, b) \in R \text{ and } (b, c) \in R\}$$

Parent<sup>2</sup> = GrandParent

$$R^0 = \{(a, a) \mid a \in A\} \quad R^0 \text{ is always equality}$$

$$R^1 = R$$

$$R^{n+1} = R^n \circ R$$

## examples

Using the relations: Parent, Child, Brother, Sister, Sibling, Father, Mother, Husband, Wife express:

Uncle:  $b$  is an uncle of  $a$

Cousin:  $b$  is a cousin of  $a$

## powers of a relation

$$R^2 = R \circ R \\ = \{(a, c) \mid \exists b \text{ such that } (a, b) \in R \text{ and } (b, c) \in R\}$$

$$R^0 = \{(a, a) \mid a \in A\}$$

$$R^1 = R$$

$$R^{n+1} = R^n \circ R$$

## matrix representation

Relation  $R$  on  $A = \{a_1, \dots, a_p\}$

$$m_{ij} = \begin{cases} 1 & \text{if } (a_i, a_j) \in R, \\ 0 & \text{if } (a_i, a_j) \notin R. \end{cases}$$

$\{(1, 1), (1, 2), (1, 4), (2, 1), (2, 3), (3, 2), (3, 3), (4, 2), (4, 3)\}$


|   | 1 | 2 | 3 | 4 |
|---|---|---|---|---|
| 1 | 1 | 1 | 0 | 1 |
| 2 | 1 | 0 | 1 | 0 |
| 3 | 0 | 1 | 1 | 0 |
| 4 | 0 | 1 | 1 | 0 |

directed graphs

$G = (V, E)$        $V$  - vertices  
                           $E$  - edges, ordered pairs of vertices

Path:  $v_0, v_1, \dots, v_k$  with  $(v_i, v_{i+1})$  in  $E$


Simple Path  
 Cycle  
 Simple Cycle



representation of relations

Directed Graph Representation (Digraph)


$\{(a, b), (a, a), (b, a), (c, a), (c, d), (c, e), (d, e)\}$



representation of relations

Directed Graph Representation (Digraph)


$\{(a, b), (a, a), (b, a), (c, a), (c, d), (c, e), (d, e)\}$



representation of relations

Directed Graph Representation (Digraph)

$\{(a, b), (a, a), (b, a), (c, a), (c, d), (c, e), (d, e)\}$



relational composition using digraphs

If  $S = \{(2,2), (2,3), (3,1)\}$  and  $R = \{(1,2), (2,1), (1,3)\}$   
 Compute  $S \circ R$

relational composition using digraphs

If  $S = \{(2,2), (2,3), (3,1)\}$  and  $R = \{(1,2), (2,1), (1,3)\}$   
 Compute  $S \circ R$

### relational composition using digraphs

If  $S = \{(2,2), (2,3), (3,1)\}$  and  $R = \{(1,2), (2,1), (1,3)\}$   
 Compute  $S \circ R$

### paths in relations and graphs

A **path** in a graph of length  $n$  is a list of edges with vertices next to each other.

Let  $R$  be a relation on a set  $A$ . There is a path of length  $n$  from  $a$  to  $b$  if and only if  $(a,b) \in R^n$

### connectivity in graphs

Two vertices in a graph are **connected** iff there is a path between them.

Let  $R$  be a relation on a set  $A$ . The connectivity relation  $R^*$  consists of the pairs  $(a,b)$  such that there is a path from  $a$  to  $b$  in  $R$ .

$$R^* = \bigcup_{k=0}^{\infty} R^k$$

Note: The book uses the wrong definition of this quantity. What the text defines (ignoring  $k=0$ ) is usually called  $R^+$

### properties of relations (repeated)


Let  $R$  be a relation on  $A$ .

$R$  is **reflexive** iff  $(a,a) \in R$  for every  $a \in A$

$R$  is **symmetric** iff  $(a,b) \in R$  implies  $(b,a) \in R$

$R$  is **transitive** iff  $(a,b) \in R$  and  $(b,c) \in R$  implies  $(a,c) \in R$


### transitive-reflexive closure



Add the minimum possible number of edges to make the relation transitive and reflexive.

The transitive-reflexive closure of a relation  $R$  is the connectivity relation  $R^*$


### transitive-reflexive closure



Add the minimum possible number of edges to make the relation transitive and reflexive.

The transitive-reflexive closure of a relation  $R$  is the connectivity relation  $R^*$

## transitive-reflexive closure



Add the minimum possible number of edges to make the relation transitive and reflexive.

The transitive-reflexive closure of a relation R is the connectivity relation  $R^*$

## n-ary relations

Let  $A_1, A_2, \dots, A_n$  be sets. An **n-ary** relation on these sets is a subset of  $A_1 \times A_2 \times \dots \times A_n$ .

## relational databases

STUDENT

| Student_Name | ID_Number | Office | GPA  |
|--------------|-----------|--------|------|
| Knuth        | 328012098 | 022    | 4.00 |
| Von Neuman   | 481080220 | 555    | 3.78 |
| Russell      | 238082388 | 022    | 3.85 |
| Einstein     | 238001920 | 022    | 2.11 |
| Newton       | 1727017   | 333    | 3.61 |
| Karp         | 348882811 | 022    | 3.98 |
| Bernoulli    | 2921938   | 022    | 3.21 |

## relational databases

STUDENT

| Student_Name | ID_Number | Office | GPA  | Course |
|--------------|-----------|--------|------|--------|
| Knuth        | 328012098 | 022    | 4.00 | CSE311 |
| Knuth        | 328012098 | 022    | 4.00 | CSE351 |
| Von Neuman   | 481080220 | 555    | 3.78 | CSE311 |
| Russell      | 238082388 | 022    | 3.85 | CSE312 |
| Russell      | 238082388 | 022    | 3.85 | CSE344 |
| Russell      | 238082388 | 022    | 3.85 | CSE351 |
| Newton       | 1727017   | 333    | 3.61 | CSE312 |
| Karp         | 348882811 | 022    | 3.98 | CSE311 |
| Karp         | 348882811 | 022    | 3.98 | CSE312 |
| Karp         | 348882811 | 022    | 3.98 | CSE344 |
| Karp         | 348882811 | 022    | 3.98 | CSE351 |
| Bernoulli    | 2921938   | 022    | 3.21 | CSE351 |

What's not so nice?

## relational databases

STUDENT

| Student_Name | ID_Number | Office | GPA  |
|--------------|-----------|--------|------|
| Knuth        | 328012098 | 022    | 4.00 |
| Von Neuman   | 481080220 | 555    | 3.78 |
| Russell      | 238082388 | 022    | 3.85 |
| Einstein     | 238001920 | 022    | 2.11 |
| Newton       | 1727017   | 333    | 3.61 |
| Karp         | 348882811 | 022    | 3.98 |
| Bernoulli    | 2921938   | 022    | 3.21 |

TAKES

| ID_Number | Course |
|-----------|--------|
| 328012098 | CSE311 |
| 328012098 | CSE351 |
| 481080220 | CSE311 |
| 238082388 | CSE312 |
| 238082388 | CSE344 |
| 238082388 | CSE351 |
| 1727017   | CSE312 |
| 348882811 | CSE311 |
| 348882811 | CSE312 |
| 348882811 | CSE344 |
| 348882811 | CSE351 |
| 2921938   | CSE351 |

Better

## database operations: projection

Find all offices:  $\Pi_{\text{Office}}(\text{STUDENT})$

| Office |
|--------|
| 022    |
| 555    |
| 333    |

Find offices and GPAs:  $\Pi_{\text{Office}, \text{GPA}}(\text{STUDENT})$

| Office | GPA  |
|--------|------|
| 022    | 4.00 |
| 555    | 3.78 |
| 022    | 3.85 |
| 022    | 2.11 |
| 333    | 3.61 |
| 022    | 3.98 |
| 022    | 3.21 |

## database operations: selection

Find students with GPA > 3.9 :  $\sigma_{\text{GPA}>3.9}(\text{STUDENT})$

| Student_Name | ID_Number | Office | GPA  |
|--------------|-----------|--------|------|
| Knuth        | 328012098 | 022    | 4.00 |
| Karp         | 348882811 | 022    | 3.98 |

Retrieve the name and GPA for students with GPA > 3.9:

$\Pi_{\text{Student\_Name,GPA}}(\sigma_{\text{GPA}>3.9}(\text{STUDENT}))$

| Student_Name | GPA  |
|--------------|------|
| Knuth        | 4.00 |
| Karp         | 3.98 |

## database operations: natural join

Student  $\bowtie$  Takes

| Student_Name | ID_Number | Office | GPA  | Course |
|--------------|-----------|--------|------|--------|
| Knuth        | 328012098 | 022    | 4.00 | CSE311 |
| Knuth        | 328012098 | 022    | 4.00 | CSE351 |
| Von Neuman   | 481080220 | 555    | 3.78 | CSE311 |
| Russell      | 238082388 | 022    | 3.85 | CSE312 |
| Russell      | 238082388 | 022    | 3.85 | CSE344 |
| Russell      | 238082388 | 022    | 3.85 | CSE351 |
| Newton       | 1727017   | 333    | 3.61 | CSE312 |
| Karp         | 348882811 | 022    | 3.98 | CSE311 |
| Karp         | 348882811 | 022    | 3.98 | CSE312 |
| Karp         | 348882811 | 022    | 3.98 | CSE344 |
| Karp         | 348882811 | 022    | 3.98 | CSE351 |
| Bernoulli    | 2921938   | 022    | 3.21 | CSE351 |