


# CSE 311: Foundations of Computing

## Lecture 2: More Logic, Equivalence & Digital Circuits

shayan



# Last class: Some Connectives & Truth Tables

---

Negation (not)

$p$	$\neg p$
T	F
F	T

Conjunction (and)

$p$	$q$	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F

Disjunction (or)

$p$	$q$	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F

Exclusive Or

$p$	$q$	$p \oplus q$
T	T	F
T	F	T
F	T	T
F	F	F

# Last Class: $p \rightarrow q$

## Implication:

- $p$  implies  $q$
- whenever  $p$  is true  $q$  must be true
- if  $p$  then  $q$
- $q$  if  $p$
- $p$  is sufficient for  $q$
- $p$  only if  $q$
- $q$  is necessary for  $p$

if  $p$  then  $q$   
Fals  $p = T$  and  $q = F$

$p$	$q$	$p \rightarrow q$
T	T	T
T	F	F
F	T	T
F	F	T

$p$  only if  $q$   
I play basketball only if  
you do as well.  
I play      you play

F	T	T
F	F	T
T	T	T
T	F	F

# Biconditional: $p \leftrightarrow q$

---

- $p$  iff  $q$
- $p$  is equivalent to  $q$
- $p$  implies  $q$  and  $q$  implies  $p$
- $p$  is necessary and sufficient for  $q$

$p$	$q$	$p \leftrightarrow q$
T	T	T
T	F	F
F	T	F
F	F	T

/

-

# Biconditional: $p \leftrightarrow q$

---

- $p$  iff  $q$
- $p$  is equivalent to  $q$
- $p$  implies  $q$  and  $q$  implies  $p$
- $p$  is necessary and sufficient for  $q$

$p$	$q$	$p \leftrightarrow q$
T	T	T
T	F	F
F	T	F
F	F	T

# Last class: Using Logical Connectives

---

Measles:

“You can get measles”

Mumps:

“You can get mumps”

MMR:

“You had the MMR vaccine”

“You can get measles and mumps if you didn’t have the MMR vaccine, but if you had the MMR vaccine then you can’t get either.”



$((\text{Measles and Mumps}) \text{ if not MMR}) \text{ and } (\text{if MMR then not } (\text{Measles or Mumps}))$



$((\text{Measles} \wedge \text{Mumps}) \text{ if } \neg \text{MMR}) \wedge (\text{if MMR then } \neg (\text{Measles} \vee \text{Mumps}))$   
 $(\neg \text{MMR} \rightarrow (\text{Measles} \wedge \text{Mumps})) \wedge (\text{MMR} \rightarrow \neg (\text{Measles} \vee \text{Mumps}))$

# Understanding the Vaccine Sentence

---

**“You can get measles and mumps if you didn’t have the MMR vaccine, but if you had the MMR vaccine you can’t get either.”**



$((\text{Measles} \wedge \text{Mumps}) \text{ if } \neg \text{MMR}) \wedge (\text{if MMR then } \neg(\text{Measles} \vee \text{Mumps}))$



$(\neg \text{MMR} \rightarrow (\text{Measles} \wedge \text{Mumps})) \wedge (\text{MMR} \rightarrow \neg(\text{Measles} \vee \text{Mumps}))$

# Understanding the Vaccine Sentence

---

“You can get measles and mumps if you didn’t have the MMR vaccine, but if you had the MMR vaccine you can’t get either.”



$((\text{Measles} \wedge \text{Mumps}) \text{ if } \neg \text{MMR}) \wedge (\text{if MMR then } \neg(\text{Measles} \vee \text{Mumps}))$



$(\neg \text{MMR} \rightarrow (\text{Measles} \wedge \text{Mumps})) \wedge (\text{MMR} \rightarrow \neg(\text{Measles} \vee \text{Mumps}))$

Define shorthand ...

$p : \text{MMR}$

$q : \text{Measles}$


$r : \text{Mumps}$



$(\neg p \rightarrow (q \wedge r)) \wedge (p \rightarrow \neg(q \vee r))$



# Analyzing the Vaccine Sentence with a Truth Table



$p$	$q$	$r$	$\neg p$	$q \wedge r$	$\neg p \rightarrow (q \wedge r)$	$q \vee r$	$\neg(q \vee r)$	$p \rightarrow \neg(q \vee r)$	$(\neg p \rightarrow (q \wedge r)) \wedge (p \rightarrow \neg(q \vee r))$
T	T	T	F	T	T	T	F	F	F
T	T	F	F	F	T	T	F	F	F
T	F	T	F	F	T	T	F	F	F
T	F	F	F	F	T	F	T	T	T
F	T	T	T	T	T	T	F	T	T
F	T	F	T	F	F	T	F	T	F
F	F	T	T	F	F	T	F	T	F
F	F	F	T	F	F	F	T	T	F

# Analyzing the Vaccine Sentence with a Truth Table

---

$p$	$q$	$r$	$\neg p$	$q \wedge r$	$\neg p \rightarrow (q \wedge r)$	$q \vee r$	$\neg(q \vee r)$	$p \rightarrow \neg(q \vee r)$	$(\neg p \rightarrow (q \wedge r)) \wedge (p \rightarrow \neg(q \vee r))$
T	T	T	F	T	T	T	F	F	F
T	T	F	F	F	T	T	F	F	F
T	F	T	F	F	T	T	F	F	F
T	F	F	F	F	T	F	T	T	T
F	T	T	T	T	T	T	F	T	T
F	T	F	T	F	F	T	F	T	F
F	F	T	T	F	F	T	F	T	F
F	F	F	T	F	F	F	T	T	F

# Converse, Contrapositive

---

Implication:

$$p \rightarrow q$$

if I take  
umbrella  
then it rains

Converse:

$$q \rightarrow p$$

If it rains  
I take my umbrella

Contrapositive:

$$\neg q \rightarrow \neg p$$

If I don't  
rain  
then I don't take  
umbrella

Inverse:

$$\neg p \rightarrow \neg q$$

Consider

$p$ :  $x$  is divisible by 2

$q$ :  $x$  is divisible by 4

$p \rightarrow q$	
$q \rightarrow p$	
$\neg q \rightarrow \neg p$	
$\neg p \rightarrow \neg q$	

# Converse, Contrapositive

---

Implication:

$$p \rightarrow q$$

Converse:

$$q \rightarrow p$$

Contrapositive:

$$\neg q \rightarrow \neg p$$

Inverse:

$$\neg p \rightarrow \neg q$$

Consider

$p$ :  $x$  is divisible by 2

$q$ :  $x$  is divisible by 4

$p \rightarrow q$	$\times$
$q \rightarrow p$	$\checkmark$
$\neg q \rightarrow \neg p$	$\times$
$\neg p \rightarrow \neg q$	$\checkmark$

$q$   
 $\neg q$

	Divisible By 2	Not Divisible By 2
Divisible By 4	4, 8	Imposs
Not Divisible By 4	2, 6	1, 3, 5

$p$        $\neg p$

# Converse, Contrapositive

---

Implication:

$$p \rightarrow q$$

Converse:

$$q \rightarrow p$$

Contrapositive:

$$\neg q \rightarrow \neg p$$

Inverse:

$$\neg p \rightarrow \neg q$$

Consider

$p$ :  $x$  is divisible by 2

$q$ :  $x$  is divisible by 4

$p \rightarrow q$	
$q \rightarrow p$	
$\neg q \rightarrow \neg p$	
$\neg p \rightarrow \neg q$	

	Divisible By 2	Not Divisible By 2
Divisible By 4	4,8,12,...	Impossible
Not Divisible By 4	2,6,10,...	1,3,5,...

# Converse, Contrapositive

---

Implication:

$$p \rightarrow q$$

Contrapositive:

$$\neg q \rightarrow \neg p$$


Converse:

$$q \rightarrow p$$

Inverse:

$$\neg p \rightarrow \neg q$$

How do these relate to each other?



$p$	$q$	$p \rightarrow q$	$q \rightarrow p$	$\neg p$	$\neg q$	$\neg p \rightarrow \neg q$	$\neg q \rightarrow \neg p$
T	T	T	T	F	F	T	T
T	F	F	T	F	T	T	F
F	T	T	F	T	F	F	T
F	F	T	T	T	T	T	T

# Converse, Contrapositive

---

Implication:

$$p \rightarrow q$$

Converse:

$$q \rightarrow p$$

Contrapositive:

$$\neg q \rightarrow \neg p$$

Inverse:

$$\neg p \rightarrow \neg q$$

An **implication** and its **contrapositive**  
have the same truth value!

$p$	$q$	$p \rightarrow q$	$q \rightarrow p$	$\neg p$	$\neg q$	$\neg p \rightarrow \neg q$	$\neg q \rightarrow \neg p$
T	T	T	T	F	F	T	T
T	F	F	T	F	T	T	F
F	T	T	F	T	F	F	T
F	F	T	T	T	T	T	T

# Tautologies!

---

**Terminology:** A compound proposition is a...

- *Tautology* if it is always true
- *Contradiction* if it is always false
- *Contingency* if it can be either true or false

$$p \vee \neg p$$

*Tautology*

$$p \oplus p$$

*Contrad.*

$$(p \rightarrow q) \wedge p$$

*Conting.*

$$\begin{array}{l} p = F \rightarrow F \\ p = T \quad q = T \rightarrow T \end{array}$$



# Tautologies!

---

**Terminology:** A compound proposition is a...

- *Tautology* if it is always true
- *Contradiction* if it is always false
- *Contingency* if it can be either true or false

$$p \vee \neg p$$

This is a tautology. It's called the "law of the excluded middle. If  $p$  is true, then  $p \vee \neg p$  is true. If  $p$  is false, then  $p \vee \neg p$  is true.

$$p \oplus p$$

This is a contradiction. It's always false no matter what truth value  $p$  takes on.

$$(p \rightarrow q) \wedge p$$

This is a contingency. When  $p=T, q=T$ ,  $(T \rightarrow T) \wedge T$  is true.

When  $p=T, q=F$ ,  $(T \rightarrow F) \wedge T$  is false.

# Logical Equivalence

---

**A = B** means **A** and **B** are identical “strings”:

$$- p \wedge q = p \wedge q$$

$$- p \wedge q \neq q \wedge p$$

# Logical Equivalence

---

**A = B** means **A** and **B** are identical “strings”:

–  $p \wedge q = p \wedge q$

These are equal, because they are character-for-character identical.

–  $p \wedge q \neq q \wedge p$

These are NOT equal, because they are different sequences of characters. They “mean” the same thing though.

**A ≡ B** means **A** and **B** have identical truth values:

–  $p \wedge q \equiv p \wedge q$  ✓

–  $p \wedge q \equiv q \wedge p$  ✓

–  $p \wedge q \neq q \vee p$

$p = T, q = F \rightarrow$   
 $\underbrace{\quad}_F \quad \underbrace{\quad}_T$

≡

# Logical Equivalence

---

**A = B** means **A** and **B** are identical “strings”:

- $p \wedge q = p \wedge q$

These are equal, because they are character-for-character identical.

- $p \wedge q \neq q \wedge p$

These are NOT equal, because they are different sequences of characters. They “mean” the same thing though.

**A ≡ B** means **A** and **B** have identical truth values:

- $p \wedge q \equiv p \wedge q$

Two formulas that are equal also are equivalent.

- $p \wedge q \equiv q \wedge p$

These two formulas have the same truth table!

- $p \wedge q \neq q \vee p$

When  $p=T$  and  $q=F$ ,  $p \wedge q$  is false, but  $p \vee q$  is true!

$A \leftrightarrow B$  vs.  $A \equiv B$

---

$p \leftrightarrow q$

~~$p \equiv q$~~

$A \equiv B$  is an **assertion** over all possible truth values that  $A$  and  $B$  always have the same truth values.

$A \leftrightarrow B$  is a **proposition** that may be true or false depending on the truth values of the variables in  $A$  and  $B$ .

$A \equiv B$  and  $(A \leftrightarrow B) \equiv T$  have the same meaning.

# De Morgan's Laws

---

$$\neg(p \wedge q) \equiv \neg p \vee \neg q$$

$$\neg(p \vee q) \equiv \neg p \wedge \neg q$$

Negate the statement:

“My code compiles or there is a bug.”

To negate the statement, ask “when is the original statement false”.

My code compiles  $\vee$  there is a bug  
 $\neg$  My code compiles  $\wedge$   $\neg$  there is a bug  
My code doesn't compile and there is not a bug.

# De Morgan's Laws

---

$$\neg(p \wedge q) \equiv \neg p \vee \neg q$$

$$\neg(p \vee q) \equiv \neg p \wedge \neg q$$

Negate the statement:

“My code compiles or there is a bug.”

To negate the statement, ask “when is the original statement false”.

It's false when not(my code compiles) AND not(there is a bug).

Translating back into English, we get:

My code doesn't compile and there is not a bug.

# De Morgan's Laws

---

Example:  $\neg(p \wedge q) \equiv (\neg p \vee \neg q)$

$p$	$q$	$\neg p$	$\neg q$	$\neg p \vee \neg q$	$p \wedge q$	$\neg(p \wedge q)$	$\neg(p \wedge q) \leftrightarrow (\neg p \vee \neg q)$
T	T	F	F	F	T	F	T
T	F	F	T	T	F	T	T
F	T	T	F	T	F	T	T
F	F	T	T	T	F	T	T

-

.

-



# De Morgan's Laws

---

Example:  $\neg(p \wedge q) \equiv (\neg p \vee \neg q)$

$p$	$q$	$\neg p$	$\neg q$	$\neg p \vee \neg q$	$p \wedge q$	$\neg(p \wedge q)$	$\neg(p \wedge q) \leftrightarrow (\neg p \vee \neg q)$
T	T	F	F	F	T	F	T
T	F	F	T	T	F	T	T
F	T	T	F	T	F	T	T
F	F	T	T	T	F	T	T

# De Morgan's Laws

---

$$\neg(p \wedge q) \equiv \neg p \vee \neg q$$

$$\neg(p \vee q) \equiv \neg p \wedge \neg q$$

```
if (!(front != null && value > front.data))  
    front = new ListNode(value, front);  
else {  
    ListNode current = front;  
    while (current.next != null && current.next.data < value))  
        current = current.next;  
    current.next = new ListNode(value, current.next);  
}
```

$front == null \quad !! \quad value \leq front.data$

# De Morgan's Laws

---

$$\neg(p \wedge q) \equiv \neg p \vee \neg q$$

$$\neg(p \vee q) \equiv \neg p \wedge \neg q$$

`!(front != null && value > front.data)`

$\equiv$

`front == null || value <= front.data`

**You've been using these for a while!**

# Law of Implication

---

$$p = T \quad q = F$$

$$\neg p = F \quad q = F$$
$$p \rightarrow q \equiv \neg p \vee q$$

$p$	$q$	$p \rightarrow q$	$\neg p$	$\neg p \vee q$	$p \rightarrow q \leftrightarrow \neg p \vee q$
T	T	T	F	T	T
T	F	F	F	F	T
F	T	T	T	T	T
F	F	T	T	T	T

# Law of Implication

---

$$p \rightarrow q \equiv \neg p \vee q$$

$p$	$q$	$p \rightarrow q$	$\neg p$	$\neg p \vee q$	$p \rightarrow q \leftrightarrow \neg p \vee q$
T	T	T	F	T	T
T	F	F	F	F	T
F	T	T	T	T	T
F	F	T	T	T	T

# Some Equivalences Related to Implication

---

$$p \rightarrow q \quad \equiv \quad \neg p \vee q$$

$$p \rightarrow q \quad \equiv \quad \neg q \rightarrow \neg p$$

$$p \leftrightarrow q \quad \equiv \quad (p \rightarrow q) \wedge (q \rightarrow p)$$

$$p \leftrightarrow q \quad \equiv \quad \neg p \leftrightarrow \neg q$$

# Properties of Logical Connectives

---

We will always give  
you this list!

- **Identity**

- $p \wedge T \equiv p$
- $p \vee F \equiv p$

- **Domination**

- $p \vee T \equiv T$
- $p \wedge F \equiv F$

- **Idempotent**

- $p \vee p \equiv p$
- $p \wedge p \equiv p$


- **Commutative**

- $p \vee q \equiv q \vee p$
- $p \wedge q \equiv q \wedge p$

- **Associative**

- $(p \vee q) \vee r \equiv p \vee (q \vee r)$
- $(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$

- **Distributive**

- 
  - $p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$
  - $p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$

- **Absorption**

- $p \vee (p \wedge q) \equiv p$
- $p \wedge (p \vee q) \equiv p$

- **Negation**

- $p \vee \neg p \equiv T$
- $p \wedge \neg p \equiv F$

# Digital Circuits

---

## Computing With Logic

- **T** corresponds to **1** or “high” voltage
- **F** corresponds to **0** or “low” voltage

## Gates

- Take inputs and produce outputs (functions)
- Several kinds of gates
- Correspond to propositional connectives (most of them)



# And Gate

---

**AND Connective**

vs.


**AND Gate**

$p \wedge q$

$p$	$q$	$p \wedge q$
T	T	T
T	F	F
F	T	F
F	F	F



$p$	$q$	OUT
1	1	1
1	0	0
0	1	0
0	0	0



“block looks like D of AND”

# Or Gate

---


**OR Connective**

**vs.**



**OR Gate**

$p \vee q$

$p$	$q$	$p \vee q$
T	T	T
T	F	T
F	T	T
F	F	F



$p$	$q$	OUT
1	1	1
1	0	1
0	1	1
0	0	0



“arrowhead block looks like V”

# Not Gates

---


**NOT Connective**

vs.

**NOT Gate**


$\neg p$

$p$	$\neg p$
T	F
F	T



Also called  
*inverter*



$p$	OUT
1	0
0	1



# Blobs are Okay!


---

You may write gates using blobs instead of shapes!



# Combinational Logic Circuits


---



**Values get sent along wires connecting gates**

# Combinational Logic Circuits

---




Values get sent along wires connecting gates

$$\neg p \wedge (\neg q \wedge (r \vee s))$$

# Combinational Logic Circuits


---



**Wires can send one value to multiple gates!**

# Combinational Logic Circuits

---



**Wires can send one value to multiple gates!**

$$(p \wedge \neg q) \vee (\neg q \wedge r)$$



# Computing Equivalence

---

Describe an algorithm for computing if two logical expressions/circuits are equivalent.

**What is the run time of the algorithm?**

Compute the entire truth table for both of them!

There are  $2^n$  entries in the column for  $n$  variables.

# Some Familiar Properties of Arithmetic

---

- $x + y = y + x$  (Commutativity)
- $x \cdot (y + z) = x \cdot y + x \cdot z$  (Distributivity)
- $(x + y) + z = x + (y + z)$  (Associativity)

# Understanding Connectives

---

- **Reflect basic rules of reasoning and logic**
- **Allow manipulation of logical formulas**
  - Simplification
  - Testing for equivalence
- **Applications**
  - Query optimization
  - Search optimization and caching
  - Artificial Intelligence
  - Program verification