

# CSE 311: Foundations of Computing I

## Strong Induction Annotated Proofs

### Relevant Definitions

Fibonacci Numbers	DEFINITION
$f_n = \begin{cases} 0 & \text{if } n = 0 \\ 1 & \text{if } n = 1 \\ f_{n-1} + f_{n-2} & \text{if } n > 1 \end{cases}$	

### Bounding the Fibonacci Numbers

Prove that for all  $n \in \mathbb{N} \setminus \{0, 1\}$ ,  $2^{n/2-1} \leq f_n < 2^n$ .

#### Proof

Let  $P(n)$  be " $2^{n/2-1} \leq f_n < 2^n$ " for all  $n \in \mathbb{N} \setminus \{0, 1\}$ . We go by strong induction on  $n$ .

#### Base Case:

Note that  $2^{2/2-1} = 2^0 = 1 \leq 1 = 0 + 1 = f_0 + f_1 = f_2 < 4 = 2^2$ . So,  $P(2)$  is true.

#### Induction Hypothesis:

Suppose that  $P(2) \wedge P(3) \wedge \dots \wedge P(k)$  is true.

#### Induction Step:

We show that  $P(k + 1)$  is true.

Case  $((k + 1) - 2 < 2 \leftrightarrow k < 3 \leftrightarrow k = 2)$ :

Note that  $2^{3/2-1} = \frac{1}{2} \leq 2 = 1 + 1 = f_1 + f_2 = f_3 < 8 = 2^3$ . So,  $P(3)$  is true.

#### Commentary & Scratch Work

*We're using strong induction because it's a recurrence.*

*An alternative, is to introduce a variable to range over the hypotheses. This would look like "Suppose that  $P(\ell)$  is true for all  $2 \leq \ell \leq k$  for some  $k \in \mathbb{N} \setminus \{0, 1\}$ ."*

*In strong induction, the IS takes careful planning. Whenever we attempt to use the IH, we need to make sure we've actually assumed it. In particular, we must ask:*

- What is the smallest value that  $k$  could be? (Here it's 2)*
- If it's the smallest value, can we plug into the recurrence for  $k + 1$ ? ( $k + 1 = 2 + 1 = 3$ ,  $f_3 = f_2 + f_1$ )*
- If any of the values we'd need to plug in (here, 2 and 1) are less than our IH, (here, this is true for 1) we need special cases.*

*Again, this case happened, because we can't apply the IH to  $f_1$ .*

Case  $((k + 1) - 2 \geq 2 \leftrightarrow k \geq 3)$ :

Since  $k \geq 3$ , we know  $f_{k+1} = f_k + f_{k-1}$ .

Furthermore, we know that  $P(k)$  and  $P(k - 1)$  are both true by our IH. We take each piece of the claim independently.

Note that

$$\begin{aligned} f_{k+1} &= f_k + f_{k-1} \\ &\geq 2^{k/2-1} + 2^{(k-1)/2-1} \\ &\geq 2^{(k-1)/2-1} + 2^{(k-1)/2-1} \\ &= 2(2^{(k-1)/2-1}) \\ &= 2^{2/2+(k-1)/2-1} \\ &= 2^{(k+1)/2-1} \end{aligned}$$

Also, we have

$$\begin{aligned} f_{k+1} &= f_k + f_{k-1} \\ &\leq 2^k + 2^{k-1} \\ &\leq 2^k + 2^k \\ &\leq 2^{k+1} \end{aligned}$$

Since the claim is true for both cases,  
 $P(k) \rightarrow P(k + 1)$ .

So, the claim is true for all  $n \geq 2$  by induction on  $n$ .

*This is just a bunch of algebra. There's nothing special here other than the idea to just use the recurrence.*