



CSE 311: Foundations of Computing

Lecture 6: More Predicate Logic



Last class: Predicates

Predicate

- A function that returns a truth value, e.g.,

$\text{Cat}(x) ::= \text{"x is a cat"}$

$\text{Prime}(x) ::= \text{"x is prime"}$

$\text{HasTaken}(x, y) ::= \text{"student x has taken course y"}$

$\text{LessThan}(x, y) ::= \text{"x < y"}$

$\text{Sum}(x, y, z) ::= \text{"x + y = z"}$

$\text{GreaterThan5}(x) ::= \text{"x > 5"}$

$\text{HasNChars}(s, n) ::= \text{"string s has length n"}$

Predicates can have varying numbers of arguments and input types.

Last class: Domain of Discourse

For ease of use, we define one “type”/“domain” that we work over. This set of objects is called the “**domain of discourse**”.

For each of the following, what might the domain be?

(1) “x is a cat”, “x barks”, “x ruined my couch”

(2) “x is prime”, “ $x = 0$ ”, “ $x < 0$ ”, “x is a power of two”

(3) “x is a pre-req for z”

Domain of Discourse

For ease of use, we define one “type”/“domain” that we work over. This set of objects is called the “**domain of discourse**”.

For each of the following, what might the domain be?

(1) “x is a cat”, “x barks”, “x ruined my couch”

“mammals” or “sentient beings” or “cats and dogs” or ...

(2) “x is prime”, “ $x = 0$ ”, “ $x < 0$ ”, “x is a power of two”

“numbers” or “integers” or “integers greater than 5” or ...

(3) “x is a pre-req for z”

“courses”

Last Class: Quantifiers

We use *quantifiers* to talk about collections of objects.

$\forall x P(x)$

$P(x)$ is true **for every** x in the domain

read as “**for all** x , P of x ”



$\exists x P(x)$

There is an x in the domain for which $P(x)$ is true

read as “**there exists** x , P of x ”

Last class: Statements with Quantifiers

Domain of Discourse

Positive Integers

Predicate Definitions

Even(x) ::= "x is even"

Greater(x, y) ::= "x > y"

Odd(x) ::= "x is odd"

Equal(x, y) ::= "x = y"

Prime(x) ::= "x is prime"

Sum(x, y, z) ::= "x + y = z"

Determine the truth values of each of these statements:

$\exists x \text{ Even}(x)$

T

$\forall x \text{ Odd}(x)$

F

$\forall x (\text{Even}(x) \vee \text{Odd}(x))$

⊕

T

$\exists x (\text{Even}(x) \wedge \text{Odd}(x))$

F

$\forall x \text{ Greater}(x+1, x)$

T

$\exists x (\text{Even}(x) \wedge \text{Prime}(x))$

T

(2)

Statements with Quantifiers

Domain of Discourse

Positive Integers

Predicate Definitions

Even(x) ::= "x is even"

Greater(x, y) ::= "x > y"

Odd(x) ::= "x is odd"

Equal(x, y) ::= "x = y"

Prime(x) ::= "x is prime"

Sum(x, y, z) ::= "x + y = z"

Determine the truth values of each of these statements:

$\exists x \text{ Even}(x)$

T e.g. 2, 4, 6, ...

$\forall x \text{ Odd}(x)$

F e.g. 2, 4, 6, ...

$\forall x (\text{Even}(x) \vee \text{Odd}(x))$

T every integer is either even or odd

$\exists x (\text{Even}(x) \wedge \text{Odd}(x))$

F no integer is both even and odd

$\forall x \text{ Greater}(x+1, x)$

T adding 1 makes a bigger number

$\exists x (\text{Even}(x) \wedge \text{Prime}(x))$

T Even(2) is true and Prime(2) is true

$\forall x (\text{Even}(x) \wedge \text{Prime}(x) \rightarrow \text{Equal}(x, 2))$

Statements with Quantifiers

Domain of Discourse

Positive Integers

Predicate Definitions

Even(x) ::= "x is even"

Greater(x, y) ::= "x > y"

Odd(x) ::= "x is odd"

Equal(x, y) ::= "x = y"

Prime(x) ::= "x is prime"

Sum(x, y, z) ::= "x + y = z"

Translate the following statements to English

$\forall x \exists y \text{ Greater}(y, x)$

$\forall x \exists y \text{ Greater}(x, y)$

$\forall x \exists y (\text{Greater}(y, x) \wedge \text{Prime}(y))$

$\forall x (\text{Prime}(x) \rightarrow (\text{Equal}(x, 2) \vee \text{Odd}(x)))$

$\exists x \exists y (\text{Sum}(x, 2, y) \wedge \text{Prime}(x) \wedge \text{Prime}(y))$

Statements with Quantifiers (Literal Translations)

Domain of Discourse

Positive Integers

Predicate Definitions

Even(x) ::= "x is even"

Greater(x, y) ::= "x > y"

Odd(x) ::= "x is odd"

Equal(x, y) ::= "x = y"

Prime(x) ::= "x is prime"

Sum(x, y, z) ::= "x + y = z"

Translate the following statements to English

$\forall x \exists y \text{ Greater}(y, x)$

For every positive integer x, there is a positive integer y, such that y > x.

$\forall x \exists y \text{ Greater}(x, y)$

For every positive integer x, there is a positive integer y, such that x > y.

$\forall x \exists y (\text{Greater}(y, x) \wedge \text{Prime}(y))$

For every positive integer x, there is a pos. int. y such that y > x and y is prime.

$\forall x (\text{Prime}(x) \rightarrow (\text{Equal}(x, 2) \vee \text{Odd}(x)))$

For each positive integer x, if x is prime, then x = 2 or x is odd.

$\exists x \exists y (\text{Sum}(x, 2, y) \wedge \text{Prime}(x) \wedge \text{Prime}(y))$

There exist positive integers x and y such that x + 2 = y and x and y are prime.

Statements with Quantifiers (Natural Translations)

Domain of Discourse

Positive Integers

Predicate Definitions

Even(x) ::= "x is even"

Greater(x, y) ::= "x > y"

Odd(x) ::= "x is odd"

Equal(x, y) ::= "x = y"

Prime(x) ::= "x is prime"

Sum(x, y, z) ::= "x + y = z"

Translate the following statements to English

$\forall x \exists y \text{ Greater}(y, x)$

There is no greatest integer.

$\forall x \exists y \text{ Greater}(x, y)$

There is no least integer.

* $\forall x \exists y (\text{Greater}(y, x) \wedge \text{Prime}(y))$

For every positive integer there is a larger number that is prime.

$\forall x (\text{Prime}(x) \rightarrow (\text{Equal}(x, 2) \vee \text{Odd}(x)))$

Every prime number is either 2 or odd.

$\exists x \exists y (\text{Sum}(x, 2, y) \wedge \text{Prime}(x) \wedge \text{Prime}(y))$

There exist prime numbers that differ by two.

$\neg \exists x (\neg \exists y y > x)$

restrict to
primes
 \wedge

English to Predicate Logic

Domain of Discourse

Mammals

Predicate Definitions

Cat(x) ::= "x is a cat"

Red(x) ::= "x is red"

LikesTofu(x) ::= "x likes tofu"

(All)

"Red cats like tofu"

$$\forall x \text{ Red}(x) \wedge \text{Cat}(x) \rightarrow \text{LikeTofu}(x)$$

(X)

(There is a)

"Some red cats don't like tofu"

$$\exists x \text{ Red}(x) \wedge \text{Cat}(x) \wedge \neg \text{LikeTofu}(x)$$

(X)

English to Predicate Logic

Domain of Discourse

Mammals

Predicate Definitions

$\text{Cat}(x) ::= \text{"x is a cat"}$

$\text{Red}(x) ::= \text{"x is red"}$

$\text{LikesTofu}(x) ::= \text{"x likes tofu"}$

“Red cats like tofu”

$\forall x ((\text{Red}(x) \wedge \text{Cat}(x)) \rightarrow \text{LikesTofu}(x))$

“Some red cats don’t like tofu”

$\exists y ((\text{Red}(y) \wedge \text{Cat}(y)) \wedge \neg \text{LikesTofu}(y))$

English to Predicate Logic

Domain of Discourse

Mammals

Predicate Definitions

$\text{Cat}(x) ::= \text{"x is a cat"}$

$\text{Red}(x) ::= \text{"x is red"}$

$\text{LikesTofu}(x) ::= \text{"x likes tofu"}$

When putting two predicates together like this, we use an "and".

"Red cats like tofu"

When restricting to a smaller domain in a "for all" we use implication.

When there's no leading quantification, it means "for all".

"Some red cats don't like tofu"

When restricting to a smaller domain in an "exists" we use and.

"Some" means "there exists".

Negations of Quantifiers

Predicate Definitions

PurpleFruit(x) ::= “x is a purple fruit”

(*) $\forall x$ PurpleFruit(x) (“All fruits are purple”)

What is the negation of (*)?

- (a) “there exists a purple fruit”
- (b) “there exists a non-purple fruit”
- (c) “all fruits are not purple”

Try your intuition! Which one “feels” right?

Key Idea: In **every** domain, exactly one of a statement and its negation should be true.

Negations of Quantifiers

Predicate Definitions

PurpleFruit(x) ::= “x is a purple fruit”

(*) $\forall x$ PurpleFruit(x) (“All fruits are purple”)

What is the negation of (*)?

- (a) “there exists a purple fruit” ✗
- (b) “there exists a non-purple fruit”
- (c) “all fruits are not purple” ✗

Key Idea: In **every** domain, exactly one of a statement and its negation should be true.

Domain of Discourse

{plum}

(*), (a)

Domain of Discourse

{apple}

(b), (c)

Domain of Discourse

{plum, apple}

(a), (b)

false: (*), (c)

Negations of Quantifiers

Predicate Definitions

PurpleFruit(x) ::= “x is a purple fruit”

(*) $\forall x$ PurpleFruit(x) (“All fruits are purple”)

What is the negation of (*)?

- (a) “there exists a purple fruit”
- (b) “there exists a non-purple fruit”
- (c) “all fruits are not purple”

Key Idea: In **every** domain, exactly one of a statement and its negation should be true.

Domain of Discourse

{plum}

(*), (a)

Domain of Discourse

{apple}

(b), (c)

Domain of Discourse

{plum, apple}

(a), (b)

The only choice that ensures exactly one of the statement and its negation is (b).

De Morgan's Laws for Quantifiers

$$\neg \forall x P(x) \equiv \exists x \neg P(x)$$
$$\neg \exists x P(x) \equiv \forall x \neg P(x)$$

← previous example

De Morgan's Laws for Quantifiers

$$\neg \forall x P(x) \equiv \exists x \neg P(x)$$

$$\neg \exists x P(x) \equiv \forall x \neg P(x)$$

“There is no largest integer”

$$\begin{aligned} & \neg \exists x \forall y (x \geq y) \\ & \equiv \forall x \neg \forall y (x \geq y) \\ & \equiv \forall x \exists y \neg (x \geq y) \\ & \equiv \forall x \exists y (y > x) \end{aligned}$$

de Morgan

“For every integer there is a larger integer”

Scope of Quantifiers

$$\exists x (P(x) \wedge Q(x)) \quad \text{vs.} \quad \underline{\exists x P(x) \wedge \exists x Q(x)}$$

scope of quantifiers

$$\exists x (P(x) \wedge Q(x)) \quad \text{vs.} \quad (\exists x P(x)) \wedge (\exists x Q(x))$$

This one asserts P
and Q of the *same* x.

This one asserts P and Q
of potentially different x's.

Scope of Quantifiers

Example: $\text{NotLargest}(x) \equiv \exists y \text{ Greater}(y, x)$
 $\equiv \exists z \text{ Greater}(z, x)$

truth value:


doesn't depend on y or z “**bound** variables”

does depend on x “**free** variable”

quantifiers only act on free variables of the formula
they quantify

$\forall x (\exists y (P(x, y) \rightarrow \forall x Q(y, x)))$


Quantifier “Style”

$$\forall x(\exists y(P(x,y) \rightarrow \forall x Q(y,x)))$$



This isn't “wrong”, it's just horrible style.
Don't confuse your reader by using the same
variable multiple times...there are a lot of letters...

Nested Quantifiers

- **Bound variable names don't matter**

$$\forall x \exists y P(x, y) \equiv \forall a \exists b P(a, b)$$


- **Positions of quantifiers can sometimes change**

$$\forall x (Q(x) \wedge \exists y P(x, y)) \equiv \forall x \exists y (Q(x) \wedge P(x, y))$$


- **But: order is important...**

Quantifier Order Can Matter

Domain of Discourse

Integers
OR
{1, 2, 3, 4}

Predicate Definitions

GreaterEq(x, y) ::= "x ≥ y"

"There is a number greater than or equal to all numbers."

$\exists x \forall y \text{ GreaterEq}(x, y)$

"Every number has a number greater than or equal to it."

$\forall y \exists x \text{ GreaterEq}(x, y)$

--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--

GreaterEq.


The purple statement requires an entire row to be true.

The red statement requires one entry in each column to be true.

1st F
2nd T

1	T	F	F	F	
2	T	T	F	F	
3	T	T	T	F	
4	T	T	T	T	

Quantification with Two Variables



expression	when true	when false
$\forall x \forall y P(x, y)$	Every pair is true.	At least one pair is false.
$\exists x \exists y P(x, y)$	At least one pair is true.	All pairs are false.
$\forall x \exists y P(x, y)$	We can find a specific y for each x. (x_1, y_1), (x_2, y_2), (x_3, y_3)	Some x doesn't have a corresponding y. row of F_5
$\exists y \forall x P(x, y)$	We can find ONE y that works no matter what x is. (x_1, y), (x_2, y), (x_3, y) col of T_5	For any candidate y, there is an x that it doesn't work for.

Logical Inference

- So far we've considered:
 - How to understand and *express* things using propositional and predicate logic
 - How to *compute* using Boolean (propositional) logic
 - How to show that different ways of expressing or computing them are *equivalent* to each other
- Logic also has methods that let us *infer* implied properties from ones that we know
 - Equivalence is a small part of this

$$\underline{P(x) \wedge Q(x)} \Rightarrow \underline{P(x)}$$

Applications of Logical Inference

- **Software Engineering**
 - Express desired properties of program as set of logical constraints
 - Use inference rules to show that program implies that those constraints are satisfied
- **Artificial Intelligence**
 - Automated reasoning
- **Algorithm design and analysis**
 - e.g., Correctness, Loop invariants.
- **Logic Programming, e.g. Prolog**
 - Express desired outcome as set of constraints
 - Automatically apply logic inference to derive solution

Proofs

- **Start with hypotheses and facts**
- **Use rules of inference to extend set of facts**
- **Result is proved when it is included in the set**

An inference rule: *Modus Ponens*

- If p and $p \rightarrow q$ are both true then q must be true
- Write this rule as
$$\frac{p, p \rightarrow q}{\therefore q}$$
- Given:
 - If it is Monday then you have a 311 class today.
 - It is Monday.
- Therefore, by Modus Ponens:
 - You have a 311 class today.

My First Proof!

Show that r follows from p , $p \rightarrow q$, and $q \rightarrow r$

1. p Given
2. $p \rightarrow q$ Given
3. $q \rightarrow r$ Given
- 4.
- 5.

My First Proof!

Show that r follows from p , $p \rightarrow q$, and $q \rightarrow r$

- | | | |
|----|-------------------|----------|
| 1. | p | Given |
| 2. | $p \rightarrow q$ | Given |
| 3. | $q \rightarrow r$ | Given |
| 4. | q | MP: 1, 2 |
| 5. | r | MP: 3, 4 |

Proofs can use equivalences too

Show that $\neg p$ follows from $p \rightarrow q$ and $\neg q$

- | | | |
|----|-----------------------------|-------------------|
| 1. | $p \rightarrow q$ | Given |
| 2. | $\neg q$ | Given |
| 3. | $\neg q \rightarrow \neg p$ | Contrapositive: 1 |
| 4. | $\neg p$ | MP: 2, 3 |

Inference Rules

- Each **inference rule** is written as:
...which means that if both A and B
are true then you can infer C and
you can infer D.
 - For rule to be correct $(A \wedge B) \rightarrow C$ and
 $(A \wedge B) \rightarrow D$ must be a tautologies

$$\frac{A, B}{\therefore C, D}$$

- Sometimes rules don't need anything to start with.
These rules are called **axioms**:
 - e.g. *Excluded Middle Axiom*

$$\frac{}{\therefore p \vee \neg p}$$

Simple Propositional Inference Rules

Excluded middle plus two inference rules per binary connective, one to eliminate it and one to introduce it

$$\frac{p \wedge q}{\therefore p, q}$$

$$\frac{p, q}{\therefore p \wedge q}$$

$$\frac{p \vee q, \neg p}{\therefore q}$$

$$\frac{p}{\therefore p \vee q, q \vee p}$$

$$\frac{p, p \rightarrow q}{\therefore q}$$

$$\frac{p \Rightarrow q}{\therefore p \rightarrow q}$$

Direct Proof Rule
Not like other rules

Proofs

Show that r follows from p , $p \rightarrow q$ and $(p \wedge q) \rightarrow r$

How To Start:

We have givens, find the ones that go together and use them. Now, treat new things as givens, and repeat.

$$\frac{p, p \rightarrow q}{\therefore q}$$

$$\frac{p \wedge q}{\therefore p, q}$$

$$\frac{p, q}{\therefore p \wedge q}$$

Proofs

Show that r follows from $p, p \rightarrow q$, and $p \wedge q \rightarrow r$

Two visuals of the same proof.
We will use the top one, but if
the bottom one helps you
think about it, that's great!

- | | | |
|----|----------------------------|-----------------------|
| 1. | p | Given |
| 2. | $p \rightarrow q$ | Given |
| 3. | q | MP: 1, 2 |
| 4. | $p \wedge q$ | Intro \wedge : 1, 3 |
| 5. | $p \wedge q \rightarrow r$ | Given |
| 6. | r | MP: 4, 5 |

$$\begin{array}{c}
 \begin{array}{c} p \quad p \rightarrow q \\ \hline q \end{array} \text{MP} \\
 \begin{array}{c} p \quad q \\ \hline p \wedge q \end{array} \text{Intro } \wedge \\
 \begin{array}{c} p \wedge q \quad p \wedge q \rightarrow r \\ \hline r \end{array} \text{MP}
 \end{array}$$

Important: Applications of Inference Rules

- You can use equivalences to make substitutions of any sub-formula.
- Inference rules only can be applied to whole formulas (not correct otherwise).

e.g. 1. $p \rightarrow q$ given
~~2. $(p \vee r) \rightarrow q$ intro \vee from 1.~~

Does not follow! e.g. $p=F, q=F, r=T$