

# CSE 311: Foundations of Computing

---

## Lecture 8: Predicate Logic Proofs

- Pick up HW1 sol's if you didn't get one on Monday
- Correction on sol's of 1(b)  
solar + hydrogen



THE AXIOM OF CHOICE ALLOWS  
YOU TO SELECT ONE ELEMENT  
FROM EACH SET IN A COLLECTION  
AND HAVE IT EXECUTED AS  
AN EXAMPLE TO THE OTHERS.

MY MATH TEACHER WAS A BIG  
BELIEVER IN PROOF BY INTIMIDATION.

# Last class: Propositional Inference Rules

Two inference rules per binary connective, one to eliminate it and one to introduce it

$$\text{Elim } \wedge \frac{A \wedge B}{\therefore A, B} \quad \checkmark$$

$$\text{Intro } \wedge \frac{A; B}{\therefore A \wedge B}, \quad \checkmark$$

$\beta \wedge A$

$$\text{Elim } \vee \frac{A \vee B; \neg A}{\therefore B}$$

↙

$$\text{Intro } \vee \frac{A}{\therefore A \vee B, B \vee A} \quad \checkmark$$

$$\text{Modus Ponens} \frac{A; A \rightarrow B}{\therefore B}$$


$$\text{Direct Proof Rule} \frac{A \Rightarrow B}{\therefore A \rightarrow B}$$

Not like other rules

# Last class: Example

---

Prove:  $((p \rightarrow q) \wedge (q \rightarrow r)) \rightarrow (p \rightarrow r)$



# One General Proof Strategy

---

1. Look at the rules for introducing connectives to see how you would build up the formula you want to prove from pieces of what is given
  
2. Use the rules for eliminating connectives to break down the given formulas so that you get the pieces you need to do 1.
  
3. Write the proof beginning with what you figured out for 2 followed by 1.

# Inference Rules for Quantifiers: First look

$$\frac{\text{Intro } \exists \quad P(c) \text{ for some } c}{\therefore \exists x P(x)}$$

$$\frac{\text{Elim } \forall \quad \forall x P(x)}{\therefore P(a) \text{ for any } a}$$

$$\frac{\text{Intro } \forall \quad \text{"Let } a \text{ be arbitrary*"} \dots P(a)}{\therefore \forall x P(x)}$$

$$\frac{\text{Elim } \exists \quad \exists x P(x)}{\therefore P(c) \text{ for some special** } c}$$

\* in the domain of P

e.g.  $\frac{\text{Let } a \text{ be arbitrary.} \quad Q(a) \vee \neg Q(a) \quad A \vdash \alpha}{\therefore \forall x (Q(x) \vee \neg Q(x))}$

\*\* By special, we mean that c is a name for a value where  $P(c)$  is true. We can't use anything else about that value, so c has to be a NEW name!

# Predicate Logic Proofs

---

- Can use
  - Predicate logic inference rules
    - whole formulas only
  - Predicate logic equivalences (De Morgan's)
    - even on subformulas
  - Propositional logic inference rules
    - whole formulas only
  - Propositional logic equivalences
    - even on subformulas

# My First Predicate Logic Proof

Prove  $\forall x P(x) \rightarrow \exists x P(x)$

$$\frac{\text{Intro } \exists \quad P(c) \text{ for some } c}{\therefore \exists x P(x)}$$

$$\frac{\text{Elim } \forall \quad \forall x P(x)}{\therefore P(a) \text{ for any } a}$$

$\forall x P(x)$       Assumption  
    }       $\exists x P(x)$

5.  $\forall x P(x) \rightarrow \exists x P(x)$



The main connective is implication  
so Direct Proof Rule seems good

# My First Predicate Logic Proof

Prove  $\forall x P(x) \rightarrow \exists x P(x)$

$$\frac{\text{Intro } \exists \quad P(c) \text{ for some } c}{\therefore \exists x P(x)}$$
$$\frac{\text{Elim } \forall \quad \forall x P(x)}{\therefore P(a) \text{ for any } a}$$

1.1.

$\forall x P(x)$

Assumption

We need an  $\exists$  we don't have  
so “intro  $\exists$ ” rule makes sense

1.5.

$\exists x P(x)$

?

1.  $\forall x P(x) \rightarrow \exists x P(x)$  Direct Proof Rule

# My First Predicate Logic Proof

Prove  $\forall x P(x) \rightarrow \exists x P(x)$

$$\begin{array}{c} \text{Intro } \exists \\ \hline P(c) \text{ for some } c \\ \therefore \exists x P(x) \end{array}$$

$$\begin{array}{c} \text{Elim } \forall \\ \hline \forall x P(x) \\ \therefore P(a) \text{ for any } a \end{array}$$

1.1.

$\forall x P(x)$

Assumption

We need an  $\exists$  we don't have  
so “intro  $\exists$ ” rule makes sense

1.5.

$\exists x P(x)$

Intro  $\exists$ :

?

That requires  $P(c)$   
for some  $c$ .

1.

$\forall x P(x) \rightarrow \exists x P(x)$

Direct Proof Rule

# My First Predicate Logic Proof

Prove  $\forall x P(x) \rightarrow \exists x P(x)$

$$\frac{\text{Intro } \exists \quad P(c) \text{ for some } c}{\therefore \exists x P(x)}$$

$$\frac{\text{Elim } \forall \quad \forall x P(x)}{\therefore P(a) \text{ for any } a}$$

1.1.  $\forall x P(x)$

1.2  $P(a)$

We could have picked any name  
or domain expression here.

Assumption

Elim  $\forall$ : 1.1

1.5.  $\exists x P(x)$

Intro  $\exists$ : ?

1. 2

That requires  $P(c)$   
for some  $c$ .

1.  $\forall x P(x) \rightarrow \exists x P(x)$  Direct Proof Rule

# My First Predicate Logic Proof

$$\frac{\text{Intro } \exists \quad P(c) \text{ for some } c}{\therefore \exists x P(x)}$$
$$\frac{\text{Elim } \forall \quad \forall x P(x)}{\therefore P(a) \text{ for any } a}$$

Prove  $\forall x P(x) \rightarrow \exists x P(x)$

No holes. Just need to clean up.

1.1.  $\forall x P(x)$  Assumption  
1.2.  $P(a)$  Elim  $\forall$ : 1.1

1.5.  $\exists x P(x)$  Intro  $\exists$ : 1.2

1.  $\forall x P(x) \rightarrow \exists x P(x)$  Direct Proof Rule

# My First Predicate Logic Proof

$$\frac{\text{Intro } \exists \quad P(c) \text{ for some } c}{\therefore \exists x P(x)}$$

$$\frac{\text{Elim } \forall \quad \forall x P(x)}{\therefore P(a) \text{ for any } a}$$

Prove  $\forall x P(x) \rightarrow \exists x P(x)$

1.1.  $\forall x P(x)$

Assumption

1.2.  $P(a)$

Elim  $\forall$ : 1.1

1.2.  $P(a)$

1.3.  $\exists x P(x)$

Intro  $\exists$ : 1.2

1.3.  $P(b)$

1.  $\forall x P(x) \rightarrow \exists x P(x)$  Direct Proof Rule

1.4.  $P(c)$   
1.5.  $P(d)$

Working forwards as well as backwards:

In applying “Intro  $\exists$ ” rule we didn’t know what expression to use. We might be able to prove  $P(c)$  for, so we worked forwards to figure out what might work.

# Predicate Logic Proofs with more content

---

- In propositional logic we could just write down other propositional logic statements as “givens”
- Here, we also want to be able to use domain knowledge so proofs are about something specific
- Example:
- Given the basic properties of arithmetic on integers, define:

Domain of Discourse
Integers

Predicate Definitions
$\text{Even}(x) \equiv \exists y (x = 2 \cdot y)$
$\text{Odd}(x) \equiv \exists y (x = 2 \cdot y + 1)$

# A Not so Odd Example

---

Domain of Discourse

Integers

Predicate Definitions

$$\text{Even}(x) \equiv \exists y (x = 2 \cdot y)$$

$$\text{Odd}(x) \equiv \exists y (x = 2 \cdot y + 1)$$

Prove “There is an even number”

Formally: prove  $\exists x \text{ Even}(x)$

# A Not so Odd Example

---

Domain of Discourse  
Integers

Predicate Definitions  
 $\text{Even}(x) \equiv \exists y (x = 2 \cdot y)$   
 $\text{Odd}(x) \equiv \exists y (x = 2 \cdot y + 1)$

Prove “There is an even number”

Formally: prove  $\exists x \text{ Even}(x)$

1.  $2 = 2 \cdot 1$  Arithmetic
2.  $\exists y (2 = 2 \cdot y)$  Intro  $\exists: 1$
3.  $\text{Even}(2)$  Definition of Even: 2
4.  $\exists x \text{ Even}(x)$  Intro  $\exists: 3$

# A Prime Example

---

**Domain of Discourse**  
Integers

**Predicate Definitions**

$$\text{Even}(x) \equiv \exists y (x = 2 \cdot y)$$

$$\text{Odd}(x) \equiv \exists y (x = 2 \cdot y + 1)$$

$\text{Prime}(x) \equiv "x > 1 \text{ and } x \neq a \cdot b \text{ for}$   
 $\text{all integers } a, b \text{ with } 1 < a < x"$

Prove “There is an even prime number”

# A Prime Example

---

Domain of Discourse  
Integers

Predicate Definitions

$$\text{Even}(x) \equiv \exists y (x = 2 \cdot y)$$

$$\text{Odd}(x) \equiv \exists y (x = 2 \cdot y + 1)$$

$\text{Prime}(x) \equiv "x > 1 \text{ and } x \neq a \cdot b \text{ for}$   
 $\text{all integers } a, b \text{ with } 1 < a < x"$

Prove “There is an even prime number”

Formally: prove  $\exists x (\text{Even}(x) \wedge \text{Prime}(x))$

1.  $2 = 2 \cdot 1$
2.  $\text{Prime}(2)^*$

Arithmetic  
Property of integers

\* Later we will further break down “Prime” using quantifiers to prove statements like this

# A Prime Example

---

Domain of Discourse  
Integers

Predicate Definitions

$$\text{Even}(x) \equiv \exists y (x = 2 \cdot y)$$

$$\text{Odd}(x) \equiv \exists y (x = 2 \cdot y + 1)$$

$\text{Prime}(x) \equiv "x > 1 \text{ and } x \neq a \cdot b \text{ for}$   
 $\text{all integers } a, b \text{ with } 1 < a < x"$

Prove “There is an even prime number”

Formally: prove  $\exists x (\text{Even}(x) \wedge \text{Prime}(x))$

- |    |   |                      |
|----|---|----------------------|
| 1. | $2 = 2 \cdot 1$                                     | Arithmetic           |
| 2. | $\text{Prime}(2)^*$                                 | Property of integers |
| 3. | $\exists y (2 = 2 \cdot y)$                         | Intro $\exists: 1$   |
| 4. | $\text{Even}(2)$                                    | Defn of Even: 3      |
| 5. | $\text{Even}(2) \wedge \text{Prime}(2)$             | Intro $\wedge: 2, 4$ |
| 6. | $\exists x (\text{Even}(x) \wedge \text{Prime}(x))$ | Intro $\exists: 5$   |

\* Later we will further break down “Prime” using quantifiers to prove statements like this

# Inference Rules for Quantifiers: First look

---

$$\frac{\text{Intro } \exists \quad P(c) \text{ for some } c}{\therefore \exists x P(x)}$$

$$\frac{\text{Elim } \forall \quad \forall x P(x)}{\therefore P(a) \text{ for any } a}$$

$$\frac{\text{Intro } \forall \quad \text{"Let } a \text{ be arbitrary*"} \dots P(a)}{\therefore \forall x P(x)}$$

\* in the domain of P

$$\frac{\text{Elim } \exists \quad \exists x P(x)}{\therefore P(c) \text{ for some special** } c}$$

\*\* By special, we mean that  $c$  is a name for a value where  $P(c)$  is true. We can't use anything else about that value, so  $c$  has to be a NEW name!

# Even and Odd

Even( $x$ )  $\equiv \exists y (x=2y)$   
Odd( $x$ )  $\equiv \exists y (x=2y+1)$   
Domain: Integers

Intro  $\forall$

"Let  $a$  be arbitrary\*" ...  $P(a)$

$\therefore \forall x P(x)$

Elim  $\exists$

$\exists x P(x)$

$\therefore P(c)$  for some special\*\*  $c$

Prove: "The square of every even number is even."

Formal proof of:  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

1. Let  $a$  be an arbitrary integer

$\text{Even}(a) \rightarrow \text{Even}(a^2)$

3.  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

~~?~~ Intro  $\forall : 1 - \exists$

# Even and Odd

Even( $x$ )  $\equiv \exists y$  ( $x=2y$ )  
Odd( $x$ )  $\equiv \exists y$  ( $x=2y+1$ )  
Domain: Integers

Intro  $\forall$

“Let  $a$  be arbitrary\*” ...  $P(a)$

$\therefore \forall x P(x)$

Elim  $\exists$

$\exists x P(x)$

$\therefore P(c)$  for some special\*\*  $c$

Prove: “The square of every even number is even.”

Formal proof of:  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

1. Let  $a$  be an arbitrary integer

2.  $\text{Even}(a) \rightarrow \text{Even}(a^2)$

3.  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$



Intro  $\forall$ : 1,2

# Even and Odd

Even( $x$ )  $\equiv \exists y$  ( $x=2y$ )  
Odd( $x$ )  $\equiv \exists y$  ( $x=2y+1$ )  
Domain: Integers

1. Let $a$ be an arbitrary integer.	Assumption
2.1 Even( $a$ )	
2.6 Even( $a^2$ )	Elim $\exists$
2. Even( $a$ ) $\rightarrow$ Even( $a^2$ )	$\exists x P(x)$
3. $\forall x$ (Even( $x$ ) $\rightarrow$ Even( $x^2$ ))	Direct proof rule Intro $\forall$ : 1,2

$\therefore P(c)$  for some *special\*\* c*

Prove: “The square of every even number is even.”

Formal proof of:  $\forall x$  (Even( $x$ )  $\rightarrow$  Even( $x^2$ ))

1. Let  $a$  be an arbitrary integer

2.1 Even( $a$ )

Assumption

2.6 Even( $a^2$ )

2. Even( $a$ )  $\rightarrow$  Even( $a^2$ )

3.  $\forall x$  (Even( $x$ )  $\rightarrow$  Even( $x^2$ ))



Direct proof rule

Intro  $\forall$ : 1,2

# Even and Odd

Even( $x$ )  $\equiv \exists y$  ( $x=2y$ )  
Odd( $x$ )  $\equiv \exists y$  ( $x=2y+1$ )  
Domain: Integers

Intro  $\forall$

“Let  $a$  be arbitrary\*” ...  $P(a)$

$\therefore \forall x P(x)$

Elim  $\exists$

$\exists x P(x)$

$\therefore P(c)$  for some special\*\*  $c$

Prove: “The square of every even number is even.”

Formal proof of:  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

1. Let  $a$  be an arbitrary integer

2.1 Even( $a$ )

Assumption

2.2  $\exists y (a = 2y)$

Definition of Even

2.5  $\exists y (a^2 = 2y)$



Definition of Even

2.6 Even( $a^2$ )

Direct proof rule

2.  $\text{Even}(a) \rightarrow \text{Even}(a^2)$

Intro  $\forall$ : 1,2

3.  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

# Even and Odd

Even( $x$ )  $\equiv \exists y$  ( $x=2y$ )  
Odd( $x$ )  $\equiv \exists y$  ( $x=2y+1$ )  
Domain: Integers

Intro  $\forall$

“Let  $a$  be arbitrary\*” ...  $P(a)$

$\therefore \forall x P(x)$

Elim  $\exists$

$\exists x P(x)$

$\therefore P(c)$  for some special\*\*  $c$

Prove: “The square of every even number is even.”

Formal proof of:  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

1. Let  $a$  be an arbitrary integer

2.1 Even( $a$ )

Assumption

2.2  $\exists y (a = 2y)$

Definition of Even

2.5  $\exists y (a^2 = 2y)$

Intro  $\exists$  rule: ?

Need  $a^2 = 2c$   
for some  $c$

2.6 Even( $a^2$ )

Definition of Even

2.  $\text{Even}(a) \rightarrow \text{Even}(a^2)$

Direct proof rule

3.  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

Intro  $\forall$ : 1,2

# Even and Odd

Even( $x$ )  $\equiv \exists y$  ( $x=2y$ )  
Odd( $x$ )  $\equiv \exists y$  ( $x=2y+1$ )  
Domain: Integers

Intro  $\forall$

“Let  $a$  be arbitrary\*” ...  $P(a)$

$\therefore \forall x P(x)$

Elim  $\exists$

$\exists x P(x)$

$\therefore P(c)$  for some special\*\*  $c$

Prove: “The square of every even number is even.”

Formal proof of:  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

1. Let  $a$  be an arbitrary integer

2.1 Even( $a$ )

Assumption

2.2  $\exists y (a = 2y)$

Definition of Even

2.3  $a = 2b$

Elim  $\exists$ :  $b$  special depends on  $a$

2.5  $\exists y (a^2 = 2y)$

Intro  $\exists$  rule: ?

Need  $a^2 = 2c$   
for some  $c$

2.6 Even( $a^2$ )

Definition of Even

2.  $\text{Even}(a) \rightarrow \text{Even}(a^2)$

Direct proof rule

3.  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

Intro  $\forall$ : 1,2

# Even and Odd

Even( $x$ )  $\equiv \exists y$  ( $x=2y$ )  
Odd( $x$ )  $\equiv \exists y$  ( $x=2y+1$ )  
Domain: Integers

Intro  $\forall$

“Let  $a$  be arbitrary\*” ...  $P(a)$

$\therefore \forall x P(x)$

Elim  $\exists$

$\exists x P(x)$

$\therefore P(c)$  for some special\*\*  $c$

Prove: “The square of every even number is even.”

Formal proof of:  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

1. Let  $a$  be an arbitrary integer

2.1 Even( $a$ )

Assumption

2.2  $\exists y (a = 2y)$

Definition of Even

2.3  $a = 2b$

Elim  $\exists$ :  $b$  special depends on  $a$

2.4  $a^2 = 4b^2 = 2(2b^2)$

Algebra

2.5  $\exists y (a^2 = 2y)$

Intro  $\exists$  rule

Used  $a^2 = 2c$  for  $c=2b^2$

2.6 Even( $a^2$ )

Definition of Even

2. Even( $a$ )  $\rightarrow$  Even( $a^2$ )

Direct proof rule

3.  $\forall x (\text{Even}(x) \rightarrow \text{Even}(x^2))$

Intro  $\forall$ : 1,2

# Why did we need to say that **b** depends on **a**?

There are extra conditions on using these rules:

$$\frac{\text{Intro } \forall \quad \boxed{\text{“Let } a \text{ be arbitrary*”} \dots P(a)} \quad \text{Elim } \exists}{\therefore \quad \forall x P(x) \qquad \qquad \qquad \therefore P(c) \text{ for some special** } c}$$

\* in the domain of  $P$

\*\*  $c$  has to be a NEW name.

Over integer domain:  $\forall x \exists y (y \geq x)$  is **True** but  $\exists y \forall x (y \geq x)$  is **False**

## BAD “PROOF”

1.  $\forall x \exists y (y \geq x)$  Given
2. Let **a** be an arbitrary integer
3.  $\exists y (y \geq a)$  Elim  $\forall$ : 1
4.  $b \geq a$  Elim  $\exists$ : **b** special depends on **a**
5.  ~~$\forall x (b \geq x)$~~  Intro  $\forall$ : 2,4
6.  $\exists y \forall x (y \geq x)$  Intro  $\exists$  : 5

# Why did we need to say that **b** depends on **a**?

There are extra conditions on using these rules:

<small>Intro <math>\forall</math></small>	<u>“Let <b>a</b> be arbitrary*” ... <math>P(a)</math></u>	<small>Elim <math>\exists</math></small>	$\exists x P(x)$
∴	$\forall x P(x)$	∴	$P(c)$ for some <b>special** c</b>
* in the domain of $P$		** $c$ has to be a NEW name.	

Over integer domain:  $\forall x \exists y (y \geq x)$  is **True** but  $\exists y \forall x (y \geq x)$  is **False**

## BAD “PROOF”

1.  $\forall x \exists y (y \geq x)$  Given
2. Let **a** be an arbitrary integer
3.  $\exists y (y \geq a)$  Elim  $\forall$ : 1
4. **b**  $\geq a$  Elim  $\exists$ : **b** special depends on **a**
5.  $\forall x (b \geq x)$  Intro  $\forall$ : 2,4
6.  $\exists y \forall x (y \geq x)$  Intro  $\exists$  : 5

Can't get rid of **a** since another name in the same line, **b**, depends on it!

# Why did we need to say that **b** depends on **a**?

There are extra conditions on using these rules:

$$\text{Intro } \forall \quad \frac{\text{“Let } a \text{ be arbitrary*”} \dots P(a)}{\therefore \forall x P(x)}$$

$$\text{Elim } \exists \quad \frac{\exists x P(x)}{\therefore P(c) \text{ for some special** } c}$$

\* in the domain of  $P$ . No other name in  $P$  depends on  $a$

\*\*  $c$  is a NEW name.  
List all dependencies for  $c$ .

Over integer domain:  $\forall x \exists y (y \geq x)$  is True but  $\exists y \forall x (y \geq x)$  is False

## BAD “PROOF”

1.  $\forall x \exists y (y \geq x)$  Given
2. Let **a** be an arbitrary integer
3.  $\exists y (y \geq a)$  Elim  $\forall$ : 1
4.  $b \geq a$  Elim  $\exists$ : **b** special depends on **a**
5.  ~~$\forall x (b \geq x)$~~  Intro  $\forall$ : 2,4
6.  $\exists y \forall x (y \geq x)$  Intro  $\exists$  : 5

Can't get rid of **a** since another name in the same line, **b**, depends on it!

# Inference Rules for Quantifiers: Full version

---

$$\frac{\text{Intro } \exists \quad P(c) \text{ for some } c}{\therefore \exists x P(x)}$$

$$\frac{\text{Elim } \forall \quad \forall x P(x)}{\therefore P(a) \text{ for any } a}$$

$$\frac{\text{Intro } \forall \quad \text{“Let } a \text{ be arbitrary*”} \dots P(a)}{\therefore \forall x P(x)}$$

\* in the domain of P. No other name in P depends on a

$$\frac{\text{Elim } \exists \quad \exists x P(x)}{\therefore P(c) \text{ for some special** } c}$$

\*\* c is a NEW name.  
List all dependencies for c.

# **English Proofs**

---

- We often write proofs in English rather than as fully formal proofs
  - They are more natural to read
- English proofs follow the structure of the corresponding formal proofs
  - Formal proof methods help to understand how proofs really work in English...  
... and give clues for how to produce them.

# An English Proof

## Predicate Definitions

$$\text{Even}(x) \equiv \exists y (x = 2 \cdot y)$$

$$\text{Odd}(x) \equiv \exists y (x = 2 \cdot y + 1)$$

Prove “There is an even integer”

Proof:

$$2 = 2 \cdot 1$$



1.  $2 = 2 \cdot 1$

Arithmetic

so **2** equals **2** times an  
integer.



2.  $\exists y (2 = 2 \cdot y)$

Intro  $\exists$ : 1

Therefore **2** is even.



3.  $\text{Even}(2)$

Defn of Even: 2

Therefore, there is an  
even integer ■



4.  $\exists x \text{ Even}(x)$

Intro  $\exists$ : 3

# English Even and Odd

Even( $x$ )  $\equiv \exists y$  ( $x=2y$ )  
Odd( $x$ )  $\equiv \exists y$  ( $x=2y+1$ )  
Domain: Integers

Prove “The square of every even integer is even.”

Proof: Let  $a$  be an arbitrary even integer.

1. Let  $a$  be an arbitrary integer  
2.1 Even( $a$ ) Assumption

Then, by definition,  $a = 2b$  for some integer  $b$   
(depending on  $a$ ).

2.2  $\exists y$  ( $a = 2y$ ) Definition  
2.3  $a = 2b$   $b$  special depends on  $a$

Squaring both sides, we get  $a^2 = 4b^2 = 2(2b^2)$ .

2.4  $a^2 = 4b^2 = 2(2b^2)$  Algebra

Since  $2b^2$  is an integer, by definition,  $a^2$  is even.

2.5  $\exists y$  ( $a^2 = 2y$ )

2.6 Even( $a^2$ ) Definition

Since  $a$  was arbitrary, it follows that the square of every even number is even. ■

2. Even( $a$ )  $\rightarrow$  Even( $a^2$ )  
3.  $\forall x$  (Even( $x$ )  $\rightarrow$  Even( $x^2$ ))

# Even and Odd

## Predicate Definitions

$$\text{Even}(x) \equiv \exists y (x = 2y)$$

$$\text{Odd}(x) \equiv \exists y (x = 2y + 1)$$

## Domain of Discourse

Integers

Prove "The square of every odd number is odd."

Proof: Let  $a$  be an arbitrary odd integer.

$\therefore a = 2b+1$  for some integer  $b$   
(depends on  $a$ )

$$\begin{aligned}\therefore a^2 &= (2b+1)^2 \\ &= 4b^2 + 4b + 1 \\ &= 2(2b^2 + b) + 1\end{aligned}$$

$$\therefore a^2 = 2c + 1 \text{ for integer } c \quad (= 2b^2 + b)$$

$\therefore a^2$  is odd

so -

# Even and Odd

## Predicate Definitions

$$\text{Even}(x) \equiv \exists y (x = 2y)$$

$$\text{Odd}(x) \equiv \exists y (x = 2y + 1)$$

## Domain of Discourse

Integers

Prove “The square of every odd number is odd.”

Proof: Let b be an arbitrary odd number.

Then,  $b = 2c+1$  for some integer c (depending on b).

Therefore,  $b^2 = (2c+1)^2 = 4c^2 + 4c + 1 = 2(2c^2 + 2c) + 1$ .

Since  $2c^2+2c$  is an integer,  $b^2$  is odd. The statement follows since b was arbitrary. ■

# Proofs

---

- Formal proofs follow simple well-defined rules and should be easy to check
  - In the same way that code should be easy to execute
- English proofs correspond to those rules but are designed to be easier for humans to read
  - Easily checkable in principle