


CSE 311: Foundations of Computing

Lecture 11: Modular Arithmetic and Applications



Last Class: Divisibility

Definition: “a divides b”

For $a \in \mathbb{Z}, b \in \mathbb{Z}$ with $a \neq 0$:

$$a \mid b \leftrightarrow \exists k \in \mathbb{Z} (b = ka)$$

Check Your Understanding. Which of the following are true?

$$5 \mid 1$$

$$5 \mid 1 \text{ iff } 1 = 5k$$

$$25 \mid 5$$

$$25 \mid 5 \text{ iff } 5 = 25k$$

$$5 \mid 0$$

$$5 \mid 0 \text{ iff } 0 = 5k$$

$$3 \mid 2$$

$$3 \mid 2 \text{ iff } 2 = 3k$$

$$1 \mid 5$$

$$1 \mid 5 \text{ iff } 5 = 1k$$

$$5 \mid 25$$

$$5 \mid 25 \text{ iff } 25 = 5k$$

$$0 \mid 5$$

$$0 \mid 5 \text{ iff } 5 = 0k$$

$$2 \mid 3$$

$$2 \mid 3 \text{ iff } 3 = 2k$$

Last Class: Division Theorem

Division Theorem

For $a \in \mathbb{Z}, d \in \mathbb{Z}$ with $d > 0$

there exist unique integers q, r with $0 \leq r < d$
such that $a = dq + r$.

$$r = a - dq \\ = a - (a \text{ div } d) d$$

To put it another way, if we divide d into a , we get a

unique quotient $q = a \text{ div } d$

and non-negative remainder

$r = a \text{ mod } d$

```
public class Test2 {  
    public static void main(String args[]) {  
        int a = -5;  
        int d = 2;  
        System.out.println(a % d);  
    }  
}
```


```
----jGRASP exec: java Test2  
-1  
----jGRASP: operation complete.
```

Note: $r \geq 0$ even if $a < 0$.
Not quite the same as $a \% d$.



Last Class: Arithmetic, mod 7

$$a +_7 b = (a + b) \bmod 7$$


$$a \times_7 b = (a \times b) \bmod 7$$



+	0	1	2	3	4	5	6
0	0	1	2	3	4	5	6
1	1	2	3	4	5	6	0
2	2	3	4	5	6	0	1
3	3	4	5	6	0	1	2
4	4	5	6	0	1	2	3
5	5	6	0	1	2	3	4
6	6	0	1	2	3	4	5



x	0	1	2	3	4	5	6
0	0	0	0	0	0	0	0
1	0	1	2	3	4	5	6
2	0	2	4	6	1	3	5
3	0	3	6	2	5	1	4
4	0	4	1	5	2	6	3
5	0	5	3	1	6	4	2
6	0	6	5	4	3	2	1



Modular Arithmetic

Definition: “a is congruent to b modulo m”

For $a, b, m \in \mathbb{Z}$ with $m > 0$

$$a \equiv b \pmod{m} \leftrightarrow m \mid (a - b)$$

Check Your Understanding. What do each of these mean?
When are they true?

$$x \equiv 0 \pmod{2}$$

$$2 \mid (x - 0) = x \quad x \text{ is even.}$$

$$-1 \equiv 19 \pmod{5}$$

$$5 \mid 19 - (-1) = 20$$

True

$$y \equiv 2 \pmod{7}$$

$$\exists k \ y = 2 - 7k$$

$$7 \mid 2 - y \leftrightarrow \exists k \ 7k = 2 - y$$

Modular Arithmetic

Definition: “a is congruent to b modulo m”

For $a, b, m \in \mathbb{Z}$ with $m > 0$

$$a \equiv b \pmod{m} \leftrightarrow m \mid (a - b)$$

**Check Your Understanding. What do each of these mean?
When are they true?**

$$x \equiv 0 \pmod{2}$$

This statement is the same as saying “x is even”; so, any x that is even (including negative even numbers) will work.

$$-1 \equiv 19 \pmod{5}$$

This statement is true. $19 - (-1) = 20$ which is divisible by 5

$$y \equiv 2 \pmod{7}$$

This statement is true for y in $\{ \dots, -12, -5, 2, 9, 16, \dots \}$. In other words, all y of the form $2+7k$ for k an integer.

Modular Arithmetic: A Property

Let a, b, m be integers with $m > 0$.

Then, $a \equiv b \pmod{m}$ if and only if $a \bmod m = b \bmod m$.

Suppose that $a \equiv b \pmod{m}$.

By defn, $m \mid a - b$, so $a - b = mk$ for some k .
Equivalently, $a = b + mk$. By Div Thm, $b = qm + r$
for some $0 \leq r < m$. Thus, $a = b + mk =$
 $(qm + r) + mk = (q + k)m + r$. By Div Thm,

Suppose that $a \bmod m = b \bmod m$. $b \bmod m = r = a \bmod m$.

Write $a = km + r$ and $b = jm + r$ where
 $r = a \bmod m = b \bmod m$. Thus, $a - b =$
 $(km + r) - (jm + r) = (k - j)m$, which proves
 $a \equiv b \pmod{m}$.

Modular Arithmetic: A Property

Let a, b, m be integers with $m > 0$.

Then, $a \equiv b \pmod{m}$ if and only if $a \bmod m = b \bmod m$.

Suppose that $a \equiv b \pmod{m}$.

Then, $m \mid (a - b)$ by definition of congruence.

So, $a - b = km$ for some integer k by definition of divides.

Therefore, $a = b + km$.

Taking both sides modulo m we get:

$$a \bmod m = (b + km) \bmod m = b \bmod m.$$

Suppose that $a \bmod m = b \bmod m$.

By the division theorem, $a = mq + (a \bmod m)$ and

$$b = ms + (b \bmod m) \text{ for some integers } q, s.$$

$$\begin{aligned} \text{Then, } a - b &= (mq + (a \bmod m)) - (ms + (b \bmod m)) \\ &= m(q - s) + (a \bmod m - b \bmod m) \\ &= m(q - s) \text{ since } a \bmod m = b \bmod m \end{aligned}$$

Therefore, $m \mid (a - b)$ and so $a \equiv b \pmod{m}$.

The mod m function vs the $\equiv \pmod{m}$ predicate

- What we have just shown
 - The mod m function takes any $a \in \mathbb{Z}$ and maps it to a remainder $a \bmod m \in \{0, 1, \dots, m - 1\}$.
 - Imagine grouping together all integers that have the same value of the mod m function
That is, the same remainder in $\{0, 1, \dots, m - 1\}$.
 - The $\equiv \pmod{m}$ predicate compares $a, b \in \mathbb{Z}$. It is true if and only if the mod m function has the same value on a and on b .
That is, a and b are in the same group.

$$\{a, a + m, a + 2m, \dots\}$$

Modular Arithmetic: Addition Property

Let m be a positive integer. If $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, then $a + c \equiv b + d \pmod{m}$

Suppose $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$

That means $a - b = km$ for some $k \in \mathbb{Z}$,

or equivalently, $a = b + km$. Likewise,

we have $c = d + lm$ for some $l \in \mathbb{Z}$

Thus, $a + c = b + km + d + lm =$

$b + d + (k + l)m$, which means

$$(a + c) - (b + d) = \underline{(k + l)m} \text{ so}$$

$$a + c \equiv b + d \pmod{m}.$$

Modular Arithmetic: Addition Property

Let m be a positive integer. If $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, then $a + c \equiv b + d \pmod{m}$

Suppose that $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$. Unrolling definitions gives us some k such that $a - b = km$, and some j such that $c - d = jm$.

Adding the equations together gives us

$(a + c) - (b + d) = m(k + j)$. Now, re-applying the definition of congruence gives us $a + c \equiv b + d \pmod{m}$.

Modular Arithmetic: Multiplication Property

Let m be a positive integer. If $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, then $ac \equiv bd \pmod{m}$

As before, $a = b + km$ and $c = d + lm$
for some $k, l \in \mathbb{Z}$. Then $ac =$
 $(b + km)(d + lm) = bd + b lm +$
 $k m d + k l m^2 = \underline{bd} + (lb + kd + kl m)m,$
which means $ac \equiv bd \pmod{m}$

Modular Arithmetic: Multiplication Property

Let m be a positive integer. If $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$, then $ac \equiv bd \pmod{m}$

Suppose that $a \equiv b \pmod{m}$ and $c \equiv d \pmod{m}$. Unrolling definitions gives us some k such that $a - b = km$, and some j such that $c - d = jm$.

Then, $a = km + b$ and $c = jm + d$. Multiplying both together gives us $ac = (km + b)(jm + d) = kjm^2 + kmd + bjm + bd$.

Re-arranging gives us $ac - bd = m(kjm + kd + bj)$.

Using the definition of congruence gives us $ac \equiv bd \pmod{m}$.

Example

Let n be an integer.

Prove that $n^2 \equiv 0 \pmod{4}$ or $n^2 \equiv 1 \pmod{4}$

Let's start by looking at a small example:

$$0^2 = 0 \equiv 0 \pmod{4}$$

$$1^2 = 1 \equiv 1 \pmod{4}$$

$$2^2 = 4 \equiv 0 \pmod{4}$$

$$3^2 = 9 \equiv 1 \pmod{4}$$

$$4^2 = 16 \equiv 0 \pmod{4}$$



Example

Let n be an integer.

Prove that $n^2 \equiv 0 \pmod{4}$ or $n^2 \equiv 1 \pmod{4}$

Case 1 (n is even):

Let's start by looking at a small example:

$$0^2 = 0 \equiv 0 \pmod{4}$$

$$1^2 = 1 \equiv 1 \pmod{4}$$

$$2^2 = 4 \equiv 0 \pmod{4}$$

$$3^2 = 9 \equiv 1 \pmod{4}$$

$$4^2 = 16 \equiv 0 \pmod{4}$$

Suppose n is even,
Then $n = 2k$ for some $k \in \mathbb{Z}$.

$$\text{So } n^2 = (2k)^2 = 4k^2,$$

which means

It looks like

Case 2 (n is odd): $n^2 \equiv 0 \pmod{4} \rightarrow n \equiv 0 \pmod{2} \rightarrow n^2 \equiv 0 \pmod{4}$, and
 $n \equiv 1 \pmod{2} \rightarrow n^2 \equiv 1 \pmod{4}$.

In this case we have

$$n = 2k + 1 \text{ for some } k \in \mathbb{Z}. \quad \text{So } n^2 = (2k + 1)^2$$

$$= 4k^2 + 2k + 2k + 1 = 4k^2 + 4k + 1$$

$$= 1 + 4(k^2 + k).$$

Example

Let n be an integer.

Prove that $n^2 \equiv 0 \pmod{4}$ or $n^2 \equiv 1 \pmod{4}$

Case 1 (n is even):

Suppose $n \equiv 0 \pmod{2}$.

Then, $n = 2k$ for some integer k .

So, $n^2 = (2k)^2 = 4k^2$. So, by

definition of congruence,

$n^2 \equiv 0 \pmod{4}$.

Let's start by looking at a small example:

$$0^2 = 0 \equiv 0 \pmod{4}$$

$$1^2 = 1 \equiv 1 \pmod{4}$$

$$2^2 = 4 \equiv 0 \pmod{4}$$

$$3^2 = 9 \equiv 1 \pmod{4}$$

$$4^2 = 16 \equiv 0 \pmod{4}$$

It looks like

Case 2 (n is odd):

Suppose $n \equiv 1 \pmod{2}$.

Then, $n = 2k + 1$ for some integer k .

So, $n^2 = (2k + 1)^2 = 4k^2 + 4k + 1 = 4(k^2 + k) + 1$.

So, by definition of congruence, $n^2 \equiv 1 \pmod{4}$.

$n \equiv 0 \pmod{2} \rightarrow n^2 \equiv 0 \pmod{4}$, and

$n \equiv 1 \pmod{2} \rightarrow n^2 \equiv 1 \pmod{4}$.

n-bit Unsigned Integer Representation

- Represent integer x as sum of powers of 2:

If $\sum_{i=0}^{n-1} b_i 2^i$ where each $b_i \in \{0,1\}$

then representation is $b_{n-1} \dots b_2 b_1 b_0$

$$99 = 64 + 32 + 2 + 1 \leftarrow$$

$$18 = 16 + 2$$

- For $n = 8$:

99: 0110 0011

18: 0001 0010

Sign-Magnitude Integer Representation

n -bit signed integers

Suppose that $-2^{n-1} < x < 2^{n-1}$

First bit as the sign, $n - 1$ bits for the value

$$99 = 64 + 32 + 2 + 1$$

$$18 = 16 + 2$$

For $n = 8$:

99: 0110 0011

-18: 1001 0010

0 6 - - 0
1 0 - - 0

Any problems with this representation?

Two's Complement Representation

n bit signed integers, first bit will still be the sign bit

Suppose that $0 \leq x < 2^{n-1}$,
 x is represented by the binary representation of x

Suppose that $0 \leq x \leq 2^{n-1}$,
 $-x$ is represented by the binary representation of $2^n - x$

$$2^n - 2^{n-1} = 2^{n-1}$$

Key property: Two's complement representation of any number y is equivalent to $y \bmod 2^n$ so arithmetic works **mod** 2^n

$$99 = 64 + 32 + 2 + 1$$

$$18 = 16 + 2$$

For $n = 8$:

$$99: \quad 0110 \ 0011$$

$$-18: \quad 1110 \ 1110$$

Sign-Magnitude vs. Two's Complement

0110

	-7	-6	-5	-4	-3	-2	-1	0	1	2	3	4	5	6	7
-	1111	1110	1101	1100	1011	1010	1001	0000	0001	0010	0011	0100	0101	0110	0111

← Sign-bit →

	-8	-7	-6	-5	-4	-3	-2	-1	0	1	2	3	4	5	6	7
	<u>1000</u>	<u>1001</u>	1010	1011	1100	1101	1110	1111	0000	0001	0010	0011	0100	0101	0110	0111

↑ Two's complement →

Two's Complement Representation

- For $0 < x \leq 2^{n-1}$, $-x$ is represented by the binary representation of $2^n - x$
 - That is, the two's complement representation of any number y has the same value as y modulo 2^n .
 - To compute this: Flip the bits of x then add 1:
 - All 1's string is $2^n - 1$, so
Flip the bits of $x \equiv$ replace x by $2^n - 1 - x$
Then add 1 to get $2^n - x$
- Handwritten notes:* $\sim x + 1$

Basic Applications of mod

- Hashing
- Pseudo random number generation
- Simple cipher

Hashing

Scenario:

Map a small number of data values from a large domain $\{0, 1, \dots, M - 1\}$...

...into a small set of locations $\{0, 1, \dots, n - 1\}$ so one can quickly check if some value is present

- $\text{hash}(x) = x \bmod p$ for p a prime close to n
 - or $\text{hash}(x) = (ax + b) \bmod p$
- Depends on all of the bits of the data
 - helps avoid collisions due to similar values
 - need to manage them if they occur

Pseudo-Random Number Generation

Linear Congruential method

$$x_{n+1} = (a x_n + c) \bmod m$$

Choose random x_0, a, c, m and produce a long sequence of x_n 's

Simple Ciphers

- **Caesar cipher**, $A = 1$, $B = 2$, . . .
 - HELLO WORLD
- **Shift cipher**
 - $f(p) = (p + k) \bmod 26$
 - $f^{-1}(p) = (p - k) \bmod 26$
- **More general**
 - $f(p) = (ap + b) \bmod 26$