


# CSE 311: Foundations of Computing

---

## Lecture 14: Induction



# Mathematical Induction

---

## Method for proving statements about all natural numbers

- A new logical inference rule!
  - It only applies over the natural numbers
  - The idea is to **use** the special structure of the naturals to prove things more easily
- Particularly useful for reasoning about programs!

```
for(int i=0; i < n; n++) { ... }
```

- Show  $P(i)$  holds after  $i$  times through the loop

```
public int f(int x) {  
    if (x == 0) { return 0; }  
    else { return f(x - 1) + 1; }  
}
```

- $f(x) = x$  for all values of  $x \geq 0$  naturally shown by induction.

**Prove**  $\forall a, b, m > 0 \forall k \in \mathbb{N} (a \equiv b \pmod{m} \rightarrow a^k \equiv b^k \pmod{m})$

---

**Let**  $a, b, m > 0 \in \mathbb{Z}$  **be arbitrary.** **Let**  $k \in \mathbb{N}$  **be arbitrary.**  
**Suppose that**  $a \equiv b \pmod{m}$ .

**We know**  $(a \equiv b \pmod{m} \wedge a \equiv b \pmod{m}) \rightarrow a^2 \equiv b^2 \pmod{m}$   
**by multiplying congruences. So, applying this repeatedly, we have:**

$$\begin{aligned} &(a \equiv b \pmod{m} \wedge a \equiv b \pmod{m}) \rightarrow a^2 \equiv b^2 \pmod{m} \\ &(a^2 \equiv b^2 \pmod{m} \wedge a \equiv b \pmod{m}) \rightarrow a^3 \equiv b^3 \pmod{m} \end{aligned}$$

...

$$\left( a^{k-1} \equiv b^{k-1} \pmod{m} \wedge a \equiv b \pmod{m} \right) \rightarrow a^k \equiv b^k \pmod{m}$$

The “...”s is a problem! We don’t have a proof rule that allows us to say “do this over and over”.

But there such a property of the natural numbers!

---

Domain: Natural Numbers

$$\frac{P(0) \quad \forall k (P(k) \rightarrow P(k + 1))}{\therefore \forall n P(n)}$$

# Induction Is A Rule of Inference

---

Domain: Natural Numbers

$$\begin{array}{c} P(0) \\ \hline \forall k (P(k) \rightarrow P(k + 1)) \\ \hline \therefore \forall n P(n) \end{array}$$

How do the givens prove  $P(5)$ ?


# Induction Is A Rule of Inference

---

Domain: Natural Numbers

$$\frac{P(0) \quad \forall k (P(k) \rightarrow P(k + 1))}{\therefore \forall n P(n)}$$

How do the givens prove P(5)?



First, we have **P(0)**.

Since  $P(k) \rightarrow P(k+1)$  for all  $k$ , we have **P(0) → P(1)**.

Since **P(0)** is true and **P(0) → P(1)**, by Modus Ponens, **P(1)** is true.

Since  $P(k) \rightarrow P(k+1)$  for all  $k$ , we have **P(1) → P(2)**.

Since **P(1)** is true and **P(1) → P(2)**, by Modus Ponens, **P(2)** is true.

## Using The Induction Rule In A Formal Proof

---

$$\frac{P(0) \quad \forall k (P(k) \rightarrow P(k + 1))}{\therefore \forall n P(n)}$$

# Using The Induction Rule In A Formal Proof

---

$$\begin{array}{c} P(0) \\ \forall k (P(k) \rightarrow P(k + 1)) \\ \hline \therefore \forall n P(n) \end{array}$$

1. Prove  $P(0)$

4.  $\forall k (P(k) \rightarrow P(k+1))$

5.  $\forall n P(n)$

Induction: 1, 4



# Using The Induction Rule In A Formal Proof

---

$$\frac{P(0) \quad \forall k (P(k) \rightarrow P(k + 1))}{\therefore \forall n P(n)}$$

1. Prove  $P(0)$
2. Let  $k$  be an arbitrary integer  $\geq 0$

3.  $P(k) \rightarrow P(k+1)$

4.  $\forall k (P(k) \rightarrow P(k+1))$

Intro  $\forall$ : 2, 3

5.  $\forall n P(n)$

Induction: 1, 4

# Using The Induction Rule In A Formal Proof

---

$$\begin{array}{c} P(0) \\ \forall k (P(k) \rightarrow P(k + 1)) \\ \hline \therefore \forall n P(n) \end{array}$$

1. Prove  $P(0)$
2. Let  $k$  be an arbitrary integer  $\geq 0$ 
  - 3.1. Assume that  $P(k)$  is true
  - 3.2. ...
  - 3.3. Prove  $P(k+1)$  is true
3.  $P(k) \rightarrow P(k+1)$       Direct Proof Rule
4.  $\forall k (P(k) \rightarrow P(k+1))$       Intro  $\forall$ : 2, 3
5.  $\forall n P(n)$       Induction: 1, 4

# Translating to an English Proof

---

$$\begin{array}{c} P(0) \\ \forall k (P(k) \rightarrow P(k + 1)) \\ \hline \therefore \forall n P(n) \end{array}$$

1. Prove  $P(0)$

**Base Case**

2. Let  $k$  be an arbitrary integer  $\geq 0$

**Inductive Hypothesis**

3.1. Assume that  $P(k)$  is true

3.2. ...

**Inductive Step**

3.3. Prove  $P(k+1)$  is true

3.  $P(k) \rightarrow P(k+1)$

Direct Proof Rule

4.  $\forall k (P(k) \rightarrow P(k+1))$

Intro  $\forall$ : 2, 3

5.  $\forall n P(n)$

Induction: 1, 4

**Conclusion**

# Translating To An English Proof

---

1. Prove $P(0)$	<b>Base Case</b>	
2. Let $k$ be an arbitrary integer $\geq 0$		<b>Inductive Hypothesis</b>
3.1. Assume that $P(k)$ is true		
3.2. ...		<b>Inductive Step</b>
3.3. Prove $P(k+1)$ is true		
3. $P(k) \rightarrow P(k+1)$		Direct Proof Rule
4. $\forall k (P(k) \rightarrow P(k+1))$		Intro $\forall$ : 2, 3
5. $\forall n P(n)$		Induction: 1, 4
		<b>Conclusion</b>

## Induction Proof Template

*[...Define  $P(n)$ ...]*

We will show that  $P(n)$  is true for every  $n \in \mathbb{N}$  by Induction.

**Base Case:** *[...proof of  $P(0)$  here...]*

**Induction Hypothesis:**

Suppose that  $P(k)$  is true for some  $k \in \mathbb{N}$ .

**Induction Step:**

We want to prove that  $P(k + 1)$  is true.

*[...proof of  $P(k + 1)$  here...]*

*The proof of  $P(k + 1)$  **must** invoke the IH somewhere.*

So, the claim is true by induction.

# Inductive Proofs In 5 Easy Steps

---

## Proof:

1. “Let  $P(n)$  be... . We will show that  $P(n)$  is true for every  $n \geq 0$  by Induction.”
2. “Base Case:” Prove  $P(0)$
3. “Inductive Hypothesis:  
Assume  $P(k)$  is true for some arbitrary integer  $k \geq 0$ ”
4. “Inductive Step:” Prove that  $P(k + 1)$  is true:  
*Use the goal to figure out what you need.*  
*Make sure you are using I.H. and point out where you are using it. (Don't assume  $P(k + 1)$  !!)*
5. “Conclusion: Result follows by induction”

## What is $1 + 2 + 4 + \dots + 2^n$ ?

---

- $1 = 1$
- $1 + 2 = 3$
- $1 + 2 + 4 = 7$
- $1 + 2 + 4 + 8 = 15$
- $1 + 2 + 4 + 8 + 16 = 31$

It sure looks like this sum is  $2^{n+1} - 1$

How can we prove it?

We could prove it for  $n = 1, n = 2, n = 3, \dots$  but that would literally take forever.

Good that we have induction!

**Prove**  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$

---

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

- 1. Let  $P(n)$  be " $1 + 2 + \dots + 2^n = 2^{n+1} - 1$ ". We will show  $P(n)$  is true for all natural numbers by induction.**



**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

- 1. Let  $P(n)$  be “ $1 + 2 + \dots + 2^n = 2^{n+1} - 1$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.**

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

- 1. Let  $P(n)$  be “ $1 + 2 + \dots + 2^n = 2^{n+1} - 1$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ .**

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

- 1. Let  $P(n)$  be “ $1 + 2 + \dots + 2^n = 2^{n+1} - 1$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ .**
- 4. Induction Step:**

**Goal: Show  $P(k+1)$ , i.e. show  $1 + 2 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$**

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

1. Let  $P(n)$  be " $1 + 2 + \dots + 2^n = 2^{n+1} - 1$ ". We will show  $P(n)$  is true for all natural numbers by induction.
2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.
3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ .

4. Induction Step:

**Goal: Show  $P(k+1)$ , i.e. show  $1 + 2 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$**

$$1 + 2 + \dots + 2^k = 2^{k+1} - 1 \quad \text{by IH}$$

Adding  $2^{k+1}$  to both sides, we get:

$$1 + 2 + \dots + 2^k + 2^{k+1} = 2^{k+1} + 2^{k+1} - 1$$

Note that  $2^{k+1} + 2^{k+1} = 2(2^{k+1}) = 2^{k+2}$ .

So, we have  $1 + 2 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$ , which is exactly  $P(k+1)$ .

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

1. Let  $P(n)$  be " $1 + 2 + \dots + 2^n = 2^{n+1} - 1$ ". We will show  $P(n)$  is true for all natural numbers by induction.
2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.
3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ .

4. Induction Step:

**Goal: Show  $P(k+1)$ , i.e. show  $1 + 2 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$**

$$\begin{aligned} 1 + 2 + \dots + 2^k + 2^{k+1} &= (1 + 2 + \dots + 2^k) + 2^{k+1} \\ &= 2^{k+1} - 1 + 2^{k+1} \quad \text{by the IH} \end{aligned}$$

Note that  $2^{k+1} + 2^{k+1} = 2(2^{k+1}) = 2^{k+2}$ .

So, we have  $1 + 2 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$ , which is exactly  $P(k+1)$ .

**Alternative way of writing the inductive step**

**Prove  $1 + 2 + 4 + \dots + 2^n = 2^{n+1} - 1$**

---

1. Let  $P(n)$  be " $1 + 2 + \dots + 2^n = 2^{n+1} - 1$ ". We will show  $P(n)$  is true for all natural numbers by induction.
2. Base Case ( $n=0$ ):  $2^0 = 1 = 2 - 1 = 2^{0+1} - 1$  so  $P(0)$  is true.
3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ .

4. Induction Step:

**Goal: Show  $P(k+1)$ , i.e. show  $1 + 2 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$**

$$\begin{aligned} 1 + 2 + \dots + 2^k + 2^{k+1} &= (1 + 2 + \dots + 2^k) + 2^{k+1} \\ &= 2^{k+1} - 1 + 2^{k+1} \quad \text{by the IH} \end{aligned}$$

Note that  $2^{k+1} + 2^{k+1} = 2(2^{k+1}) = 2^{k+2}$ .

So, we have  $1 + 2 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$ , which is exactly  $P(k+1)$ .

5. Thus  $P(n)$  is true for all  $n \in \mathbb{N}$ , by induction.

**Prove**  $1 + 2 + 3 + \dots + n = n(n + 1)/2$

---

**Prove  $1 + 2 + 3 + \dots + n = n(n + 1)/2$**

---

- 1. Let  $P(n)$  be " $0 + 1 + 2 + \dots + n = n(n+1)/2$ ". We will show  $P(n)$  is true for all natural numbers by induction.**



**Prove  $1 + 2 + 3 + \dots + n = n(n + 1)/2$**

---

- 1. Let  $P(n)$  be “ $0 + 1 + 2 + \dots + n = n(n+1)/2$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $0 = 0(0+1)/2$ . Therefore  $P(0)$  is true.**

**Prove  $1 + 2 + 3 + \dots + n = n(n + 1)/2$**

---

- 1. Let  $P(n)$  be “ $0 + 1 + 2 + \dots + n = n(n+1)/2$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):  $0 = 0(0+1)/2$ . Therefore  $P(0)$  is true.**
- 3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ .**
- 4. Induction Step:**

**Goal: Show  $P(k+1)$ , i.e. show  $1 + 2 + \dots + k + (k+1) = (k+1)(k+2)/2$**

## Prove $1 + 2 + 3 + \dots + n = n(n + 1)/2$

---

1. Let  $P(n)$  be " $0 + 1 + 2 + \dots + n = n(n+1)/2$ ". We will show  $P(n)$  is true for all natural numbers by induction.
2. Base Case ( $n=0$ ):  $0 = 0(0+1)/2$ . Therefore  $P(0)$  is true.
3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ .

### 4. Induction Step:

**Goal: Show  $P(k+1)$ , i.e. show  $1 + 2 + \dots + k + (k+1) = (k+1)(k+2)/2$**

$$\begin{aligned} 1 + 2 + \dots + k + (k+1) &= (1 + 2 + \dots + k) + (k+1) \\ &= k(k+1)/2 + (k+1) \text{ by IH} \end{aligned}$$

Now  $k(k+1)/2 + (k+1) = (k+1)(k/2 + 1) = (k+1)(k+2)/2$ .

So, we have  $1 + 2 + \dots + k + (k+1) = (k+1)(k+2)/2$ , which is exactly  $P(k+1)$ .

5. Thus  $P(n)$  is true for all  $n \in \mathbb{N}$ , by induction.

## Another example of a pattern

---

- $2^0 - 1 = 1 - 1 = 0 = 3 \cdot 0$
- $2^2 - 1 = 4 - 1 = 3 = 3 \cdot 1$
- $2^4 - 1 = 16 - 1 = 15 = 3 \cdot 5$
- $2^6 - 1 = 64 - 1 = 63 = 3 \cdot 21$
- $2^8 - 1 = 256 - 1 = 255 = 3 \cdot 85$
- ...

**Prove:  $3 \mid (2^{2n} - 1)$  for all  $n \geq 0$**

---

**Prove:  $3 \mid (2^{2n} - 1)$  for all  $n \geq 0$**

---

- 1. Let  $P(n)$  be “ $3 \mid (2^{2n} - 1)$ ”. We will show  $P(n)$  is true for all natural numbers by induction.**
- 2. Base Case ( $n=0$ ):**

## Prove: $3 \mid (2^{2n} - 1)$ for all $n \geq 0$

---

1. Let  $P(n)$  be " $3 \mid (2^{2n} - 1)$ ". We will show  $P(n)$  is true for all natural numbers by induction.
2. Base Case ( $n=0$ ):  $2^{2 \cdot 0} - 1 = 1 - 1 = 0 = 3 \cdot 0$  Therefore  $P(0)$  is true.
3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ .
4. Induction Step:  

**Goal: Show  $P(k+1)$ , i.e. show  $3 \mid (2^{2(k+1)} - 1)$**

## Prove: $3 \mid (2^{2n} - 1)$ for all $n \geq 0$

---

1. Let  $P(n)$  be " $3 \mid (2^{2n} - 1)$ ". We will show  $P(n)$  is true for all natural numbers by induction.
2. Base Case ( $n=0$ ):  $2^{2 \cdot 0} - 1 = 1 - 1 = 0 = 3 \cdot 0$  Therefore  $P(0)$  is true.
3. Induction Hypothesis: Suppose that  $P(k)$  is true for some arbitrary integer  $k \geq 0$ .

### 4. Induction Step:

Goal: Show  $P(k+1)$ , i.e. show  $3 \mid (2^{2(k+1)} - 1)$

By IH,  $3 \mid (2^{2k} - 1)$  so  $2^{2k} - 1 = 3j$  for some integer  $j$

$$\begin{aligned} \text{So } 2^{2(k+1)} - 1 &= 2^{2k+2} - 1 = 4(2^{2k}) - 1 = 4(3j+1) - 1 \\ &= 12j+3 = 3(4j+1) \end{aligned}$$

Therefore  $3 \mid (2^{2(k+1)} - 1)$  which is exactly  $P(k+1)$ .



5. Thus  $P(n)$  is true for all  $n \in \mathbb{N}$ , by induction.



# Checkerboard Tiling

---

- Prove that a  $2^n \times 2^n$  checkerboard with one square removed can be tiled with:



# Checkerboard Tiling

---

1. Let  $P(n)$  be any  $2^n \times 2^n$  checkerboard with one square removed can be tiled with  .

We prove  $P(n)$  for all  $n \geq 1$  by induction on  $n$ .

2. Base Case:  $n=1$     

# Checkerboard Tiling

---



1. Let  $P(n)$  be any  $2^n \times 2^n$  checkerboard with one square removed can be tiled with  .

We prove  $P(n)$  for all  $n \geq 1$  by induction on  $n$ .

2. Base Case:  $n=1$     

3. Inductive Hypothesis: Assume  $P(k)$  for some arbitrary integer  $k \geq 1$

4. Inductive Step: Prove  $P(k+1)$



Apply IH to each quadrant then fill with extra tile.