

CSE 311 Section 3

Quantifiers and Proofs

Administrivia & Introductions



Announcements & Reminders

- HW1
 - If you think something was graded incorrectly, submit a regrade request!
- HW2 is due TOMORROW 10/13 @ 10PM on Gradescope
 - Use late days if you need them!
- HW3
 - Due Friday 10/20 @ 10pm

References

- Helpful reference sheets can be found on the course website!
 - <https://courses.cs.washington.edu/courses/cse311/23au/resources/>
- How to LaTeX (found on Assignments page of website):
 - <https://courses.cs.washington.edu/courses/cse311/23au/assignments/HowToLaTeX.pdf>
- Equivalence Reference Sheet
 - https://courses.cs.washington.edu/courses/cse311/23au/resources/reference-logical_equiv.pdf
 - <https://courses.cs.washington.edu/courses/cse311/23au/resources/logicalConnectPoster.pdf>
- Boolean Algebra Reference Sheet
 - <https://courses.cs.washington.edu/courses/cse311/23au/resources/reference-boolean-alg.pdf>
- Plus more!

Predicates & Quantifiers



Predicates & Quantifiers Review

- **Predicate:** a function that outputs true or false
 - $\text{Cat}(x) := \text{“}x \text{ is a cat”}$
 - $\text{LessThan}(x, y) := \text{“}x < y\text{”}$
- **Domain of Discourse:** the types of inputs allowed in predicates
 - Numbers, mammals, cats and dogs, people in this class, etc.
- **Quantifiers**
 - Universal Quantifier $\forall x$: for all x , for every x
 - Existential Quantifier $\exists x$: there is an x , there exists an x , for some x
- **Domain Restriction**
 - Universal Quantifier $\forall x$: add the restriction as the hypothesis to an **implication**
 - Existential Quantifier $\exists x$: **AND** in the restriction

Problem 1 – Domain Restriction

Translate each of the following sentences into logical notation. These translations require some of our quantifier tricks. You may use the operators $+$ and \cdot which take two numbers as input and evaluate to their sum or product, respectively.

- a) Domain: Positive integers; Predicates: *Even, Prime, Equal*
“There is only one positive integer that is prime and even.”
- b) Domain: Real numbers; Predicates: *Even, Prime, Equal*
“There are two different prime numbers that sum to an even number.”
- c) Domain: Real numbers; Predicates: *Even, Prime, Equal*
“The product of two distinct prime numbers is not prime.”
- d) Domain: Real numbers; Predicates: *Even, Prime, Equal, Positive, Greater, Integer*
“For every positive integer, there is a greater even integer”

Work on parts (a) and (b) with the people around you, and then we'll go over it together!

Problem 1 – Domain Restriction

- a) Domain: Positive integers; Predicates: *Even*, *Prime*, *Equal*
“There is only one positive integer that is prime and even.”

Problem 1 – Domain Restriction

- a) Domain: Positive integers; Predicates: *Even*, *Prime*, *Equal*
“There is only one positive integer that is prime and even.”

We can start out with:

$$\exists x(\text{Prime}(x) \wedge \text{Even}(x))$$

Problem 1 – Domain Restriction

- a) Domain: Positive integers; Predicates: *Even*, *Prime*, *Equal*
“There is only one positive integer that is prime and even.”

We can start out with:

$$\exists x(\text{Prime}(x) \wedge \text{Even}(x))$$

But now we need to add in the restriction that this x is the ONLY positive integer that is prime and even. This is a technique you'll use whenever you need to have only one of something:

$$\exists x(\text{Prime}(x) \wedge \text{Even}(x) \wedge \forall y[\neg \text{Equal}(x, y) \rightarrow \neg(\text{Even}(y) \wedge \text{Prime}(y))])$$

Or, we could use the contrapositive:

$$\exists x(\text{Prime}(x) \wedge \text{Even}(x) \wedge \forall y[(\text{Even}(y) \wedge \text{Prime}(y)) \rightarrow \text{Equal}(x, y)])$$

Problem 1 – Domain Restriction

- b) Domain: Real numbers; Predicates: *Even*, *Prime*, *Equal*
“There are two different prime numbers that sum to an even number.”

Problem 1 – Domain Restriction

b) Domain: Real numbers; Predicates: *Even*, *Prime*, *Equal*

“There are two different prime numbers that sum to an even number.”

Seems like maybe we should be able to say something like:

$$\exists x \exists y (\text{Prime}(x) \wedge \text{Prime}(y) \wedge \text{Even}(x + y))$$

Problem 1 – Domain Restriction

- b) Domain: Real numbers; Predicates: *Even*, *Prime*, *Equal*
“There are two different prime numbers that sum to an even number.”

Seems like maybe we should be able to say something like:

$$\exists x \exists y (\text{Prime}(x) \wedge \text{Prime}(y) \wedge \text{Even}(x + y))$$

But this leaves open the possibility of x and y being equal (so they won't be two DIFFERENT numbers). So, we need to explicitly add in that x and y are not equal:

$$\exists x \exists y (\text{Prime}(x) \wedge \text{Prime}(y) \wedge \text{Even}(x + y) \wedge \neg \text{Equal}(x, y))$$

Problem 2 – ctrl-z

Translate these logical expressions to English. For each of the translations, assume that domain restriction is being used and take that into account in your English versions.

Let your domain be all UW Students. Predicates $143Student(x)$ and $311Student(x)$ mean the student is in CSE 143 and 311, respectively. $BioMajor(x)$ means x is a bio major, $DidHomeworkOne(x)$ means the student did homework 1 (of 311). Finally, $KnowsJava(x)$ and $KnowsDeMorgan(x)$ mean x knows Java and knows DeMorgan's Laws, respectively.

- a) $\forall x(143Student(x) \rightarrow KnowsJava(x))$
- b) $\exists x(143Student(x) \wedge BioMajor(x))$
- c) $\forall x([311Student(x) \wedge DidHomeworkOne(x)] \rightarrow KnowsDeMorgan(x))$

Work on parts (a) and (c) with the people around you, and then we'll go over it together!

Problem 2 – ctrl-z

a) $\forall x(143\text{Student}(x) \rightarrow \text{KnowsJava}(x))$

Problem 2 – ctrl-z

a) $\forall x(143\text{Student}(x) \rightarrow \text{KnowsJava}(x))$

Every 143 student knows java.

“If a UW student is a 143 student, then they know java” is a valid translation of the original sentence, but it is not taking advantage of the domain restriction.

Problem 2 – ctrl-z

c) $\forall x([311Student(x) \wedge DidHomeworkOne(x)] \rightarrow KnowsDeMorgan(x))$

Problem 2 – ctrl-z

c) $\forall x([311\text{Student}(x) \wedge \text{DidHomeworkOne}(x)] \rightarrow \text{KnowsDeMorgan}(x))$

All 311 students who do Homework 1 know DeMorgan's Laws.

“If a UW student is a 311 student and they did Homework 1, then they know DeMorgan's Laws” is a valid translation of the original sentence, but it is not taking advantage of the domain restriction.

Problem 3 – Quantifier Switch

Consider the following pairs of sentences. For each pair, determine if one implies the other, if they are equivalent, or neither.

a) $\forall x \forall y P(x, y)$ $\forall y \forall x P(x, y)$

b) $\exists x \exists y P(x, y)$ $\exists y \exists x P(x, y)$

c) $\forall x \exists y P(x, y)$ $\forall y \exists x P(x, y)$

d) $\forall x \exists y P(x, y)$ $\exists x \forall y P(x, y)$

e) $\forall x \exists y P(x, y)$ $\exists y \forall x P(x, y)$

Work on parts (d) and (e) with the people around you, and then we'll go over it together!

Problem 3 – Quantifier Switch

d) $\forall x \exists y P(x, y)$ $\exists x \forall y P(x, y)$

Problem 3 – Quantifier Switch

d) $\forall x \exists y P(x, y)$ $\exists x \forall y P(x, y)$

Different

For all x , there is a y vs. there exists an x that, for all y

Everyone likes someone vs. someone likes everyone

Problem 3 – Quantifier Switch

e) $\forall x \exists y P(x, y)$ $\exists y \forall x P(x, y)$

Problem 3 – Quantifier Switch

e) $\forall x \exists y P(x, y)$ $\exists y \forall x P(x, y)$

The second implies the first


For all x, there is a y, vs. there exists a y that, for all x

The second is stronger since a specific y must work for all x whereas the for the first, the y value does not have to be the same for every x

Everyone likes someone

vs.

There is someone who is liked by everyone



Direct Proofs



Direct Proofs

- Very common form of proof, sometimes written as a symbolic proof and sometimes written as an English proof
- Use direct proofs to prove implications
- Steps to prove $p \rightarrow q$
 - Assume p is true
 - Write down all the facts we know (including p)
 - Combine the things we know to derive new facts
 - Continue until we directly show q is true

Writing a Proof (symbolically or in English)

- Don't just jump right in!
- Look at the **claim**, and make sure you know:
 - What every word in the claim means
 - What the claim as a whole means
- Translate the claim in predicate logic.
- Next, write down the **Proof Skeleton**:
 - Where to start
 - What your target is
- Then once you know what claim you are proving and your starting point and ending point, you can finally write the proof!

Helpful Tips for English Proofs

- Start by introducing your assumptions
 - Introduce variables with “let”
 - “Let x be an arbitrary prime number...”
 - Introduce assumptions with “suppose”
 - “Suppose that $y \in A \wedge y \notin B...$ ”
- When you supply a value for an existence proof, use “Consider”
 - “Consider $x = 2...$ ”
- **ALWAYS** state what type your variable is (integer, set, etc.)
- Universal Quantifier means variable must be arbitrary
- Existential Quantifier means variable can be specific

Problem 6 – Direct Proof

- a) Let the domain of discourse be integers. Define the predicates $\text{Odd}(x) := \exists k(x = 2k + 1)$, and $\text{Even}(x) := \exists k(x = 2k)$. Translate the following claim into predicate logic:

The sum of an even and odd integer is odd.

- b) Prove that the claim holds.

Problem 6 – Direct Proof

- a) Let the domain of discourse be integers. Define the predicates $\text{Odd}(x) := \exists k(x = 2k + 1)$, and $\text{Even}(x) := \exists k(x = 2k)$. Translate the following claim into predicate logic:

The sum of an even and odd integer is odd.

Work on part (a) of this problem with the people around you, and then we'll go over it together!

Problem 6 – Direct Proof

- a) Let the domain of discourse be integers. Define the predicates $\text{Odd}(x) := \exists k(x = 2k + 1)$, and $\text{Even}(x) := \exists k(x = 2k)$. Translate the following claim into predicate logic:

The sum of an even and odd integer is odd.

Problem 6 – Direct Proof

- a) Let the domain of discourse be integers. Define the predicates $\text{Odd}(x) := \exists k(x = 2k + 1)$, and $\text{Even}(x) := \exists k(x = 2k)$. Translate the following claim into predicate logic:

The sum of an even and odd integer is odd.

$$\forall n \forall m ((\text{Even}(n) \wedge \text{Odd}(m)) \rightarrow \text{Odd}(n + m))$$

Problem 6 – Direct Proof

- a) Let the domain of discourse be integers. Define the predicates $\text{Odd}(x) := \exists k(x = 2k + 1)$, and $\text{Even}(x) := \exists k(x = 2k)$. Translate the following claim into predicate logic:

The sum of an even and odd integer is odd.

- b) Prove that the claim holds.

Lets walk through part (b) together!

Problem 6 – Direct Proof

b) Prove that the claim holds.

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n \forall m ((\text{Even}(n) \wedge \text{Odd}(m)) \rightarrow \text{Odd}(n + m))$$

Problem 6 – Direct Proof

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n \forall m ((\text{Even}(n) \wedge \text{Odd}(m)) \rightarrow \text{Odd}(n + m))$$

b) Prove that the claim holds.

Let n and m be arbitrary integers.

...

Since n and m were arbitrary, the sum of any even and odd integer is odd.

Problem 6 – Direct Proof

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n \forall m ((\text{Even}(n) \wedge \text{Odd}(m)) \rightarrow \text{Odd}(n + m))$$

b) Prove that the claim holds.

Let n and m be arbitrary integers. Suppose n is even and m is odd.

...

Thus by (some reasoning here), $n + m$ is odd. Since n and m were arbitrary, the sum of any even and odd integer is odd.

Problem 6 – Direct Proof

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n \forall m ((\text{Even}(n) \wedge \text{Odd}(m)) \rightarrow \text{Odd}(n + m))$$

b) Prove that the claim holds.

Let n and m be arbitrary integers. Suppose n is even and m is odd. Then by definition of even, $n = 2k$ for some integer k . By definition of odd, $m = 2j + 1$ for some integer j .

...

Then $n + m$ is 2 times an integer plus 1. Thus by definition of odd, $n + m$ is odd. Since n and m were arbitrary, the sum of any even and odd integer is odd.

Problem 6 – Direct Proof

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n \forall m ((\text{Even}(n) \wedge \text{Odd}(m)) \rightarrow \text{Odd}(n + m))$$

b) Prove that the claim holds.

Let n and m be arbitrary integers. Suppose n is even and m is odd. Then by definition of even, $n = 2k$ for some integer k . By definition of odd, $m = 2j + 1$ for some integer j .

Then consider $n + m$:

$$n + m = 2k + 2j + 1$$

...

Then $n + m$ is 2 times an integer plus 1. Thus by definition of odd, $n + m$ is odd. Since n and m were arbitrary, the sum of any even and odd integer is odd.

Problem 6 – Direct Proof

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n \forall m ((\text{Even}(n) \wedge \text{Odd}(m)) \rightarrow \text{Odd}(n + m))$$

b) Prove that the claim holds.

Let n and m be arbitrary integers. Suppose n is even and m is odd. Then by definition of even, $n = 2k$ for some integer k . By definition of odd, $m = 2j + 1$ for some integer j .

Then consider $n + m$:

$$\begin{aligned} n + m &= 2k + 2j + 1 \\ &= 2(k + j) + 1 \end{aligned}$$

...

Then $n + m$ is 2 times an integer plus 1. Thus by definition of odd, $n + m$ is odd. Since n and m were arbitrary, the sum of any even and odd integer is odd.

Problem 6 – Direct Proof

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n \forall m ((\text{Even}(n) \wedge \text{Odd}(m)) \rightarrow \text{Odd}(n + m))$$

b) Prove that the claim holds.

Let n and m be arbitrary integers. Suppose n is even and m is odd. Then by definition of even, $n = 2k$ for some integer k . By definition of odd, $m = 2j + 1$ for some integer j .

Then consider $n + m$:

$$\begin{aligned} n + m &= 2k + 2j + 1 \\ &= 2(k + j) + 1 \end{aligned}$$

Since k and j are integers, $k + j$ is an integer.

Then $n + m$ is 2 times an integer plus 1. Thus by definition of odd, $n + m$ is odd.

Since n and m were arbitrary, the sum of any even and odd integer is odd.

Problem 7 – Proof of Biconditional

- a) Let the domain of discourse be integers. Define the predicates $\text{Odd}(x) := \exists k(x = 2k + 1)$, and $\text{Even}(x) := \exists k(x = 2k)$. Translate the following claim into predicate logic:

For all integers n , $n - 4$ is even if and only if $n + 17$ is odd.

- b) Prove that the claim holds.

Problem 7 – Proof of Biconditional

- a) Let the domain of discourse be integers. Define the predicates $\text{Odd}(x) := \exists k(x = 2k + 1)$, and $\text{Even}(x) := \exists k(x = 2k)$. Translate the following claim into predicate logic:

For all integers n , $n - 4$ is even if and only if $n + 17$ is odd.

Work on part (a) of this problem with the people around you, and then we'll go over it together!

Problem 7 – Proof of Biconditional

- a) Let the domain of discourse be integers. Define the predicates $\text{Odd}(x) := \exists k(x = 2k + 1)$, and $\text{Even}(x) := \exists k(x = 2k)$. Translate the following claim into predicate logic:

For all integers n , $n - 4$ is even if and only if $n + 17$ is odd.

$$\forall n(\text{Even}(n - 4) \leftrightarrow \text{Odd}(n + 17))$$

Problem 7 – Proof of Biconditional

- a) Let the domain of discourse be integers. Define the predicates $\text{Odd}(x) := \exists k(x = 2k + 1)$, and $\text{Even}(x) := \exists k(x = 2k)$. Translate the following claim into predicate logic:

For all integers n , $n - 4$ is even if and only if $n + 17$ is odd.

- b) Prove that the claim holds.

Lets walk through part (b) together!

Problem 7 – Proof of Biconditional

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n(\text{Even}(n - 4) \leftrightarrow \text{Odd}(n + 17))$$

b) Prove that the claim holds.

Problem 7 – Proof of Biconditional

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n(\text{Even}(n - 4) \leftrightarrow \text{Odd}(n + 17))$$

b) Prove that the claim holds.

We know that a biconditional $p \leftrightarrow q$ can be equivalently expressed as two implications anded together: $p \rightarrow q \wedge q \rightarrow p$. So, in order to prove a biconditional, we need to prove both implications hold.

For this problem, we need to prove both the forward direction:

$$\forall n(\text{Even}(n - 4) \rightarrow \text{Odd}(n + 17))$$

And the backward direction:

$$\forall n(\text{Odd}(n + 17) \rightarrow \text{Even}(n - 4))$$

By showing both implications hold, we prove that the biconditional holds.

Problem 7 – Proof of Biconditional

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n(\text{Even}(n - 4) \leftrightarrow \text{Odd}(n + 17))$$

b) Prove that the claim holds.

⇒ Let n be an arbitrary integer.

...

Since n was arbitrary, we have shown that for all integers n that if $n - 4$ is even, then $n + 17$ is odd.

Problem 7 – Proof of Biconditional

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n(\text{Even}(n - 4) \leftrightarrow \text{Odd}(n + 17))$$

b) Prove that the claim holds.

⇒ Let n be an arbitrary integer. Suppose that $n - 4$ is even. Then by definition of even, $n - 4 = 2k$ for some integer k . Then observe that:

$$n - 4 = 2k$$

$$n + 17 = 2k + 21 \quad \text{Adding 21 to both sides}$$

$$n + 17 = 2(k + 10) + 1 \quad \text{Factoring}$$

Thus $n + 17 = 2(k + 10) + 1$.

Since k is an integer, $k + 10$ is an integer. So $n + 17$ is 2 times an integer plus 1. Thus by definition of odd, $n + 17$ is odd.

Since n was arbitrary, we have shown that for all integers n that if $n - 4$ is even, then $n + 17$ is odd.

Problem 7 – Proof of Biconditional

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n(\text{Even}(n - 4) \leftrightarrow \text{Odd}(n + 17))$$

b) Prove that the claim holds.

← Let n be an arbitrary integer.

...

Since n was arbitrary, we have shown that for all integers n , if $n + 17$ is odd, then $n - 4$ is even.

Problem 7 – Proof of Biconditional

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n(\text{Even}(n - 4) \leftrightarrow \text{Odd}(n + 17))$$

b) Prove that the claim holds.

\Leftarrow Let n be an arbitrary integer. Suppose $n + 17$ is odd. Then by definition of odd, $n + 17 = 2k + 1$ for some integer k . Then observe that:

$$n + 17 = 2k + 1$$

$$n - 4 = 2k + 1 - 21$$

Subtracting 21 from both sides

$$n - 4 = 2(k - 10)$$

Factoring

Thus $n - 4 = 2(k - 10)$.

Since k is an integer, $k - 10$ is an integer. So $n - 4$ is 2 times an integer.

So by definition of even, $n - 4$ is even. Since n was arbitrary, we have shown that for all integers n , if $n + 17$ is odd, then $n - 4$ is even.

Problem 7 – Proof of Biconditional

$$\text{Odd}(x) := \exists k(x = 2k + 1)$$

$$\text{Even}(x) := \exists k(x = 2k)$$

Claim:

$$\forall n(\text{Even}(n - 4) \leftrightarrow \text{Odd}(n + 17))$$

b) Prove that the claim holds.

\Rightarrow Let n be an arbitrary integer. Suppose that $n - 4$ is even. Then by definition of even, $n - 4 = 2k$ for some integer k . Then observe that:

$$n - 4 = 2k$$

$$n + 17 = 2k + 21 \quad \text{Adding 21 to both sides}$$

$$n + 17 = 2(k + 10) + 1 \quad \text{Factoring}$$

Thus $n + 17 = 2(k + 10) + 1$. Since k is an integer, $k + 10$ is an integer. So $n + 17$ is 2 times an integer plus 1. Thus by definition of odd, $n + 17$ is odd. Since n was arbitrary, we have shown that for all integers n that if $n - 4$ is even, then $n + 17$ is odd.

\Leftarrow Let n be an arbitrary integer. Suppose $n + 17$ is odd. Then by definition of odd, $n + 17 = 2k + 1$ for some integer k . Then observe that:

$$n + 17 = 2k + 1$$

$$n - 4 = 2k + 1 - 21 \quad \text{Subtracting 21 from both sides}$$

$$n - 4 = 2(k - 10) \quad \text{Factoring}$$

Thus $n - 4 = 2(k - 10)$. Since k is an integer, $k - 10$ is an integer. So $n - 4$ is 2 times an integer. So by definition of even, $n - 4$ is even. Since n was arbitrary, we have shown that for all integers n , if $n + 17$ is odd, then $n - 4$ is even.

That's All, Folks!

Thanks for coming to section this week!
Any questions?