

CSE 312

Autumn 2012

MLE: Maximum Likelihood Estimators

Reminder: population or distribution versus sample

Population

mean $\mu = \sum_{1 \leq i \leq 6} ip_i$ $\mu = \int_{\mathbb{R}} xf(x)dx$

variance $\sigma^2 = \sum_{1 \leq i \leq 6} (i - \mu)^2 p_i$ $\sigma^2 = \int_{\mathbb{R}} (x - \mu)^2 f(x)dx$

Sample

mean $\bar{x} = \sum_{1 \leq i \leq n} x_i/n$

variance $\bar{s}^2 = \sum_{1 \leq i \leq n} (x_i - \bar{x})^2/n$

Learning From Data: MLE

Maximum Likelihood Estimators

Parameter Estimation

Assuming sample x_1, x_2, \dots, x_n is from a parametric distribution $f(x|\theta)$, estimate θ .

E.g.: Given sample HHTTTTTHTHTTTHH of (possibly biased) coin flips, estimate

$\theta =$ probability of Heads

$f(x|\theta)$ is the Bernoulli probability mass function with parameter θ

Likelihood

$P(x | \theta)$: Probability of event x given *model* θ

Viewed as a function of x (fixed θ), it's a *probability*

E.g., $\sum_x P(x | \theta) = 1$

Viewed as a function of θ (fixed x), it's a *likelihood*

E.g., $\sum_{\theta} P(x | \theta)$ can be anything; *relative* values of interest.

E.g., if θ = prob of heads in a sequence of coin flips then

$$P(\text{HHTHH} | .6) > P(\text{HHTHH} | .5),$$


I.e., event HHTHH is *more likely* when $\theta = .6$ than $\theta = .5$

And **what θ make HHTHH *most likely*?**

Likelihood Function

$P(\text{HHTHH} \mid \theta)$:
Probability of HHTHH,
given $P(H) = \theta$:

θ	$\theta^4(1-\theta)$
0.2	0.0013
0.5	0.0313
0.8	0.0819
0.95	0.0407



Maximum Likelihood Parameter Estimation

One (of many) approaches to param. est.

Likelihood of (indp) observations x_1, x_2, \dots, x_n

$$L(x_1, x_2, \dots, x_n \mid \theta) = \prod_{i=1}^n f(x_i \mid \theta)$$

As a function of θ , what θ maximizes the likelihood of the data actually observed

Typical approach: $\frac{\partial}{\partial \theta} L(\vec{x} \mid \theta) = 0$ or $\frac{\partial}{\partial \theta} \log L(\vec{x} \mid \theta) = 0$

Example 1

n coin flips, x_1, x_2, \dots, x_n ; n_0 tails, n_1 heads, $n_0 + n_1 = n$;

θ = probability of heads

$$L(x_1, x_2, \dots, x_n \mid \theta) = (1 - \theta)^{n_0} \theta^{n_1}$$


$$\log L(x_1, x_2, \dots, x_n \mid \theta) = n_0 \log(1 - \theta) + n_1 \log \theta$$

$$\frac{\partial}{\partial \theta} \log L(x_1, x_2, \dots, x_n \mid \theta) = \frac{-n_0}{1 - \theta} + \frac{n_1}{\theta}$$

Setting to zero and solving:

$$\hat{\theta} = \frac{n_1}{n}$$

Observed fraction of
successes in *sample* is
MLE of success
probability in *population*



(Also verify it's max, not min, & not better on boundary)


Parameter Estimation

Assuming sample x_1, x_2, \dots, x_n is from a parametric distribution $f(x|\theta)$, estimate θ .


E.g.: Given n normal samples, estimate mean & variance

$$f(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-(x-\mu)^2 / (2\sigma^2)}$$

$$\theta = (\mu, \sigma^2)$$




Ex2: I got data; a little birdie tells me
it's normal, and promises $\sigma^2 = 1$




Which is more likely: (a) this?

μ unknown, $\sigma^2 = 1$




Which is more likely: (b) or this?

μ unknown, $\sigma^2 = 1$



Which is more likely: (c) or *this*?


μ unknown, $\sigma^2 = 1$



Which is more likely: (c) or this?

μ unknown, $\sigma^2 = 1$

Looks good by eye, but how do I optimize my estimate of μ ?



Ex. 2: $x_i \sim N(\mu, \sigma^2)$, $\sigma^2 = 1$, μ unknown


$$L(x_1, x_2, \dots, x_n | \theta) = \prod_{1 \leq i \leq n} \frac{1}{\sqrt{2\pi}} e^{-(x_i - \theta)^2 / 2}$$

$$\ln L(x_1, x_2, \dots, x_n | \theta) = \sum_{1 \leq i \leq n} -\frac{1}{2} \ln 2\pi - \frac{(x_i - \theta)^2}{2}$$

$$\frac{d}{d\theta} \ln L(x_1, x_2, \dots, x_n | \theta) = \sum_{1 \leq i \leq n} (x_i - \theta)$$

$$= \left(\sum_{1 \leq i \leq n} x_i \right) - n\theta = 0$$

And verify it's max,
not min & not better
on boundary



$$\hat{\theta} = \left(\sum_{1 \leq i \leq n} x_i \right) / n = \bar{x}$$

**Sample mean is MLE of
population mean**

Hmm ..., density \neq probability


So why is “likelihood” function equal to product of *densities*??

a) for maximizing likelihood, we really only care about *relative* likelihoods, and density captures that

and/or


b) if density at x is $f(x)$, for any small $\delta > 0$, the probability of a sample within $\pm\delta/2$ of x is $\approx \delta f(x)$, but δ is *constant* wrt θ , so it just drops out of $d/d\theta \log L(\dots) = 0$.

Ex3: I got data; a little birdie tells me it's normal (but does *not* tell me σ^2)




Which is more likely: (a) this?

μ, σ^2 both unknown




Which is more likely: (b) or this?

μ, σ^2 both unknown




Which is more likely: (c) or this?

μ, σ^2 both unknown



Which is more likely: (d) or *this*?


μ, σ^2 both unknown



Which is more likely: (d) or *this*?

μ, σ^2 both unknown


Looks good by eye, but how do I optimize my estimates of μ & $\underline{\underline{\sigma^2}}$?



Ex 3: $x_i \sim N(\mu, \sigma^2)$, μ, σ^2 both unknown

$$\ln L(x_1, x_2, \dots, x_n | \theta_1, \theta_2) = \sum_{1 \leq i \leq n} -\frac{1}{2} \ln 2\pi\theta_2 - \frac{(x_i - \theta_1)^2}{2\theta_2}$$

$$\frac{\partial}{\partial \theta_1} \ln L(x_1, x_2, \dots, x_n | \theta_1, \theta_2) = \sum_{1 \leq i \leq n} \frac{(x_i - \theta_1)}{\theta_2} = 0$$



$$\hat{\theta}_1 = \left(\sum_{1 \leq i \leq n} x_i \right) / n = \bar{x}$$

Sample mean is MLE of population mean, again

In general, a problem like this results in 2 equations in 2 unknowns. Easy in this case, since θ_2 drops out of the $\partial/\partial\theta_1 = 0$ equation 23

Ex. 3, (cont.)

$$\ln L(x_1, x_2, \dots, x_n | \theta_1, \theta_2) = \sum_{1 \leq i \leq n} -\frac{1}{2} \ln 2\pi\theta_2 - \frac{(x_i - \theta_1)^2}{2\theta_2}$$

$$\frac{\partial}{\partial \theta_2} \ln L(x_1, x_2, \dots, x_n | \theta_1, \theta_2) = \sum_{1 \leq i \leq n} -\frac{1}{2} \frac{2\pi}{2\pi\theta_2} + \frac{(x_i - \theta_1)^2}{2\theta_2^2} = 0$$

$$\hat{\theta}_2 = \left(\sum_{1 \leq i \leq n} (x_i - \hat{\theta}_1)^2 \right) / n = \bar{s}^2$$

*Sample variance is MLE of
population variance*

Summary

MLE is *one* way to estimate *parameters* from *data*

You choose the *form* of the model (normal, binomial, ...)

Math chooses the *value(s)* of parameter(s)

Has the intuitively appealing property that the parameters maximize the *likelihood* of the observed data; basically just assumes your sample is “representative”

Of course, unusual samples will give bad estimates (estimate normal human heights from a sample of NBA stars?) but that is an unlikely event

Often, but not always, MLE has other desirable properties like being *unbiased*, or at least *consistent*