

CSE 312: Foundations of Computing II

Section 5: Variance, Independence of RVs; Zoo of discrete R.V.s

1. Review of Main Concepts - Discrete R.V.s

- (a) **Variance:** Let X be a random variable and $\mu = \mathbb{E}[X]$. The variance of X is defined to be $\text{Var}(X) = \mathbb{E}[(X - \mu)^2]$. Notice that since this is an expectation of a nonnegative random variable, i.e., $(X - \mu)^2$, variance is always nonnegative. With some algebra, we can simplify this to $\text{Var}(X) = \mathbb{E}[X^2] - \mathbb{E}[X]^2$.
- (b) **Standard Deviation:** Let X be a random variable. We define the standard deviation of X to be the square root of the variance, and denote it $\sigma = \sqrt{\text{Var}(X)}$.
- (c) **Property of Variance:** Let $a, b \in \mathbb{R}$ and let X be a random variable. Then, $\text{Var}(aX + b) = a^2\text{Var}(X)$.
- (d) **Independence:** Random variable X and event E are independent iff

$$\forall x, \quad \mathbb{P}(X = x \cap E) = \mathbb{P}(X = x)\mathbb{P}(E)$$

- (e) **Independence:** Random variables X and Y are independent iff

$$\forall x \forall y, \quad \mathbb{P}(X = x \cap Y = y) = \mathbb{P}(X = x)\mathbb{P}(Y = y)$$

In this case, we have $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$ (the converse is not necessarily true).

- (f) **Independence of functions of a r.v.:** If X and Y are independent and $g(\cdot), h(\cdot)$ are functions mapping real numbers to real numbers, then $g(X)$ and $h(Y)$ are independent. (See if you can prove this!)
- (g) **i.i.d. (independent and identically distributed):** Random variables X_1, \dots, X_n are i.i.d. (or iid) iff they are independent and have the same probability mass function.
- (h) **Variance of Independent Variables:** If X is independent of Y , $\text{Var}(X + Y) = \text{Var}(X) + \text{Var}(Y)$. This depends on independence, whereas linearity of expectation always holds. Note that this combined with the above shows that $\forall a, b, c \in \mathbb{R}$ and if X is independent of Y , $\text{Var}(aX + bY + c) = a^2\text{Var}(X) + b^2\text{Var}(Y)$.

2. Review: Zoo of Discrete Random Variables

- (a) **Uniform:** $X \sim \text{Uniform}(a, b)$ ($\text{Unif}(a, b)$ for short), for integers $a \leq b$, iff X has the following probability mass function:

$$p_X(k) = \frac{1}{b - a + 1}, \quad k = a, a + 1, \dots, b$$

$\mathbb{E}[X] = \frac{a+b}{2}$ and $\text{Var}(X) = \frac{(b-a)(b-a+1)}{12}$. This represents each integer from $[a, b]$ to be equally likely. For example, a single roll of a fair die is $\text{Uniform}(1, 6)$.

- (b) **Bernoulli (or indicator):** $X \sim \text{Bernoulli}(p)$ ($\text{Ber}(p)$ for short) iff X has the following probability mass function:

$$p_X(k) = \begin{cases} p, & k = 1 \\ 1 - p, & k = 0 \end{cases}$$

$\mathbb{E}[X] = p$ and $\text{Var}(X) = p(1 - p)$. An example of a Bernoulli r.v. is one flip of a coin with $\mathbb{P}(\text{head}) = p$.

- (c) **Binomial:** $X \sim \text{Binomial}(n, p)$ ($\text{Bin}(n, p)$ for short) iff X is the sum of n iid Bernoulli(p) random variables. X has probability mass function

$$p_X(k) = \binom{n}{k} p^k (1 - p)^{n-k}, \quad k = 0, 1, \dots, n$$

$\mathbb{E}[X] = np$ and $\text{Var}(X) = np(1 - p)$. An example of a Binomial r.v. is the number of heads in n independent flips of a coin with $\mathbb{P}(\text{head}) = p$. Note that $\text{Bin}(1, p) \equiv \text{Ber}(p)$. As $n \rightarrow \infty$ and $p \rightarrow 0$, with $np = \lambda$, then $\text{Bin}(n, p) \rightarrow \text{Poi}(\lambda)$. If X_1, \dots, X_n are independent Binomial r.v.'s, where $X_i \sim \text{Bin}(N_i, p)$, then $X = X_1 + \dots + X_n \sim \text{Bin}(N_1 + \dots + N_n, p)$.

(d) **Geometric:** $X \sim \text{Geometric}(p)$ (Geo(p) for short) iff X has the following probability mass function:

$$p_X(k) = (1-p)^{k-1} p, \quad k = 1, 2, \dots$$

$\mathbb{E}[X] = \frac{1}{p}$ and $\text{Var}(X) = \frac{1-p}{p^2}$. An example of a Geometric r.v. is the number of independent coin flips up to and including the first head, where $\mathbb{P}(\text{head}) = p$.

(e) **Poisson:** $X \sim \text{Poisson}(\lambda)$ (Poi(λ) for short) iff X has the following probability mass function:

$$p_X(k) = e^{-\lambda} \frac{\lambda^k}{k!}, \quad k = 0, 1, \dots$$

$\mathbb{E}[X] = \lambda$ and $\text{Var}(X) = \lambda$. An example of a Poisson r.v. is the number of people born during a particular minute, where λ is the average birth rate per minute. If X_1, \dots, X_n are independent Poisson r.v.'s, where $X_i \sim \text{Poi}(\lambda_i)$, then $X = X_1 + \dots + X_n \sim \text{Poi}(\lambda_1 + \dots + \lambda_n)$.

(f) **Negative Binomial :** $X \sim \text{NegativeBinomial}(r, p)$ (NegBin(r, p) for short) iff X is the sum of r iid Geometric(p) random variables. X has probability mass function

$$p_X(k) = \binom{k-1}{r-1} p^r (1-p)^{k-r}, \quad k = r, r+1, \dots$$

$\mathbb{E}[X] = \frac{r}{p}$ and $\text{Var}(X) = \frac{r(1-p)}{p^2}$. An example of a Negative Binomial r.v. is the number of independent coin flips up to and including the r^{th} head, where $\mathbb{P}(\text{head}) = p$. If X_1, \dots, X_n are independent Negative Binomial r.v.'s, where $X_i \sim \text{NegBin}(r_i, p)$, then $X = X_1 + \dots + X_n \sim \text{NegBin}(r_1 + \dots + r_n, p)$.

(g) **Hypergeometric :** $X \sim \text{HyperGeometric}(N, K, n)$ (HypGeo(N, K, n) for short) iff X has the following probability mass function:

$$p_X(k) = \frac{\binom{K}{k} \binom{N-K}{n-k}}{\binom{N}{n}}, \quad k = \max\{0, n+K-N\}, \dots, \min\{K, n\}$$

$\mathbb{E}[X] = n \frac{K}{N}$. This represents the number of successes drawn, when n items are drawn from a bag with N items (K of which are successes, and $N-K$ failures) without replacement. If we did this with replacement, then this scenario would be represented as $\text{Bin}(n, \frac{K}{N})$.

3. Review of Main Concepts - Continuous R.V.s

(a) **Cumulative Distribution Function (cdf):** For any random variable (discrete or continuous) X , the cumulative distribution function is defined as $F_X(x) = \mathbb{P}(X \leq x)$. Notice that this function must be monotonically nondecreasing: if $x < y$ then $F_X(x) \leq F_X(y)$, because $\mathbb{P}(X \leq x) \leq \mathbb{P}(X \leq y)$. Also notice that since probabilities are between 0 and 1, that $0 \leq F_X(x) \leq 1$ for all x , with $\lim_{x \rightarrow -\infty} F_X(x) = 0$ and $\lim_{x \rightarrow +\infty} F_X(x) = 1$.

(b) **Continuous Random Variable:** A continuous random variable X is one for which its cumulative distribution function $F_X(x) : \rightarrow$ is continuous everywhere. A continuous random variable has an uncountably infinite number of values.

(c) **Probability Density Function (pdf or density):** Let X be a continuous random variable. Then the probability density function $f_X(x) : \rightarrow$ of X is defined as $f_X(x) = \frac{d}{dx} F_X(x)$. Turning this around, it means that $F_X(x) = \mathbb{P}(X \leq x) = \int_{-\infty}^x f_X(t) dt$. From this, it follows that $\mathbb{P}(a \leq X \leq b) = F_X(b) - F_X(a) = \int_a^b f_X(x) dx$ and that $\int_{-\infty}^{\infty} f_X(x) dx = 1$. From the fact that $F_X(x)$ is monotonically nondecreasing it follows that $f_X(x) \geq 0$ for every real number x .

If X is a continuous random variable, note that in general $f_X(a) \neq \mathbb{P}(X = a)$, since $\mathbb{P}(X = a) = F_X(a) - F_X(a) = 0$ for all a . However, the probability that X is close to a is proportional to $f_X(a)$: for small δ , $\mathbb{P}(a - \frac{\delta}{2} < X < a + \frac{\delta}{2}) \approx \delta f_X(a)$.

- (d) **i.i.d. (independent and identically distributed)**: Random variables X_1, \dots, X_n are i.i.d. (or iid) if they are independent and have the same probability mass function or probability density function.
- (e) **Discrete to Continuous**:

	Discrete	Continuous
PMF/PDF	$p_X(x) = \mathbb{P}(X = x)$	$f_X(x) \neq \mathbb{P}(X = x) = 0$
CDF	$F_X(x) = \sum_{t < x} p_X(t)$	$F_X(x) = \int_{-\infty}^x f_X(t) dt$
Normalization	$\sum_x p_X(x) = 1$	$\int_{-\infty}^{\infty} f_X(x) dx = 1$
Expectation	$\mathbb{E}[X] = \sum_x x p_X(x)$	$\mathbb{E}[X] = \int_{-\infty}^{\infty} x f_X(x) dx$
LOTUS	$\mathbb{E}[g(X)] = \sum_x g(x) p_X(x)$	$\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(x) f_X(x) dx$

- (f) **Standardizing**: Let X be any random variable (discrete or continuous, not necessarily normal), with $\mathbb{E}[X] = \mu$ and $Var(X) = \sigma^2$. If we let $Y = \frac{X - \mu}{\sigma}$, then $\mathbb{E}[Y] = 0$ and $Var(Y) = 1$.

4. Pond Fishing

Suppose I am fishing in a pond with B blue fish, R red fish, and G green fish, where $B + R + G = N$. For each of the following scenarios, identify the most appropriate distribution (with parameter(s)):

- how many of the next 10 fish I catch are blue, if I catch and release
- how many fish I had to catch until my first green fish, if I catch and release
- how many red fish I catch in the next five minutes, if I catch on average r red fish per minute
- whether or not my next fish is blue
- how many of the next 10 fish I catch are blue, if I do not release the fish back to the pond after each catch
- how many fish I have to catch until I catch three red fish, if I catch and release

5. Best Coach Ever!!

You are a hardworking boxer. Your coach tells you that the probability of your winning a boxing match is 0.2 independently of every other match.

- How many matches do you expect to fight until you win 10 times and what kind of random variable is this?
- You only get to play 12 matches every year. To win a spot in the Annual Boxing Championship, a boxer needs to win at least 10 matches in a year. What is the probability that you will go to the Championship this year and what kind of random variable is the number of matches you win out of the 12?
- Let p be your answer to part (b). How many times can you expect to go to the Championship in your 20 year career?

6. Variance of a Product

Let X, Y, Z be independent random variables with means μ_X, μ_Y, μ_Z and variances $\sigma_X^2, \sigma_Y^2, \sigma_Z^2$, respectively. Find $Var(XY - Z)$.

7. True or False?

Identify the following statements as true or false (true means always true). Justify your answer.

- For any random variable X , we have $\mathbb{E}[X^2] \geq \mathbb{E}[X]^2$.

- (b) Let X, Y be random variables. Then, X and Y are independent if and only if $\mathbb{E}[XY] = \mathbb{E}[X] \mathbb{E}[Y]$.
- (c) Let $X \sim \text{Binomial}(n, p)$ and $Y \sim \text{Binomial}(m, p)$ be independent. Then, $X + Y \sim \text{Binomial}(n + m, p)$.
- (d) Let X_1, \dots, X_{n+1} be independent Bernoulli(p) random variables. Then, $\mathbb{E}[\sum_{i=1}^n X_i X_{i+1}] = np^2$.
- (e) Let X_1, \dots, X_{n+1} be independent Bernoulli(p) random variables. Then, $Y = \sum_{i=1}^n X_i X_{i+1} \sim \text{Binomial}(n, p^2)$.
- (f) If $X \sim \text{Bernoulli}(p)$, then $nX \sim \text{Binomial}(n, p)$.
- (g) If $X \sim \text{Binomial}(n, p)$, then $\frac{X}{n} \sim \text{Bernoulli}(p)$.
- (h) For any two independent random variables X, Y , we have $\text{Var}(X - Y) = \text{Var}(X) + \text{Var}(Y)$.

8. Fun with Poissons

Let $X \sim \text{Poisson}(\lambda_1)$ and $Y \sim \text{Poisson}(\lambda_2)$, and X and Y are independent.

- (a) Show that $X + Y \sim \text{Poisson}(\lambda_1 + \lambda_2)$
- (b) Show that $P(X = k \mid X + Y = n) = P(W = k)$ where $W \sim \text{Bin}(n, \frac{\lambda_1}{\lambda_1 + \lambda_2})$

9. Memorylessness

We say that a random variable X is memoryless if $\mathbb{P}(X > k + i \mid X > k) = \mathbb{P}(X > i)$ for all non-negative integers k and i . The idea is that X does not *remember* its history. Let $X \sim \text{Geo}(p)$. Show that X is memoryless.

10. New PDF?

Aleks came up with a function that he thinks could represent a probability density function. He defined the potential pdf for X as $f(x) = \frac{1}{1+x^2}$ defined on $[0, \infty)$. Is this a valid pdf? If not, find a constant c such that the pdf $f_X(x) = \frac{c}{1+x^2}$ is valid. Then find $\mathbb{E}[X]$. (Hints: $\frac{d}{dx}(\tan^{-1} x) = \frac{1}{1+x^2}$, $\tan \frac{\pi}{2} = \infty$, and $\tan 0 = 0$.)

11. Throwing a dart

Consider the closed unit circle of radius r , i.e., $S = \{(x, y) : x^2 + y^2 \leq r^2\}$. Suppose we throw a dart onto this circle and are guaranteed to hit it, but the dart is equally likely to land anywhere in S . Concretely this means that the probability that the dart lands in any particular area of size A (that is entirely inside the circle of radius R), is equal to $\frac{A}{\text{Area of whole circle}}$. The density outside the circle of radius r is 0.

Let X be the distance the dart lands from the center. What is the CDF and pdf of X ? What is $\mathbb{E}[X]$ and $\text{Var}(X)$?

12. A square dartboard?

You throw a dart at an $s \times s$ square dartboard. The goal of this game is to get the dart to land as close to the lower left corner of the dartboard as possible. However, your aim is such that the dart is equally likely to land at any point on the dartboard. Let random variable X be the length of the side of the smallest *square* B in the lower left corner of the dartboard that contains the point where the dart lands. That is, the lower left corner of B must be the same point as the lower left corner of the dartboard, and the dart lands somewhere along the upper or right edge of B . For X , find the CDF, PDF, $\mathbb{E}[X]$, and $\text{Var}(X)$.