

CSE 312

Foundations of Computing II




Lecture 8: Introduction to Random Variables



Aleks Jovcic

Slide Credit: Based on Stefano Tessaro's slides for 312 19au incorporating ideas from Anna Karlin, Alex Tsun, Rachel Lin, Hunter Schafer & Anna 😊

Agenda

- Random Variables 
- Probability Mass Function (PMF) 
- Cumulative Distribution Function (CDF) 
- Expectation

Random Variables (Idea)

Often: We want to **capture quantitative properties** of the outcome of a random experiment, e.g.:


- *What is the total of two dice rolls?*
- *What is the number of coin tosses needed to see the first head?*
- *What is the number of heads among 2 coin tosses?*

Random Variables

Definition. A **random variable (RV)** for a probability space (Ω, \mathbb{P}) is a function $X: \Omega \rightarrow \mathbb{R}$.

The set of values that X can take on is called its range/support Ω_X

Example. Number of heads in 2 independent coin flips $\Omega = \{HH, HT, TH, TT\}$



RV Example

$$\binom{20}{3} = |\Omega|$$

20 balls labeled 1, 2, ..., 20 in a bin

- Draw a subset of 3 uniformly at random
- Let $X =$ maximum of the 3 numbers on the balls
 - Example: $X(2, 7, 5) = 7$
 - Example: $X(15, 3, 8) = 15$
- What is $|\Omega_X|$?

$$|\Omega_X| = 18$$

A. 20^3

B. 20

C. 18

~~D. $\binom{20}{3}$~~

Agenda

- Random Variables
- Probability Mass Function (PMF) ◀
- Cumulative Distribution Function (CDF)
- Expectation


Probability Mass Function (PMF)

$\{H, \bar{H}\}$

$\frac{|E|}{|\Omega|}$

Random variables partition the sample space.

$$\sum_{x \in X(\Omega)} \mathbb{P}(X = x) = 1$$



Probability Mass Function (PMF)


Definition. For a RV $X: \Omega \rightarrow \mathbb{R}$, we define the event

$$\{X = x\} \stackrel{\text{def}}{=} \{\omega \in \Omega \mid X(\omega) = x\}$$

We write $\mathbb{P}(X = x) = \mathbb{P}(\{X = x\}) = \mathbb{P}(\{\omega \in \Omega \mid X(\omega) = x\})$ where $\mathbb{P}(X = x)$ is the **probability mass function (PMF)** of X

Random variables
partition the
sample space.

$$\sum_{x \in X(\Omega)} \mathbb{P}(X = x) = 1$$



Probability Mass Function (PMF)

Definition. For a RV $X: \Omega \rightarrow \mathbb{R}$, we define the event

$$\{X = x\} \stackrel{\text{def}}{=} \{\omega \in \Omega \mid X(\omega) = x\}$$

We write $\mathbb{P}(X = x) = \mathbb{P}(\{X = x\}) = \mathbb{P}(\{\omega \in \Omega \mid X(\omega) = x\})$ where $\mathbb{P}(X = x)$ is the **probability mass function** (PMF) of X

Random variables
partition the
sample space.

$$\sum_{x \in X(\Omega)} \mathbb{P}(X = x) = 1$$

**You also see this
notation (e.g. in
book):**

$$\mathbb{P}(X = x) = p_X(x)$$

Probability Mass Function

$$\frac{|\mathcal{E}|}{|\Omega|} = \frac{1}{4}$$

Flipping two independent coins

fair

$$\Omega = \{\text{HH}, \text{HT}, \text{TH}, \text{TT}\}$$

X = number of heads in the two flips

$$X(\text{HH}) = \underline{2}$$

$$X(\text{HT}) = \underline{1}$$

$$X(\text{TH}) = \underline{1}$$

$$X(\text{TT}) = \underline{0}$$

$\mathbb{P}(X=0)$

$$\Omega_X = \{0, 1, 2\}$$

What is $\Pr(X = k)$?

$$\mathbb{P}(X = k) = \begin{cases} 1/4 & k=0 \\ 1/2 & k=1 \\ 1/4 & k=2 \end{cases}$$

RV Example

Ω

20 balls labeled 1, 2, ..., 20 in a bin

- Draw a subset of 3 uniformly at random
- Let $X =$ maximum of the 3 numbers on the balls

What is $\Pr(X = 20)$?

3...20

$$\frac{|X = 20|}{|\Omega|} = \frac{\binom{19}{2}}{\binom{20}{3}}$$

- A. $\frac{\binom{20}{2}}{\binom{20}{3}}$ ←
- B. $\frac{\binom{19}{2}}{\binom{20}{3}}$
- C. $\frac{19^2}{\binom{20}{3}}$ ←
- D. $\frac{19 \cdot 18}{\binom{20}{3}}$ ←

Agenda

- Random Variables
- Probability Mass Function (PMF)
- Cumulative Distribution Function (CDF) ◀
- Expectation


Cumulative Distribution Function (CDF)

Definition. For a RV $X: \Omega \rightarrow \mathbb{R}$, the **cumulative distribution function** of where X specifies for any real number x , the probability that $X \leq x$.

$$F_X(x) = \Pr(X \leq x)$$

Go back to 2 coin clips, where X is the number of heads

$$\Pr(X = x) = \begin{cases} \frac{1}{4}, & x = 0 \\ \frac{1}{2}, & x = 1 \\ \frac{1}{4}, & x = 2 \end{cases} \quad F_X(x) = \left. \vphantom{\Pr(X = x)} \right\}$$




Cumulative Distribution Function (CDF)

Definition. For a RV $X: \Omega \rightarrow \mathbb{R}$, the **cumulative distribution function** of where X specifies for any real number x , the probability that $X \leq x$.

$$F_X(x) = \Pr(X \leq x)$$

Go back to 2 coin clips, where X is the number of heads

$$\Pr(X = x) = \begin{cases} \frac{1}{4}, & x = 0 \\ \frac{1}{2}, & x = 1 \\ \frac{1}{4}, & x = 2 \end{cases} \quad F_X(x) = \begin{cases} 0, & x < 0 \\ \frac{1}{4}, & 0 \leq x < 1 \\ \frac{3}{4}, & 1 \leq x < 2 \\ 1, & 2 \leq x \end{cases}$$



Example: Returning Homeworks

- Class with 3 students, randomly hand back homeworks. All permutations equally likely.
- Let X be the number of students who get their own HW

$\Pr(\omega)$	ω	$X(\omega)$
1/6	1, 2, 3	3
1/6	1, 3, 2	1
1/6	2, 1, 3	1
1/6	2, 3, 1	0
1/6	3, 1, 2	0
1/6	3, 2, 1	1

Agenda

- Random Variables
- Probability Mass Function (PMF)
- Cumulative Distribution Function (CDF)
- Expectation ◀

Expectation (Idea)

What is the *expected* number of heads in 2 independent flips of a fair coin?

Cumulative Distribution Function (CDF)

Definition. Given a discrete RV $X: \Omega \rightarrow \mathbb{R}$, the **expectation or expected value** of X is

$$E[X] = \sum_{\omega \in \Omega} X(\omega) \cdot \Pr(\omega)$$

or equivalently

$$E[X] = \sum_{x \in X(\Omega)} x \cdot \Pr(X = x)$$

Intuition: “Weighted average” of the possible outcomes (weighted by probability)

Example: Returning Homeworks

- Class with 3 students, randomly hand back homeworks. All permutations equally likely.
- Let X be the number of students who get their own HW

$\Pr(\omega)$	ω	$X(\omega)$
1/6	1, 2, 3	3
1/6	1, 3, 2	1
1/6	2, 1, 3	1
1/6	2, 3, 1	0
1/6	3, 1, 2	0
1/6	3, 2, 1	1

Flip a Biased Coin Until Heads (Independent Flips)

Suppose a coin has probability p of being heads. Keep flipping independent flips until heads. Let X be the number of flips until heads.

What is: $\Pr(X = 1) =$

What is: $\Pr(X = 2) =$

What is: $\Pr(X = k) =$

Flip a Biased Coin Until Heads (Independent Flips)

Suppose a coin has probability p of being heads. Keep flipping independent flips until heads. Let X be the number of flips until heads. What is $E[X]$?

Students on a bus

A group of 120 students are driven on 3 buses to a football game. There are 36 students in the first bus, 40 in the second and 44 in the third. Let Y be the number of students on a uniformly random bus. What is the pmf of Y and $E(Y)$? When the buses arrive, one of the 120 students is randomly chosen. Let X denote the number of students on the bus of the randomly chosen student. What is the pmf of X and what is $E(X)$?

Coin flipping again

Suppose we flip a coin with probability p of coming up Heads n times. Let X be the number of Heads in the n coin flips. What is the pmf of X ?