

Section 3

Review

1) **Conditional Probability.** $\mathbb{P}(\mathcal{B} \mid \mathcal{A}) = \underline{\hspace{2cm}}$

2) **Independent Events.** Two events \mathcal{A}, \mathcal{B} are **independent** if $\mathbb{P}(\mathcal{A} \cap \mathcal{B}) = \underline{\hspace{2cm}}$.

If $\mathbb{P}(\mathcal{A}) \neq 0$, this is equivalent to $\mathbb{P}(\mathcal{B} \mid \mathcal{A}) = \underline{\hspace{2cm}}$.

If $\mathbb{P}(\mathcal{B}) \neq 0$, this is equivalent to $\mathbb{P}(\mathcal{A} \mid \mathcal{B}) = \underline{\hspace{2cm}}$.

3) **Partition.** Nonempty events $\mathcal{E}_1, \dots, \mathcal{E}_n$ partition the sample space Ω iff

(1) $\underline{\hspace{2cm}}$, (2) $\underline{\hspace{2cm}}$

(1) , (2)

4) **Bayes Rule.** For any events \mathcal{A} and \mathcal{B} , $\mathbb{P}(\mathcal{A} \mid \mathcal{B}) = \underline{\hspace{2cm}}$.

5) **Chain Rule:** Suppose $\mathcal{A}_1, \dots, \mathcal{A}_n$ are events. Then,

$$\mathbb{P}(\mathcal{A}_1 \cap \dots \cap \mathcal{A}_n) = \underline{\hspace{2cm}}$$

6) **Law of Total Probability (LTP):** Suppose $\mathcal{E}_1, \dots, \mathcal{E}_n$ is a partition of Ω and let \mathcal{B} be any event. Then

$$\mathbb{P}(\mathcal{B}) = \sum_{i=1}^n \mathbb{P}(\mathcal{B} \cap \mathcal{E}_i) = \underline{\hspace{2cm}}$$

7) **Bayes Theorem with LTP:** $\mathcal{E}_1, \dots, \mathcal{E}_n$ is a partition of Ω and let \mathcal{B} be any event. Then

$$\mathbb{P}(\mathcal{E}_1 \mid \mathcal{B}) = \frac{\underline{\hspace{2cm}}}{\sum_{i=1}^n \underline{\hspace{2cm}}}.$$

Task 1 – Naive Bayes

Most of Section 3 will be an introduction to an application of Bayes' Theorem called the Naive Bayes Classifier.

Task 2 – Flipping Coins

We consider two independent tosses of the same coin. The coin is “heads” one quarter of the time.

- What is the probability that the second toss is “heads” given that the first toss is “tails”?
- What is the probability that the second toss is “heads” given that at least one of the tosses is “tails”?
- In the probability space of this task, give an example of two events that are disjoint but not independent.
- In the probability space of this task, give an example of two events that are independent but not disjoint.

Task 3 – Balls from an Urn – Take 2

Say an urn contains three red balls and four blue balls. Imagine we draw three balls without replacement. (You can assume every ball is uniformly selected among those remaining in the urn.)

- What is the probability that all three balls are all of the same color?

- b) What is the probability that we get more than one red ball given the first ball is red?

Task 4 – Game Show

Corrupted by their power, the judges running the popular game show America's Next Top Mathematician have been taking bribes from many of the contestants. During each of two episodes, a given contestant is either allowed to stay on the show or is kicked off. If the contestant has been bribing the judges, they will be allowed to stay with probability 1. If the contestant has not been bribing the judges, they will be allowed to stay with probability $1/3$, independent of what happens in earlier episodes. Suppose that $1/4$ of the contestants have been bribing the judges. The same contestants bribe the judges in both rounds.

- a) If you pick a random contestant, what is the probability that they are allowed to stay during the first episode?
- b) If you pick a random contestant, what is the probability they are allowed to stay during both episodes?
- c) If you pick a random contestant who was allowed to stay during the first episode, what is the probability that they get kicked off during the second episode?
- d) If you pick a random contestant who was allowed to stay during the first episode, what is the probability that they were bribing the judges?

Task 5 – Allergy Season

In a certain population, everyone is equally susceptible to colds. Each person, in particular, catches a cold with probability 0.2.

The number of colds suffered by each person during each winter season ranges from 0 to 4, with probability 0.2 for each value (see table below). A new cold prevention drug is introduced that, for people for whom the drug is effective, changes the probabilities as shown in the table. Unfortunately, the effects of the drug last only the duration of one winter season, and the drug is only effective in 20% of people, independently.

number of colds	no drug or ineffective	drug effective
0	0.2	0.4
1	0.2	0.3
2	0.2	0.2
3	0.2	0.1
4	0.2	0.0

- a) Sneezzy decides to take the drug. Given that he gets 1 cold that winter, what is the probability that the drug is effective for Sneezzy?
- b) The next year he takes the drug again. Given that he gets 2 colds in this winter, what is the updated probability that the drug is effective for Sneezzy?
- c) Why is the answer to (b) the same as the answer to (a)?

Task 6 – Coins

There are three coins, C_1 , C_2 , and C_3 . The probability of "heads" is 1 for C_1 , 0 for C_2 , and p for C_3 . A coin is picked among these three uniformly at random, and then flipped a certain number of times.

- a) What is the probability that the first n flips are tails?
- b) Given that the first n flips were tails, what is the probability that C_1 was flipped / C_2 was flipped / C_3 was flipped?

Task 7 – Parallel Systems

A parallel system functions whenever at least one of its components works. Consider a parallel system of n components and suppose that each component works with probability p independently.

- a) What is the probability the system is functioning?
- b) If the system is functioning, what is the probability that component 1 is working?
- c) If the system is functioning and component 2 is working, what is the probability that component 1 is working?

Task 8 – Marbles in Pockets

A girl has 5 blue and 3 white marbles in her left pocket, and 4 blue and 4 white marbles in her right pocket. If she transfers a randomly chosen marble from left pocket to right pocket without looking, and then draws a randomly chosen marble from her right pocket, what is the probability that it is blue?

Task 9 – A game

Pemi and Shreya are playing the following game: A 6-sided die is thrown and each time it's thrown, regardless of the history, it is equally likely to show any of the six numbers.

- If it shows 5, Pemi wins.
- If it shows 1, 2, or 6, Shreya wins.
- Otherwise, they play a second round and so on.

What is the probability that Shreya wins on the 4th round?

Task 10 – Another game

Leiyi and Luxi are playing a tournament in which they stop as soon as one of them wins n games. Luxi wins each game with probability p and Leiyi wins with probability $1 - p$, independently of other games. What is the probability that Luxi wins and that when the match is over, Leiyi has won k games?

Task 11 – Random Variables

(The material for this problem will most likely be covered early next week.)

Assume that we roll a fair 3-sided die three times. Here, the sides have values 1, 2, 3.

- a) Describe the PMF of the random variable X giving the sum of the first two rolls.
- b) Give the expectation $\mathbb{E}[X]$.
- c) Compute $\mathbb{P}(X > 3)$.
- d) Let Y be the random variable describing the sum of the three rolls. Compute $\mathbb{P}(X = 5 \mid Y = 7)$.