

# CSE 312 Section 2

**Intro to Discrete Probability**

# Administrivia



# Announcements & Reminders

- Office Hours
  - Offered every day! Mostly in-person, a few zoom options
  - Times posted on the calendar on the course website
- HW1
  - Was due yesterday, Wednesday 4/3 @ 11:59pm
  - Late deadline Saturday 4/6 @ 11:59pm (max of 3 late days per homework)
- HW2
  - Released on the course website
  - Due Wednesday 4/10 @ 11:59pm

# Homework

- Submissions
  - LaTeX (highly encouraged)
    - overleaf.com
    - template and LaTeX guide posted on course website!
  - Word Editor that supports mathematical equations
  - Handwritten neatly and scanned
- Homework will typically be due on Wednesdays at 11:59pm on Gradescope
- Each assignment can be submitted a max of 72 hours late
- You have **8 late days total** to use throughout the quarter
  - Anything beyond that will result in a deduction on further late assignments

# Review



**Any lingering questions from this last week?**

# Kahoot!

These questions are also Task 1 on the section handout!

# Problem 6 – Powers and Divisibility





## 6 – Powers and Divisibility

Prove that there exist two powers of 7 whose difference is divisible by 2003. (You may want to use the Pigeonhole principle.)

Work on this problem with the people around you, and then we'll go over it together!

## 6 – Powers and Divisibility

Prove that there exist two powers of 7 whose difference is divisible by 2003. (You may want to use the Pigeonhole principle.)

HINT: think about what are the pigeons, and what are the pigeonholes?

Work on this problem with the people around you, and then we'll go over it together!

## 6 – Powers and Divisibility

Prove that there exist two powers of 7 whose difference is divisible by 2003. (You may want to use the Pigeonhole principle.)

## 6 – Powers and Divisibility

Prove that there exist two powers of 7 whose difference is divisible by 2003. (You may want to use the Pigeonhole principle.)

There are 2003 possible remainders of division by 2003 (0 through 2002).

## 6 – Powers and Divisibility

Prove that there exist two powers of 7 whose difference is divisible by 2003. (You may want to use the Pigeonhole principle.)

There are 2003 possible remainders of division by 2003 (0 through 2002).

Consider a sequence of powers  $1, 7, 7^2, \dots, 7^{2003}$ , it contains 2004 members.

Each of which can be expressed as  $7^i = 2003x_i + r_i$ .

## 6 – Powers and Divisibility

Prove that there exist two powers of 7 whose difference is divisible by 2003. (You may want to use the Pigeonhole principle.)

There are 2003 possible remainders of division by 2003 (0 through 2002).

Consider a sequence of powers  $1, 7, 7^2, \dots, 7^{2003}$ , it contains 2004 members.

Each of which can be expressed as  $7^i = 2003x_i + r_i$ .

There are 2004 such  $r_i$ s (pigeons) which need to be mapped to 2003 possible remainders (pigeonholes)

# 6 – Powers and Divisibility

Prove that there exist two powers of 7 whose difference is divisible by 2003. (You may want to use the Pigeonhole principle.)

There are 2003 possible remainders of division by 2003 (0 through 2002).

Consider a sequence of powers  $1, 7, 7^2, \dots, 7^{2003}$ , it contains 2004 members.

Each of which can be expressed as  $7^i = 2003x_i + r_i$ .

There are 2004 such  $r_i$ s (pigeons) which need to be mapped to 2003 possible remainders (pigeonholes)

Therefore, by Pigeonhole principle, there exists a pair of members, say  $7^n$  and  $7^m$ , where  $n > m$ , such that  $r_n = r_m$ .

# 6 – Powers and Divisibility

Prove that there exist two powers of 7 whose difference is divisible by 2003. (You may want to use the Pigeonhole principle.)

There are 2003 possible remainders of division by 2003 (0 through 2002). Consider a sequence of powers  $1, 7, 7^2, \dots, 7^{2003}$ , it contains 2004 members. Each of which can be expressed as  $7^i = 2003x_i + r_i$ .

There are 2004 such  $r_i$ s (pigeons) which need to be mapped to 2003 possible remainders (pigeonholes)

Therefore, by Pigeonhole principle, there exists a pair of members, say  $7^n$  and  $7^m$ , where  $n > m$ , such that  $r_n = r_m$ .

Then their difference  $7^n - 7^m = 2003x_n + r_n - 2003x_m - r_m = 2003(x_n - x_m)$  is divisible by 2003.



# Problem 10 – Balls from an Urn



# 10 – Balls from an Urn

Say an urn contains one red ball, one blue ball, and one green ball. (Other than for their colors, balls are identical.) Imagine we draw two balls with replacement, i.e., after drawing one ball, we put it back into the urn, before we draw the second one. (In particular, each ball is equally likely to be drawn.)

- a) Give a probability space describing the experiment.
- b) What is the probability that both balls are red? (Describe the event first, before you compute its probability.)
- c) What is the probability that at most one ball is red?
- d) What is the probability that we get at least one green ball?
- e) Repeat b)-d) for the case where the balls are drawn without replacement, i.e., when the first ball is drawn, it is not placed back from the urn.

Work on this problem with the people around you, and then we'll go over it together!

# 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

a) Give a probability space describing the experiment.

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

a) Give a probability space describing the experiment.

$$\Omega = \{BB, BR, BG, RB, RR, RG, GB, GR, GG\} = \{B, R, G\}^2$$

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

a) Give a probability space describing the experiment.

$$\Omega = \{BB, BR, BG, RB, RR, RG, GB, GR, GG\} = \{B, R, G\}^2$$

$$\mathbb{P}(\omega) = 1/9 \text{ for all } \omega \in \Omega$$

# 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

- b) What is the probability that both balls are red? (Describe the event first, before you compute its probability.)

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

b) What is the probability that both balls are red? (Describe the event first, before you compute its probability.)

The event is  $B = \{RR\}$

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

b) What is the probability that both balls are red? (Describe the event first, before you compute its probability.)

The event is  $B = \{RR\}$

The probability of the event is  $\mathbb{P}(B) = \frac{|B|}{9} = 1/9$



## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

c) What is the probability that at most one ball is red?

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

c) What is the probability that at most one ball is red?

The event is  $C = \{BB, BR, BG, RB, RG, GB, GR, GG\}$

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

c) What is the probability that at most one ball is red?

The event is  $C = \{BB, BR, BG, RB, RG, GB, GR, GG\}$

The probability of the event is  $\mathbb{P}(C) = \frac{|C|}{9} = 8/9$

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

c) What is the probability that at most one ball is red?

The event is  $C = \{BB, BR, BG, RB, RG, GB, GR, GG\}$

The probability of the event is  $\mathbb{P}(C) = \frac{|C|}{9} = 8/9$

Alternatively, this is just  $B^C$ , the complement of  $B$ .

We know that  $\mathbb{P}(B^C) = 1 - \mathbb{P}(B) = 1 - 1/9 = 8/9$

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

d) What is the probability that we get at least one green ball?

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

d) What is the probability that we get at least one green ball?

The event is  $D = \{BG, GB, RG, GR, GG\}$

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball. Draw two balls with replacement.

d) What is the probability that we get at least one green ball?

The event is  $D = \{BG, GB, RG, GR, GG\}$

The probability of the event is  $\mathbb{P}(D) = \frac{|D|}{9} = 5/9$

# 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball.

e) Repeat b)-d) for the case where **the balls are drawn without replacement.**

a) Does the probability space change? If so, what is it now?

a) What is the probability that both balls are red?

a) What is the probability that at most one ball is red?

a) What is the probability that we get at least one green ball?



# 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball.

e) Repeat b)-d) for the case where **the balls are drawn without replacement**.

a) Does the probability space change? If so, what is it now?

$$\Omega = \{BR, BG, RB, RG, GB, GR\}$$

$$\mathbb{P}(\omega) = \frac{1}{3} \cdot \frac{1}{2} = 1/6 \text{ for all } \omega \in \Omega \text{ (3 choices for the first ball, 2 for the second)}$$

a) What is the probability that both balls are red?

a) What is the probability that at most one ball is red?

a) What is the probability that we get at least one green ball?

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball.

e) Repeat b)-d) for the case where **the balls are drawn without replacement**.

a) Does the probability space change? If so, what is it now?

$$\Omega = \{BR, BG, RB, RG, GB, GR\}$$

$$\mathbb{P}(\omega) = \frac{1}{3} \cdot \frac{1}{2} = 1/6 \text{ for all } \omega \in \Omega \text{ (3 choices for the first ball, 2 for the second)}$$

a) What is the probability that both balls are red?

$$B = \{\} \text{ We can't have both be red, so } \mathbb{P}(B) = 0$$

a) What is the probability that at most one ball is red?

a) What is the probability that we get at least one green ball?

## 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball.

e) Repeat b)-d) for the case where **the balls are drawn without replacement**.

a) Does the probability space change? If so, what is it now?

$$\Omega = \{BR, BG, RB, RG, GB, GR\}$$

$$\mathbb{P}(\omega) = \frac{1}{3} \cdot \frac{1}{2} = 1/6 \text{ for all } \omega \in \Omega \text{ (3 choices for the first ball, 2 for the second)}$$

a) What is the probability that both balls are red?

$$B = \{\} \text{ We can't have both be red, so } \mathbb{P}(B) = 0$$

a) What is the probability that at most one ball is red?

$$C = B^c \text{ so } \mathbb{P}(B^c) = 1 - \mathbb{P}(B) = 1$$

a) What is the probability that we get at least one green ball?

# 10 – Balls from an Urn

An urn contains one red ball, one blue ball, and one green ball.

e) Repeat b)-d) for the case where **the balls are drawn without replacement**.

a) Does the probability space change? If so, what is it now?

$$\Omega = \{BR, BG, RB, RG, GB, GR\}$$

$$\mathbb{P}(\omega) = \frac{1}{3} \cdot \frac{1}{2} = 1/6 \text{ for all } \omega \in \Omega \text{ (3 choices for the first ball, 2 for the second)}$$

a) What is the probability that both balls are red?

$$B = \{\} \text{ We can't have both be red, so } \mathbb{P}(B) = 0$$

a) What is the probability that at most one ball is red?

$$C = B^c \text{ so } \mathbb{P}(B^c) = 1 - \mathbb{P}(B) = 1$$

a) What is the probability that we get at least one green ball?

$$D = \{BG, GB, RG, GR\} \text{ so } \mathbb{P}(D) = \frac{4}{6} = \frac{2}{3}$$

# Problem 11 – Weighted Die



# 11 – Weighted Die

Consider a weighted die such that

- $\mathbb{P}(1) = \mathbb{P}(2)$
- $\mathbb{P}(3) = \mathbb{P}(4) = \mathbb{P}(5) = \mathbb{P}(6)$
- $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$

What is the probability that the outcome is 3 or 4?

Work on this problem with the people around you, and then we'll go over it together!

# 11 – Weighted Die

Consider a weighted die such that

- $\mathbb{P}(1) = \mathbb{P}(2)$
- $\mathbb{P}(3) = \mathbb{P}(4) = \mathbb{P}(5) = \mathbb{P}(6)$
- $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$

What is the probability that the outcome is 3 or 4?

# 10 – Weighted Die

Consider a weighted die such that

- $\mathbb{P}(1) = \mathbb{P}(2)$
- $\mathbb{P}(3) = \mathbb{P}(4) = \mathbb{P}(5) = \mathbb{P}(6)$
- $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$

What is the probability that the outcome is 3 or 4?

By the second axiom of probability, the sum of probabilities for the sample space must equal 1.



# 10 – Weighted Die

Consider a weighted die such that

- $\mathbb{P}(1) = \mathbb{P}(2)$
- $\mathbb{P}(3) = \mathbb{P}(4) = \mathbb{P}(5) = \mathbb{P}(6)$
- $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$

What is the probability that the outcome is 3 or 4?

By the second axiom of probability, the sum of probabilities for the sample space must equal 1.

That is,  $\sum_{i=1}^6 \mathbb{P}(i) = 1$ .

# 10 – Weighted Die

Consider a weighted die such that

- $\mathbb{P}(1) = \mathbb{P}(2)$
- $\mathbb{P}(3) = \mathbb{P}(4) = \mathbb{P}(5) = \mathbb{P}(6)$
- $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$

What is the probability that the outcome is 3 or 4?

By the second axiom of probability, the sum of probabilities for the sample space must equal 1.

That is,  $\sum_{i=1}^6 \mathbb{P}(i) = 1$ .

Since  $\mathbb{P}(1) = \mathbb{P}(2)$  and  $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$ , we have that:

# 10 – Weighted Die

Consider a weighted die such that

- $\mathbb{P}(1) = \mathbb{P}(2)$
- $\mathbb{P}(3) = \mathbb{P}(4) = \mathbb{P}(5) = \mathbb{P}(6)$
- $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$

What is the probability that the outcome is 3 or 4?

By the second axiom of probability, the sum of probabilities for the sample space must equal 1.

That is,  $\sum_{i=1}^6 \mathbb{P}(i) = 1$ .

Since  $\mathbb{P}(1) = \mathbb{P}(2)$  and  $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$ , we have that:

$$\begin{aligned} 1 &= \mathbb{P}(1) + \mathbb{P}(2) + \mathbb{P}(3) + \mathbb{P}(4) + \mathbb{P}(5) + \mathbb{P}(6) \\ &= 3 \cdot \mathbb{P}(3) + 3 \cdot \mathbb{P}(3) + \mathbb{P}(3) + \mathbb{P}(3) + \mathbb{P}(3) = 10 \cdot \mathbb{P}(3) \end{aligned}$$

# 10 – Weighted Die

Consider a weighted die such that

- $\mathbb{P}(1) = \mathbb{P}(2)$
- $\mathbb{P}(3) = \mathbb{P}(4) = \mathbb{P}(5) = \mathbb{P}(6)$
- $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$

What is the probability that the outcome is 3 or 4?

By the second axiom of probability, the sum of probabilities for the sample space must equal 1.

That is,  $\sum_{i=1}^6 \mathbb{P}(i) = 1$ .

Since  $\mathbb{P}(1) = \mathbb{P}(2)$  and  $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$ , we have that:

$$\begin{aligned} 1 &= \mathbb{P}(1) + \mathbb{P}(2) + \mathbb{P}(3) + \mathbb{P}(4) + \mathbb{P}(5) + \mathbb{P}(6) \\ &= 3 \cdot \mathbb{P}(3) + 3 \cdot \mathbb{P}(3) + \mathbb{P}(3) + \mathbb{P}(3) + \mathbb{P}(3) = 10 \cdot \mathbb{P}(3) \end{aligned}$$

Thus, solving algebraically,  $\mathbb{P}(3) = 0.1$ , so  $\mathbb{P}(3) = \mathbb{P}(4) = 0.1$ .

# 10 – Weighted Die

Consider a weighted die such that

- $\mathbb{P}(1) = \mathbb{P}(2)$
- $\mathbb{P}(3) = \mathbb{P}(4) = \mathbb{P}(5) = \mathbb{P}(6)$
- $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$

What is the probability that the outcome is 3 or 4?

By the second axiom of probability, the sum of probabilities for the sample space must equal 1.

That is,  $\sum_{i=1}^6 \mathbb{P}(i) = 1$ .


Since  $\mathbb{P}(1) = \mathbb{P}(2)$  and  $\mathbb{P}(1) = 3 \cdot \mathbb{P}(3)$ , we have that:

$$\begin{aligned} 1 &= \mathbb{P}(1) + \mathbb{P}(2) + \mathbb{P}(3) + \mathbb{P}(4) + \mathbb{P}(5) + \mathbb{P}(6) \\ &= 3 \cdot \mathbb{P}(3) + 3 \cdot \mathbb{P}(3) + \mathbb{P}(3) + \mathbb{P}(3) + \mathbb{P}(3) = 10 \cdot \mathbb{P}(3) \end{aligned}$$

Thus, solving algebraically,  $\mathbb{P}(3) = 0.1$ , so  $\mathbb{P}(3) = \mathbb{P}(4) = 0.1$ .

Since rolling a 3 and 4 are disjoint events, then  $\mathbb{P}(3 \text{ or } 4) = \mathbb{P}(3) + \mathbb{P}(4) = 0.1 + 0.1 = 0.2$

# Problem 2 - Subsubset



## 2 – Subsubset

Let  $[n] = \{1, 2, \dots, n\}$  denote the first  $n$  natural numbers. How many (ordered) pairs of subsets  $(A, B)$  are there such that  $A \subseteq B \subseteq [n]$ ?

Work on this problem with the people around you, and then we'll go over it together!

## 2 – Subsubset

Let  $[n] = \{1, 2, \dots, n\}$  denote the first  $n$  natural numbers. How many (ordered) pairs of subsets  $(A, B)$  are there such that  $A \subseteq B \subseteq [n]$ ?



## 2 – Subsubset

Let  $[n] = \{1, 2, \dots, n\}$  denote the first  $n$  natural numbers. How many (ordered) pairs of subsets  $(A, B)$  are there such that  $A \subseteq B \subseteq [n]$ ?

Realize that, if there are no restrictions, for each element  $i$  of  $1, \dots, n$ , there are four possibilities: it can be in only  $A$ , only  $B$ , both, or neither. In our case, there is only one that is not valid (violates  $A \subseteq B$ ): being in  $A$  but not  $B$ . Hence there are 3 choices for each element, so the total number of such ordered pairs of subsets is

## 2 – Subsubset

Let  $[n] = \{1, 2, \dots, n\}$  denote the first  $n$  natural numbers. How many (ordered) pairs of subsets  $(A, B)$  are there such that  $A \subseteq B \subseteq [n]$ ?

Realize that, if there are no restrictions, for each element  $i$  of  $1, \dots, n$ , there are four possibilities: it can be in only  $A$ , only  $B$ , both, or neither. In our case, there is only one that is not valid (violates  $A \subseteq B$ ): being in  $A$  but not  $B$ . Hence there are 3 choices for each element, so the total number of such ordered pairs of subsets is

$$3^n$$

## 2 – Subsubset

Let  $[n] = \{1, 2, \dots, n\}$  denote the first  $n$  natural numbers. How many (ordered) pairs of subsets  $(A, B)$  are there such that  $A \subseteq B \subseteq [n]$ ?

Alternately, apply the sum rule by adding up the number of ways of doing this where  $B$  has size  $k$ , where  $k$  is any integer between 0 and  $n$ . Now apply the product rule to find the number of ways to choose  $B$  of size exactly  $k$  (there are  $\binom{n}{k}$  possibilities for  $B$ ), and then once  $B$  is selected, count the number of ways of choosing  $A$  which has to be a subset of  $B$  ( $2^k$  ways).

Hence, by the Binomial Theorem, the number of such ordered pairs of subsets is

## 2 – Subsubset

Let  $[n] = \{1, 2, \dots, n\}$  denote the first  $n$  natural numbers. How many (ordered) pairs of subsets  $(A, B)$  are there such that  $A \subseteq B \subseteq [n]$ ?

Alternately, apply the sum rule by adding up the number of ways of doing this where  $B$  has size  $k$ , where  $k$  is any integer between 0 and  $n$ . Now apply the product rule to find the number of ways to choose  $B$  of size exactly  $k$  (there are  $\binom{n}{k}$  possibilities for  $B$ ), and then once  $B$  is selected, count the number of ways of choosing  $A$  which has to be a subset of  $B$  ( $2^k$  ways).

Hence, by the Binomial Theorem, the number of such ordered pairs of subsets is

$$\sum_{k=0}^n \binom{n}{k} 2^k = \sum_{k=0}^n \binom{n}{k} 2^k 1^{n-k} = 3^n$$

# Problem 4 – GREED INNIT



## 4 – GREED INNIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other. For example, “INGREEDINT” is invalid because the two E’s are adjacent.

Repeat the question for the letters “AAAAABBB”.

Work on this problem with the people around you, and then we’ll go over it together!

## 4 – GREED INNIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

## 4 – GREED INNIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

We use inclusion-exclusion.



## 4 – GREED INNIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

We use inclusion-exclusion.

Let  $\Omega$  be the set of all anagrams (permutations) of “INGREDIENT” and  $A_I$  be the set of all anagrams with two consecutive I’s. Define  $A_E$  and  $A_N$  similarly.

## 4 – GREED ININIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

We use inclusion-exclusion.

Let  $\Omega$  be the set of all anagrams (permutations) of “INGREDIENT” and  $A_I$  be the set of all anagrams with two consecutive I’s. Define  $A_E$  and  $A_N$  similarly.

$A_I \cup A_E \cup A_N$  clearly are the set of anagrams we don’t want.

## 4 – GREED ININIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

We use inclusion-exclusion.

Let  $\Omega$  be the set of all anagrams (permutations) of “INGREDIENT” and  $A_I$  be the set of all anagrams with two consecutive I’s. Define  $A_E$  and  $A_N$  similarly.

$A_I \cup A_E \cup A_N$  clearly are the set of anagrams we don’t want.

We use complementing to count the size of  $\Omega \setminus (A_I \cup A_E \cup A_N)$ .

## 4 – GREED ININIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

We use inclusion-exclusion.

Let  $\Omega$  be the set of all anagrams (permutations) of “INGREDIENT” and  $A_I$  be the set of all anagrams with two consecutive I’s. Define  $A_E$  and  $A_N$  similarly.

$A_I \cup A_E \cup A_N$  clearly are the set of anagrams we don’t want.

We use complementing to count the size of  $\Omega \setminus (A_I \cup A_E \cup A_N)$ .

By inclusion exclusion,  $|A_I \cup A_E \cup A_N| = \text{singles} - \text{doubles} + \text{triples}$ , and by complementing,  $|\Omega \setminus (A_I \cup A_E \cup A_N)| = |\Omega| - |A_I \cup A_E \cup A_N|$ .

## 4 – GREED ININIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

$$\text{Singles} = |A_I| + |A_E| + |A_N|$$

$$\text{Doubles} = |A_I \cap A_E| + |A_I \cap A_N| + |A_E \cap A_N|$$

$$\text{Triples} = |A_I \cap A_E \cap A_N|$$

## 4 – GREED ININIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

$$\text{Singles} = |A_I| + |A_E| + |A_N|$$

$$\text{Doubles} = |A_I \cap A_E| + |A_I \cap A_N| + |A_E \cap A_N|$$

$$\text{Triples} = |A_I \cap A_E \cap A_N|$$

First,  $|\Omega| = \frac{10!}{2!2!2!}$  because there are 2 of each of I,E,N's (multinomial coefficient).

## 4 – GREED ININIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

$$\text{Singles} = |A_I| + |A_E| + |A_N|$$

$$\text{Doubles} = |A_I \cap A_E| + |A_I \cap A_N| + |A_E \cap A_N|$$

$$\text{Triples} = |A_I \cap A_E \cap A_N|$$

First,  $|\Omega| = \frac{10!}{2!2!2!}$  because there are 2 of each of I,E,N's (multinomial coefficient).

$|A_I| = \frac{9!}{2!2!}$  because we treat the two adjacent I's as one entity.

Clearly,  $|A_I| = |A_E| = |A_N|$ , so  $\text{Singles} = 3 \cdot \frac{9!}{2!2!}$

## 4 – GREED INNTIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

$$\text{Singles} = |A_I| + |A_E| + |A_N|$$

$$\text{Doubles} = |A_I \cap A_E| + |A_I \cap A_N| + |A_E \cap A_N|$$

$$\text{Triples} = |A_I \cap A_E \cap A_N|$$

First,  $|\Omega| = \frac{10!}{2!2!2!}$  because there are 2 of each of I,E,N's (multinomial coefficient).

$|A_I| = \frac{9!}{2!2!}$  because we treat the two adjacent I's as one entity.

Clearly,  $|A_I| = |A_E| = |A_N|$ , so  $\text{Singles} = 3 \cdot \frac{9!}{2!2!}$

$|A_I \cap A_E| = \frac{8!}{2!}$  because we treat the two adjacent I's as one entity and the two adjacent E's as one entity.  $|A_I \cap A_E| = |A_I \cap A_N| = |A_E \cap A_N|$ , so  $\text{Doubles} = 3 \cdot \frac{8!}{2!}$



## 4 – GREED ININIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

$$\text{Singles} = |A_I| + |A_E| + |A_N|$$

$$\text{Doubles} = |A_I \cap A_E| + |A_I \cap A_N| + |A_E \cap A_N|$$

$$\text{Triples} = |A_I \cap A_E \cap A_N|$$

First,  $|\Omega| = \frac{10!}{2!2!2!}$  because there are 2 of each of I,E,N's (multinomial coefficient).

$|A_I| = \frac{9!}{2!2!}$  because we treat the two adjacent I's as one entity.

Clearly,  $|A_I| = |A_E| = |A_N|$ , so  $\text{Singles} = 3 \cdot \frac{9!}{2!2!}$

$|A_I \cap A_E| = \frac{8!}{2!}$  because we treat the two adjacent I's as one entity and the two adjacent E's as one entity.  $|A_I \cap A_E| = |A_I \cap A_N| = |A_E \cap A_N|$ , so  $\text{Doubles} = 3 \cdot \frac{8!}{2!}$

Finally,  $|A_I \cap A_E \cap A_N| = 7!$  since we treat each pair of adjacent I's, E's, and N's as one entity.

## 4 – GREED INNIT

Find the number of ways to rearrange the word “INGREDIENT”, such that no two identical letters are adjacent to each other.

$$\text{Singles} = |A_I| + |A_E| + |A_N|$$

$$\text{Doubles} = |A_I \cap A_E| + |A_I \cap A_N| + |A_E \cap A_N|$$

$$\text{Triples} = |A_I \cap A_E \cap A_N|$$

First,  $|\Omega| = \frac{10!}{2!2!2!}$  because there are 2 of each of I,E,N's (multinomial coefficient).

$|A_I| = \frac{9!}{2!2!}$  because we treat the two adjacent I's as one entity.

Clearly,  $|A_I| = |A_E| = |A_N|$ , so  $\text{Singles} = 3 \cdot \frac{9!}{2!2!}$

$|A_I \cap A_E| = \frac{8!}{2!}$  because we treat the two adjacent I's as one entity and the two adjacent E's as one entity.  $|A_I \cap A_E| = |A_I \cap A_N| = |A_E \cap A_N|$ , so  $\text{Doubles} = 3 \cdot \frac{8!}{2!}$

Finally,  $|A_I \cap A_E \cap A_N| = 7!$  since we treat each pair of adjacent I's, E's, and N's as one entity.

Putting this together gives  $\frac{10!}{2!2!2!} - \left( 3 \cdot \frac{9!}{2!2!} - 3 \cdot \frac{8!}{2!} + 7! \right)$

## 4 – GREED INNIT

Repeat the question for the letters “AAAAABBB”.

## 4 – GREED INNIT

Repeat the question for the letters “AAAAABBB”.

Note that no A’s and no B’s can be adjacent. So let us put the B’s down first:

\_B\_B\_B\_

By the pigeonhole principle, two A’s must go in the same slot, but then they would be adjacent, so there are no ways .

# Problem 12 – Shuffling Cards



## 12 – Shuffling Cards

We have a deck of cards, with 4 suits, with 13 cards in each. Within each suit, the cards are ordered Ace > King > Queen > Jack > 10 > . . . > 2. Also, suppose we perfectly shuffle the deck (i.e., all possible shuffles are equally likely). What is the probability the first card on the deck is (strictly) larger than the second one?

Work on this problem with the people around you, and then we'll go over it together!

## 12 – Shuffling Cards

We have a deck of cards, with 4 suits, with 13 cards in each. Within each suit, the cards are ordered Ace > King > Queen > Jack > 10 > . . . > 2. Also, suppose we perfectly shuffle the deck (i.e., all possible shuffles are equally likely). What is the probability the first card on the deck is (strictly) larger than the second one?

## 12 – Shuffling Cards

We have a deck of cards, with 4 suits, with 13 cards in each. Within each suit, the cards are ordered Ace > King > Queen > Jack > 10 > . . . > 2. Also, suppose we perfectly shuffle the deck (i.e., all possible shuffles are equally likely). What is the probability the first card on the deck is (strictly) larger than the second one?

First off, the sample space  $\Omega$  here consists of all pairs of cards – which we can represent by their value and suit, e.g.,  $(4\clubsuit, A\spadesuit)$ . There are  $52 \cdot 51 = 2652$  possible outcomes, therefore  $\mathbb{P}(\omega) = \frac{1}{2652}$  for all  $\omega \in \Omega$ .

Let us now look at the size of the event  $E$  containing all pairs where the first card is strictly larger than the second. Then, the number of pairs of values of cards  $a$  and  $b$  where  $a < b$  is exactly  $\binom{13}{2} = 13 \cdot 6 = 78$ . We can then assign suits to each of them – given the cards are different, all suits are possible for each, so there are  $4^2 = 16$  choices. Thus, overall,

$$|E| = 16 \cdot 78 = 1248$$

$$\text{Therefore, } \mathbb{P}(E) = \frac{|E|}{|\Omega|} = \frac{16 \cdot 78}{52 \cdot 51} = \frac{13 \cdot 3 \cdot 2^5}{13 \cdot 3 \cdot 2^2 \cdot 17} = \frac{8}{17} \approx 0.47$$



# Problem 5 - Friendships



## 5 – Friendships

Show that in any group  $n$  people there are two who have an identical number of friends within the group. (Friendship is bi-directional – i.e., if  $A$  is friend of  $B$ , then  $B$  is friend of  $A$  – and nobody is a friend of themselves.) Solve in particular the following two cases individually:

- a) Everyone has at least one friend.
- b) At least one person has no friends.

Work on this problem with the people around you, and then we'll go over it together!

# 5 – Friendships

- a) Everyone has at least one friend.

## 5 – Friendships

a) Everyone has at least one friend.

Everyone has between 1 and  $n - 1$  friends (i.e.,  $n - 1$  holes), and there are  $n$  people (the “pigeons”). Therefore, two of them will have the same number of friends.

## 5 – Friendships

b) At least one person has no friends.

## 5 – Friendships

b) At least one person has no friends.

Here, we need to observe that if someone has 0 friends, then nobody has  $n - 1$  friends (by the symmetry of the friendship relation). Then, possible choices are now between 0 and  $n - 2$  friends (i.e.,  $n - 1$  holes), and there are  $n$  people (the “pigeons”). Therefore, two of them will have the same number of friends.

# **That's All, Folks!**

**Thanks for coming to section this week!  
Any questions?**