


CSE 312 Section 1

Combinatorics

Administrivia & Introductions



Announcements & Reminders

- Section Materials
 - Handouts will be provided in at each section
 - Worksheets and sample solutions will be available on the course calendar later this evening
- Office Hours
 - We'll start hosting office hours Monday!
 - Exact times and locations will be posted on the calendar on the course website
- HW1
 - Released later today!
 - Due Wednesday, October 1st @ 11:59pm

Homework

- Submissions
 - LaTeX (highly encouraged)
 - Parchmynt (easier-to-learn interface for subset of LaTeX)
 - overleaf.com (full-power of LaTeX).
 - Login through UW's SSO for free Overleaf Premium!
 - template and LaTeX guide posted on course website!
 - Word Editor that supports mathematical equations
 - Handwritten neatly and scanned
- Homework will typically be due on Wednesdays at 11:59pm on Gradescope
- Each assignment can be submitted a max of 72 hours late
- You have **6 late days total** to use throughout the quarter
 - Anything beyond that will result in a 15% deduction per day

Your TAs

- TA 1
 - Intro
- TA 2
 - Intro

Our Office Hours

- TA 1
 - OH
- TA 2
 - OH

Icebreaker

- Small groups of 4-6ish
- Please share with your group
 - Your name
 - Number of years in department/ at UW
 - What was something fun you did over Spring break?
 - What are you concerned about for 312 / what are you excited about?
- Then, share how you like to eat your potatoes (baked, fried, chips, etc)
- We'll go around and see what style of potato is most popular!



Review




Any lingering questions from this last week?

Each week in section, we'll be reviewing the main concepts from this week by going through some practice problems together. But before that, we'll try to start off each section with some time for you to ask questions.

Was anything particularly confusing this week? Is there anything we can clarify before we dive into the review? This is your chance to clear things up!

Problem 1 - Seating



1 – Seating

How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if ...

- a) ... all couples are to get adjacent seats?
- b) ... anyone can sit anywhere, except that one couple insists on not sitting in adjacent seats?

Work on this problem with the people around you, and then we'll go over it together!

1 – Seating

- a) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if all couples are to get adjacent seats?

1 – Seating

- a) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if all couples are to get adjacent seats?

Consider each couple as a unit.

1 – Seating

- a) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if all couples are to get adjacent seats?

Consider each couple as a unit.

Count the number of ways we can order the 5 couples in the row.

1 – Seating

- a) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if all couples are to get adjacent seats?

Consider each couple as a unit.

Count the number of ways we can order the 5 couples in the row.

We can choose any one of the 5 couples for the first pair of seats, then any of the 4 remaining couples for the second pair of seats, then any of the 3 remaining couples for the third pair of seats, then one of the 2 remaining couples for the fourth pair of seats. The last couple must go into the last pair of seats.

1 – Seating

- a) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if all couples are to get adjacent seats?

Consider each couple as a unit.

Count the number of ways we can order the 5 couples in the row.

We can choose any one of the 5 couples for the first pair of seats, then any of the 4 remaining couples for the second pair of seats, then any of the 3 remaining couples for the third pair of seats, then one of the 2 remaining couples for the fourth pair of seats. The last couple must go into the last pair of seats.

Applying the product rule, this gives us $5 \cdot 4 \cdot 3 \cdot 2 \cdot 1$ options for ordering the couples.

Note that this equals 5 factorial, or $5!$. We'll go over factorials and counting sequences in lecture tomorrow

1 – Seating

- a) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if all couples are to get adjacent seats?

Applying the product rule, this gives us $5 \cdot 4 \cdot 3 \cdot 2 \cdot 1$ options for ordering the couples.

Now, apply the product rule to count the number of ways the people in each couple can be assigned to their seat pair. Each couple can be assigned to the seat pair in 2 different ways, and we have 5 couples, giving us 2^5 .

Use the product rule to combine the number of ways to order the couples with the number of ways to assign the people in the couple to their seats. Thus, the final answer is

$$5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \cdot 2^5$$

Or, more concisely,

$$5! \cdot 2^5$$

1 – Seating

- b) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if anyone can sit anywhere, except that one couple insists on not sitting in adjacent seats?

1 – Seating

- b) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if anyone can sit anywhere, except that one couple insists on not sitting in adjacent seats?

One way we can solve this is to consider by cases.

Name the two people in the couple A and B. There are two cases: A can sit on one of the ends, or not.

1 – Seating

- b) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if anyone can sit anywhere, except that one couple insists on not sitting in adjacent seats?

One way we can solve this is to consider by cases.

Name the two people in the couple A and B. There are two cases: A can sit on one of the ends, or not.

Case 1: If A sits on an end seat, A has 2 choices and B has 8 possible seats.

Case 2: If A doesn't sit on the end, A has 8 choices and B only has 7.

So, there are a total of $2 \cdot 8 + 8 \cdot 7$ ways A and B can sit.

1 – Seating

- b) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if anyone can sit anywhere, except that one couple insists on not sitting in adjacent seats?

One way we can solve this is to consider by cases.

Name the two people in the couple A and B. There are two cases: A can sit on one of the ends, or not.

Case 1: If A sits on an end seat, A has 2 choices and B has 8 possible seats.

Case 2: If A doesn't sit on the end, A has 8 choices and B only has 7.

So, there are a total of $2 \cdot 8 + 8 \cdot 7$ ways A and B can sit.

Once they do, the other 8 people can sit in $8!$ ways since there are no other restrictions.

1 – Seating

- b) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if anyone can sit anywhere, except that one couple insists on not sitting in adjacent seats?

One way we can solve this is to consider by cases.

Name the two people in the couple A and B. There are two cases: A can sit on one of the ends, or not.

Case 1: If A sits on an end seat, A has 2 choices and B has 8 possible seats.

Case 2: If A doesn't sit on the end, A has 8 choices and B only has 7.

So, there are a total of $2 \cdot 8 + 8 \cdot 7$ ways A and B can sit.

Once they do, the other 8 people can sit in $8!$ ways since there are no other restrictions. Hence the total number of ways is

$$(2 \cdot 8 + 8 \cdot 7)8! = 9 \cdot 8 \cdot 8! = 8 \cdot 9!$$

1 – Seating

- b) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if anyone can sit anywhere, except that one couple insists on not sitting in adjacent seats?

Another way to solve this is to apply complementary counting to first compute the total number of arrangements of the 10 people, and then subtract from this the number of arrangements in which that particular couple does get adjacent seats.

1 – Seating

- b) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if anyone can sit anywhere, except that one couple insists on not sitting in adjacent seats?

Another way to solve this is to apply complementary counting to first compute the total number of arrangements of the 10 people, and then subtract from this the number of arrangements in which that particular couple does get adjacent seats.

Total arrangements: $10!$

1 – Seating

- b) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if anyone can sit anywhere, except that one couple insists on not sitting in adjacent seats?

Another way to solve this is to apply complementary counting to first compute the total number of arrangements of the 10 people, and then subtract from this the number of arrangements in which that particular couple does get adjacent seats.

Total arrangements: $10!$

Ways for a particular couple to get adjacent seats. Since you can treat the couple as a unit, permute the 9 “individuals” (consisting of 8 people plus the couple) and then consider the 2 permutations for that couple: $9! \cdot 2$

1 – Seating

- b) How many ways are there to seat 10 people, consisting of 5 couples, in a row of 10 seats if anyone can sit anywhere, except that one couple insists on not sitting in adjacent seats?

Another way to solve this is to apply complementary counting to first compute the total number of arrangements of the 10 people, and then subtract from this the number of arrangements in which that particular couple does get adjacent seats.

Total arrangements: $10!$

Ways for a particular couple to get adjacent seats. Since you can treat the couple as a unit, permute the 9 “individuals” (consisting of 8 people plus the couple) and then consider the 2 permutations for that couple: $9! \cdot 2$

That means the answer to the question is

$$10! - 9! \cdot 2 = 9! (10 - 2) = 8 \cdot 9!$$

Problem 5 – Birthday Cake



5 – Birthday Cake

A chef is preparing desserts for the week, starting on a Sunday. On each day, only one of five desserts (apple pie, cherry pie, strawberry pie, pineapple pie, and cake) may be served. On Thursday there is a birthday, so cake must be served that day. On no two consecutive days can the chef serve the same dessert. How many dessert menus are there for the week?

Work on this problem with the people around you, and then we'll go over it together!

5 – Birthday Cake

A chef is preparing desserts for the week, starting on a Sunday. On each day, only one of five desserts (apple pie, cherry pie, strawberry pie, pineapple pie, and cake) may be served. On Thursday there is a birthday, so cake must be served that day. On no two consecutive days can the chef serve the same dessert. How many dessert menus are there for the week?

5 – Birthday Cake

A chef is preparing desserts for the week, starting on a Sunday. On each day, only one of five desserts (apple pie, cherry pie, strawberry pie, pineapple pie, and cake) may be served. On Thursday there is a birthday, so cake must be served that day. On no two consecutive days can the chef serve the same dessert. How many dessert menus are there for the week?

Apply the product rule. Start from Thursday and work forward and backward in the week:

5 – Birthday Cake

A chef is preparing desserts for the week, starting on a Sunday. On each day, only one of five desserts (apple pie, cherry pie, strawberry pie, pineapple pie, and cake) may be served. On Thursday there is a birthday, so cake must be served that day. On no two consecutive days can the chef serve the same dessert. How many dessert menus are there for the week?

Apply the product rule. Start from Thursday and work forward and backward in the week:

Given the 1 choice on Thursday, for each of Wednesday and Friday, there are 4 choices (the different pie options).

5 – Birthday Cake

A chef is preparing desserts for the week, starting on a Sunday. On each day, only one of five desserts (apple pie, cherry pie, strawberry pie, pineapple pie, and cake) may be served. On Thursday there is a birthday, so cake must be served that day. On no two consecutive days can the chef serve the same dessert. How many dessert menus are there for the week?

Apply the product rule. Start from Thursday and work forward and backward in the week:

Given the 1 choice on Thursday, for each of Wednesday and Friday, there are 4 choices (the different pie options).

Given the choice on Wednesday, there are 4 choices for Tuesday, and given the choice on Tuesday, there are 4 choices for Monday, and given the choice on Monday, there are 4 choices on Sunday. Similarly, given the choice on Friday, there are 4 choices on Saturday.

5 – Birthday Cake

A chef is preparing desserts for the week, starting on a Sunday. On each day, only one of five desserts (apple pie, cherry pie, strawberry pie, pineapple pie, and cake) may be served. On Thursday there is a birthday, so cake must be served that day. On no two consecutive days can the chef serve the same dessert. How many dessert menus are there for the week?

Apply the product rule. Start from Thursday and work forward and backward in the week:

Given the 1 choice on Thursday, for each of Wednesday and Friday, there are 4 choices (the different pie options).

Given the choice on Wednesday, there are 4 choices for Tuesday, and given the choice on Tuesday, there are 4 choices for Monday, and given the choice on Monday, there are 4 choices on Sunday. Similarly, given the choice on Friday, there are 4 choices on Saturday. Therefore the answer is

$$4 \cdot 4 \cdot 4 \cdot 4 \cdot 1 \cdot 4 \cdot 4 = 4^6$$

Problem 10 – Extended Family Portrait



10 – Extended Family Portrait

A group of n families, each with m members, are to be lined up for a photograph. In how many ways can the nm people be arranged if members of a family must stay together?

Work on this problem with the people around you, and then we'll go over it together!

10 – Extended Family Portrait

A group of n families, each with m members, are to be lined up for a photograph. In how many ways can the nm people be arranged if members of a family must stay together?

10 – Extended Family Portrait

A group of n families, each with m members, are to be lined up for a photograph. In how many ways can the nm people be arranged if members of a family must stay together?

Apply the product rule.

10 – Extended Family Portrait

A group of n families, each with m members, are to be lined up for a photograph. In how many ways can the nm people be arranged if members of a family must stay together?

Apply the product rule.

First order the families; there are $n!$ ways to do this.

10 – Extended Family Portrait

A group of n families, each with m members, are to be lined up for a photograph. In how many ways can the nm people be arranged if members of a family must stay together?

Apply the product rule.

First order the families; there are $n!$ ways to do this.

Then consider each of the n families one by one and reorder their members. Within each family, there are $m!$ ways to order their members.

10 – Extended Family Portrait

A group of n families, each with m members, are to be lined up for a photograph. In how many ways can the nm people be arranged if members of a family must stay together?

Apply the product rule.

First order the families; there are $n!$ ways to do this.

Then consider each of the n families one by one and reorder their members. Within each family, there are $m!$ ways to order their members.

So, the total number of ways to line these people up according to the given constraints is

$$n! (m!)^n$$

Problem 9 – Rabbits!



9 – Rabbits!

Rabbits Peter and Pauline have three offspring: Flopsie, Mopsie, and Cotton-tail. These five rabbits are to be distributed to four different pet stores so that no store gets both a parent and a child. It is not required that every store gets a rabbit. In how many different ways can this be done?

Work on this problem with the people around you, and then we'll go over it together!

9 – Rabbits!

Rabbits Peter and Pauline have three offspring: Flopsie, Mopsie, and Cotton-tail. These 5 rabbits are to be distributed to 4 different pet stores so that no store gets both a parent and a child. It is not required that every store gets a rabbit. In how many different ways can this be done?

9 – Rabbits!

Rabbits Peter and Pauline have three offspring: Flopsie, Mopsie, and Cotton-tail. These 5 rabbits are to be distributed to 4 different pet stores so that no store gets both a parent and a child. It is not required that every store gets a rabbit. In how many different ways can this be done?

Find the number of possibilities when Peter and Pauline go to the same store and find the number of possibilities when they go to different stores, and then use the sum rule to get the final answer.

If Peter and Pauline go to the same store, there are 4 stores it could be. For each such choice, there are 3 choices of store for each of the 3 offspring, so 3^3 choices for all the offspring. If Peter and Pauline go to different stores, there are $4 \cdot 3 = 12$ pairs of stores they could go to. For each such choice, there are 2 choices of store for each of the 3 offspring, so 2^3 choices for all the offspring. Therefore the answer is

9 – Rabbits!

Rabbits Peter and Pauline have three offspring: Flopsie, Mopsie, and Cotton-tail. These 5 rabbits are to be distributed to 4 different pet stores so that no store gets both a parent and a child. It is not required that every store gets a rabbit. In how many different ways can this be done?

Find the number of possibilities when Peter and Pauline go to the same store and find the number of possibilities when they go to different stores, and then use the sum rule to get the final answer.

If Peter and Pauline go to the same store, there are 4 stores it could be. For each such choice, there are 3 choices of store for each of the 3 offspring, so 3^3 choices for all the offspring. If Peter and Pauline go to different stores, there are $4 \cdot 3 = 12$ pairs of stores they could go to. For each such choice, there are 2 choices of store for each of the 3 offspring, so 2^3 choices for all the offspring. Therefore the answer is

$$4 \cdot 3^3 + 12 \cdot 2^3$$

Problem 4 – Escape the Professor



4 – Escape the Professor

There are 6 security professors and 7 theory professors taking part in an escape room. If 4 security professors and 4 theory professors are chosen and paired off, how many pairings are possible?

Work on this problem with the people around you, and then we'll go over it together!

4 – Escape the Professor

There are 6 security professors and 7 theory professors taking part in an escape room. If 4 security professors and 4 theory professors are chosen and paired off, how many pairings are possible?

4 – Escape the Professor

There are 6 security professors and 7 theory professors taking part in an escape room. If 4 security professors and 4 theory professors are chosen and paired off, how many pairings are possible?

Apply the product rule to first choose 4 of the security professors, then 4 of the theory professors. Then assign each theory professor to a security professor (4 choices for the first, 3 for the second and so on).

The answer is

4 – Escape the Professor

There are 6 security professors and 7 theory professors taking part in an escape room. If 4 security professors and 4 theory professors are chosen and paired off, how many pairings are possible?

Apply the product rule to first choose 4 of the security professors, then 4 of the theory professors. Then assign each theory professor to a security professor (4 choices for the first, 3 for the second and so on).

The answer is

$$\binom{6}{4} \binom{7}{4} 4!$$

Problem 11 - Subsubset



11 – Subsubset

Let $[n] = \{1, 2, \dots, n\}$ denote the first n natural numbers. How many (ordered) pairs of subsets (A, B) are there such that $A \subseteq B \subseteq [n]$?

Work on this problem with the people around you, and then we'll go over it together!

11 – Subsubset

Let $[n] = \{1, 2, \dots, n\}$ denote the first n natural numbers. How many (ordered) pairs of subsets (A, B) are there such that $A \subseteq B \subseteq [n]$?

11 – Subsubset

Let $[n] = \{1, 2, \dots, n\}$ denote the first n natural numbers. How many (ordered) pairs of subsets (A, B) are there such that $A \subseteq B \subseteq [n]$?

Realize that, if there are no restrictions, for each element i of $1, \dots, n$, there are four possibilities: it can be in only A , only B , both, or neither. In our case, there is only one that is not valid (violates $A \subseteq B$): being in A but not B . Hence there are 3 choices for each element, so the total number of such ordered pairs of subsets is

11 – Subsubset

Let $[n] = \{1, 2, \dots, n\}$ denote the first n natural numbers. How many (ordered) pairs of subsets (A, B) are there such that $A \subseteq B \subseteq [n]$?

Realize that, if there are no restrictions, for each element i of $1, \dots, n$, there are four possibilities: it can be in only A , only B , both, or neither. In our case, there is only one that is not valid (violates $A \subseteq B$): being in A but not B . Hence there are 3 choices for each element, so the total number of such ordered pairs of subsets is

$$3^n$$

That's All, Folks!

Thanks for coming to section this week!
Any questions?