Continuous Zoo CSE 312 Winter 25 Lecture 15

Let's start with the pmf

For discrete random variables, we defined the pmf: $p_Y(k) = \mathbb{P}(Y = k)$.

We can't have a pmf quite like we did for discrete random variables. Let *X* be a random real number between 0 and 1.

$$\mathbb{P}(X=.1)=\frac{1}{\infty}??$$

Let's try to maintain as many rules as we can...

Discrete	Continuous	
$p_Y(k) \ge 0$	$f_X(k) \ge 0$	
$\sum_{\omega} p_Y(\omega) = 1$	$\int_{-\infty}^{\infty} f_X(k) \mathrm{d}k = 1$	Use f_X instead of p_X to remember it's different .

Comparing Discrete and Continuous

		Discrete Random Variables	Continuous Random Variables
	Probability 0	Equivalent to impossible	All impossible events have probability 0, but not conversely.
	Relative Chances	PMF: $p_X(k) = \mathbb{P}(X = k)$	PDF $f_X(k)$ gives chances relative to $f_X(k')$
	Events	Sum over PMF to get probability	Integrate PDF to get probability
	Convert from CDF to PMF	Sum up PMF to get CDF. Look for "breakpoints" in CDF to get PMF.	Integrate PDF to get CDF. Differentiate CDF to get PDF.
	$\mathbb{E}[X]$	$\sum_{\omega} X(\omega) \cdot p_X(\omega)$	$\int_{-\infty}^{\infty} z \cdot f_X(z) \mathrm{d}z$
)	$\mathbb{E}[\boldsymbol{g}(\boldsymbol{X})]$	$\sum_{\omega} g(X(\omega)) \cdot p_X(\omega)$	$\int_{-\infty}^{\infty} g(z) \cdot f_X(z) \mathrm{d}z$
/	Var(X)	$\mathbb{E}[X^2] - (\mathbb{E}[X])^2$	$\mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \int_{-\infty}^{\infty} (z - \mathbb{E}[X])^2 f_X(z) \mathrm{d}z$

What about expectation?

For a random variable *X*, we define:





Just replace summing over the pmf with integrating the pdf.

It still represents the average value of X.

Expectation of a function

For any function g and any continuous random variable, X: $\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(X(z)) \cdot f_X(z) \, \mathrm{d}z$

Again, analogous to the discrete case; just replace summation with integration and pmf with the pdf.

We're going to treat this as a definition.

Technically, this is really a theorem; since f() is the pdf of X and it only gives relative likelihoods for X, we need a proof to guarantee it "works" for g(X).

Sometimes called "Law of the Unconscious Statistician."



Linearity of Expectation

Still true!

 $\mathbb{E}[aX + bY + c] = a\mathbb{E}[X] + b\mathbb{E}[Y] + c$ For all *X*, *Y*; even if they're continuous.

Won't show you the proof – for just $\mathbb{E}[aX + b]$, it's $\mathbb{E}[aX + b] = \int_{-\infty}^{\infty} [aX(k) + b] f_X(k) dk$

- $= \int_{-\infty}^{\infty} aX(k) f_X(k) dk + \int_{-\infty}^{\infty} bf_X(k) dk$
- $= a \int_{-\infty}^{\infty} X(k) f_X(k) dk + b \int_{-\infty}^{\infty} f_X(k) dk$
- $= a\mathbb{E}[X] + b$







Let's calculate an expectation $f_X(z) = \begin{cases} \frac{1}{b-a} & \text{if } a \leq z \leq b \\ 0 & \text{otherwise} \end{cases}$ Let X be a uniform random number between a and b. $\mathbb{E}[X] = \int_{-\infty}^{\infty} z \cdot f_X(z) \, \mathrm{d}z$ $= \int_{-\infty}^{a} z \cdot \underbrace{0}_{a} dz + \int_{a}^{b} z \cdot \frac{1}{b-a} dz + \int_{b}^{\infty} z \cdot \underbrace{0}_{a} dz$ $= 0 + \int_a^b \frac{z}{b-a} \, \mathrm{d}z + 0$ $= \frac{z^2}{2(b-a)} \bigg|_{z=a}^{b} = \frac{b^2}{2(b-a)} - \frac{a^2}{2(b-a)} = \frac{b^2 - a^2}{2(b-a)} = \frac{(b+a)(b-a)}{2(b-a)}$



Let's assemble the variance



Continuous Uniform Distribution

X~Unif(*a*, *b*) (uniform real number between *a* and *b*)

 $\begin{array}{l} \label{eq:pdf} \mathsf{PDF:} \ f_X(k) = \begin{cases} \frac{1}{b-a} & \text{if } a \leq k \leq b \\ 0 & \text{otherwise} \end{cases} \end{cases}$ $\mathsf{CDF:} \ F_X(k) = \begin{cases} 0 & \text{if } k < a \\ \frac{k-a}{b-a} & \text{if } a \le k \le b \\ 1 & \text{if } k \ge b \end{cases}$ $\mathbb{E}[X] = \frac{a+b}{2}$ $Var(X) = \frac{(b-a)^2}{12}$

Continuous Zoo



Exponential Random Variable

Like a geometric random variable, but continuous time. How long do we wait until an event happens? (instead of "how many flips until a heads")

Where waiting doesn't make the event happen any sooner.

Geometric:
$$\mathbb{P}(X = k + 1 | X \ge 1) = \mathbb{P}(X = k)$$

When the first flip is tails, the coin doesn't remember it came up tails, you've made no progress.

For an exponential random variable:

 $\mathbb{P}(X \ge k+1 \mid X \ge 1) = \mathbb{P}(Y \ge k)$

Are these memoryless?

You arrive to a bus stop at a (uniformly) random time, to a bus that arrives every 10 minutes. How long until the bus arrives? How long conditioned on you've already waited 8 minutes?

You put everyone in class into a random order. You'll iterate through that list. What is the probability of being next? Probability of being next conditioned on not selected yet AND half the class has gone?

You flip a coin (independently) until you see a heads. How many flips do you need? How many additional flips after seeing 4 tails?

Are these memoryless?

You arrive to a bus stop at a (uniformly) random time, to a bus that arrives every 10 minutes. How long until the bus arrives? How long conditioned on you've already waited 8 minutes?

Not memoryless! (bus must arrive in 10 minutes total, must be soon!)

You put everyone in class into a random order. You'll iterate through that list. What is the probability of being next? Probability of being next conditioned on not selected yet AND half the class has gone?

Not memoryless (1/n of being first 1/(n/2) after half class gone)

You flip a coin (independently) until you see a heads

Memoryless!

A continuous memoryless RV?

Poisson random variables come from a memoryless-type process. Number of earthquakes (people in bakery, days with snow) would be memoryless under assumption that events are independent of each other!

Same experiments, but now ask a different question:

Poisson: how many incidents occur in fixed interval?

Exponential: how long do I have to wait to see the next incident?

Exponential random variable

If you take a Poisson random variable and ask "what's the time until the next event" you get an exponential distribution!

Let's find the CDF for an exponential.

Let $Y \sim \text{Exp}(\lambda)$, be the time until the first event, when we see an average of λ events per time unit.

What's $\mathbb{P}(Y > t)$? What Poisson are we waiting on, and what event for it tells you that Y > t?

Exponential random variable

If you take a Poisson random variable and ask "what's the time until the next event" you get an exponential distribution!

Let's find the CDF for an exponential.

Let $Y \sim \text{Exp}(\lambda)$, be the time until the first event, when we see an average of λ events per time unit. What's $\mathbb{P}(Y > t)$?

What Poisson are we waiting on? For $X \sim Poi(\lambda t) \mathbb{P}(Y > t) = \mathbb{P}(X = 0)$

$$\mathbb{P}(X=0) = \frac{(\lambda t)^0 e^{-\lambda t}}{0!} = e^{-\lambda t}$$

$$F_Y(t) = \mathbb{P}(Y \le t) = 1 - e^{-\lambda t} \text{ (for } t \ge 0, F_Y(x) = 0 \text{ for } x < 0)$$

Where did the *t* come from?

Why did we switch from $Exp(\lambda)$ to $Poi(\lambda t)$?

Let's make our units "incidents/second", so $\lambda = 3$ says we average 3 incidents per second.

What if I want to know the probability of waiting at least 5 seconds? Well then on average how many incidents do we see in a 5 second period?

15 incidents! So the Poisson (how many incidents in fixed interval) now refers to a larger interval, so averages more events; specifically λt .

Find the density

We know the CDF, $F_Y(t) = \mathbb{P}(Y \le t) = 1 - e^{-\lambda t}$ What's the density?

 $f_Y(t) =$

Find the density

We know the CDF, $F_Y(t) = \mathbb{P}(Y \le t) = 1 - e^{-\lambda t}$

What's the density?

$$f_Y(t) = \frac{d}{dt} \left(1 - e^{-\lambda t} \right) = 0 - \frac{d}{dt} \left(e^{-\lambda t} \right) = \lambda e^{-\lambda t}.$$

For $t \ge 0$ it's that expression For t < 0 it's just 0.

Exponential PDF



Memorylessness

$$\mathbb{P}(X \ge k+1 | X \ge 1) = \frac{\mathbb{P}(X \ge k+1 \cap X \ge 1)}{\mathbb{P}(X \ge 1)} = \frac{\mathbb{P}(X \ge k+1)}{1 - (1 - e^{-\lambda \cdot 1})}$$
$$= \frac{e^{-\lambda(k+1)}}{e^{-\lambda}} = e^{-\lambda k}$$

What about $\mathbb{P}(X \ge k)$ (without conditioning on the first step)? $1 - (1 - e^{-\lambda k}) = e^{-\lambda k}$

It's the same!!!

More generally, for an exponential $r \vee X$, $\mathbb{P}(X \ge s + t | X \ge s) = \mathbb{P}(X \ge t)$

Side note

I hid a trick in that algebra,

 $\mathbb{P}(X \ge 1) = 1 - \mathbb{P}(X < 1) = 1 - \mathbb{P}(X \le 1)$

The first step is the complementary law.

The second step is using that $\int_1^1 f_X(z) dz = 0$

In general, for continuous random variables we can switch out \leq and < without anything changing. We can't make those switches for discrete random variables.

Expectation of an exponential

Let $X \sim \text{Exp}(\lambda)$ $\mathbb{E}[X] = \int_{-\infty}^{\infty} z \cdot f_X(z) \mathrm{d}z$ $=\int_0^\infty z\cdot\lambda e^{-\lambda z}\,dz$ Let u = z: $dv = \lambda e^{-\lambda z} dz$ ($v = -e^{-\lambda z}$) Integrate by parts: $-ze^{-\lambda z} - \int -e^{-\lambda z} dz = -ze^{-\lambda z} - \frac{1}{2}e^{-\lambda z}$ Definite Integral: $-ze^{-\lambda z} - \frac{1}{\lambda}e^{-\lambda z}|_{z=0}^{\infty} = (\lim_{z \to \infty} -ze^{-\lambda z} - \frac{1}{\lambda}e^{-\lambda z}) - (0 - \frac{1}{\lambda})$ By L'Hopital's Rule $(\lim_{z \to \infty} -\frac{z}{e^{\lambda z}} - \frac{1}{\lambda e^{\lambda z}}) - (0 - \frac{1}{\lambda}) = (\lim_{z \to \infty} -\frac{1}{\lambda e^{\lambda z}}) + \frac{1}{\lambda} = \frac{1}{\lambda}$

Don't worry about the derivation (it's here if you're interested; you're not responsible for the derivation. Just the value.

Variance of an exponential

If X~Exp(λ)then Var(X) = $\frac{1}{\lambda^2}$

Similar calculus tricks will get you there.

Exponential

 $X \sim \text{Exp}(\lambda)$ Parameter $\lambda \geq 0$ is the average number of events in a unit of time. $f_X(k) = \begin{cases} \lambda e^{-\lambda k} & \text{if } k \ge 0\\ 0 & \text{otherwise} \end{cases}$