

Discrete Structures

Graphs

Chapter 7, Sections 7.1 - 7.3

Dieter Fox

Undirected Graphs

- ◇ A **simple graph** $G = (V, E)$ consists of V , a nonempty set of **vertices**, and E , a set of unordered pairs of distinct elements of V called **edges**.
- ◇ A **multigraph** $G = (V, E)$ consists of a set V of vertices, a set E of edges, and a function f from E to $\{\{u, v\} \mid u, v \in V, u \neq v\}$. The edges e_1 and e_2 are called **multiple** or **parallel edges** if $f(e_1) = f(e_2)$.
- ◇ A **pseudograph** $G = (V, E)$ consists of a set V of vertices, a set E of edges, and a function f from E to $\{\{u, v\} \mid u, v \in V\}$. An edge is a **loop** if $f(e) = \{u, u\} = \{u\}$ for some $u \in V$.

Directed Graphs

- ◇ A **directed graph** $G = (V, E)$ consists of a set V of vertices and a set of edges E that are ordered pairs of elements of V .
- ◇ A **directed multigraph** $G = (V, E)$ consists of a set V of vertices, a set E of edges, and a function f from E to $\{(u, v) \mid u, v \in V\}$. The edges e_1 and e_2 are **multiple edges** if $f(e_1) = f(e_2)$.

Undirected Graph Terminology

- ◇ Two vertices u and v in an undirected graph G are called **adjacent** (or **neighbors**) in G if $\{u, v\}$ is an edge of G . If $e = \{u, v\}$, the edge e is called **incident with** the vertices u and v . The edge e is also said to **connect** u and v . The vertices u and v are called **endpoints** of the edges $\{u, v\}$.
- ◇ The **degree** of a vertex in an undirected graph is the number of edges incident with it, except that a loop at a vertex contributes twice to the degree of that vertex. The degree of the vertex v is denoted by $\deg(v)$.
- ◇ **The Handshaking Theorem** : Let $G = (V, E)$ be an undirected graph with e edges. Then
$$2e = \sum_{v \in V} \deg(v).$$
- ◇ **Theorem** : An undirected graph has an even number of vertices of odd degree.

Directed Graph Terminology

- ◇ When (u, v) is an edge of the graph G with directed edges, u is said to be **adjacent to** v and v is said to be **adjacent from** u . The vertex u is called the **initial vertex** of (u, v) , and v is called the **terminal** or **end vertex** of (u, v) . The initial vertex and terminal vertex of a loop are the same.
- ◇ In a graph with directed edges the **in-degree** of a vertex v , denoted by $\deg^-(v)$, is the number of edges with v as their terminal vertex. The **out-degree** of v , denoted by $\deg^+(v)$, is the number of edges with v as their initial vertex.
- ◇ **Theorem:** Let $G = (V, E)$ be a graph with directed edges. Then

$$\sum_{v \in V} \deg^-(v) = \sum_{v \in V} \deg^+(v) = |E|.$$

More Definitions ...

- ◇ A simple graph G is called **bipartite** if its vertex V can be partitioned into two disjoint nonempty sets V_1 and V_2 such that every edge in the graph connects a vertex in V_1 and a vertex in V_2 (so that no edge in G connects either two vertices in V_1 or two vertices in V_2).
- ◇ A **subgraph** of a graph $G = (V, E)$ is a graph $H = (W, F)$ where $W \subseteq V$ and $F \subseteq E$.
- ◇ The **union** of two simple graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ is the simple graph with vertex set $V_1 \cup V_2$ and edge set $E_1 \cup E_2$. The union of G_1 and G_2 is denoted by $G_1 \cup G_2$.
- ◇ The simple graphs $G_1 = (V_1, E_1)$ and $G_2 = (V_2, E_2)$ are **isomorphic** if there is a one-to-one and onto function f from V_1 to V_2 with the property that a and b are adjacent in G_1 if and only if $f(a)$ and $f(b)$ are adjacent in G_2 , for all a and b in V_1 . Such a function f is called an **isomorphism**.