

# Discrete Structures

## Graphs

---

Chapter 9, Sections 9.1 - 9.5

*Dieter Fox*

# Undirected Graphs

---

- ◇ A **simple graph**  $G = (V, E)$  consists of  $V$ , a nonempty set of **vertices**, and  $E$ , a set of unordered pairs of distinct elements of  $V$  called **edges**.
- ◇ A **multigraph**  $G = (V, E)$  consists of a set  $V$  of vertices, a set  $E$  of edges, and a function  $f$  from  $E$  to  $\{\{u, v\} \mid u, v \in V, u \neq v\}$ . The edges  $e_1$  and  $e_2$  are called **multiple** or **parallel edges** if  $f(e_1) = f(e_2)$ .
- ◇ A **pseudograph**  $G = (V, E)$  consists of a set  $V$  of vertices, a set  $E$  of edges, and a function  $f$  from  $E$  to  $\{\{u, v\} \mid u, v \in V\}$ . An edge is a **loop** if  $f(e) = \{u, u\} = \{u\}$  for some  $u \in V$ .

# Directed Graphs

---

- ◇ A **directed graph**  $G = (V, E)$  consists of a set  $V$  of vertices and a set of edges  $E$  that are ordered pairs of elements of  $V$ .
- ◇ A **directed multigraph**  $G = (V, E)$  consists of a set  $V$  of vertices, a set  $E$  of edges, and a function  $f$  from  $E$  to  $\{(u, v) \mid u, v \in V\}$ . The edges  $e_1$  and  $e_2$  are **multiple edges** if  $f(e_1) = f(e_2)$ .

# Undirected Graph Terminology

---

◇ Two vertices  $u$  and  $v$  in an undirected graph  $G$  are called **adjacent** (or **neighbors**) in  $G$  if  $\{u, v\}$  is an edge of  $G$ . If  $e = \{u, v\}$ , the edge  $e$  is called **incident with** the vertices  $u$  and  $v$ . The edge  $e$  is also said to **connect**  $u$  and  $v$ . The vertices  $u$  and  $v$  are called **endpoints** of the edges  $\{u, v\}$ .

◇ The **degree** of a vertex in an undirected graph is the number of edges incident with it, except that a loop at a vertex contributes twice to the degree of that vertex. The degree of the vertex  $v$  is denoted by  $\deg(v)$ .

◇ **The Handshaking Theorem** : Let  $G = (V, E)$  be an undirected graph with  $e$  edges. Then

$$2e = \sum_{v \in V} \deg(v).$$

◇ **Theorem** : An undirected graph has an even number of vertices of odd degree.

# Directed Graph Terminology

---

- ◇ When  $(u, v)$  is an edge of the graph  $G$  with directed edges,  $u$  is said to be **adjacent to**  $v$  and  $v$  is said to be **adjacent from**  $u$ . The vertex  $u$  is called the **initial vertex** of  $(u, v)$ , and  $v$  is called the **terminal** or **end vertex** of  $(u, v)$ . The initial vertex and terminal vertex of a loop are the same.
- ◇ In a graph with directed edges the **in-degree** of a vertex  $v$ , denoted by  $\deg^-(v)$ , is the number of edges with  $v$  as their terminal vertex. The **out-degree** of  $v$ , denoted by  $\deg^+(v)$ , is the number of edges with  $v$  as their initial vertex.
- ◇ **Theorem:** Let  $G = (V, E)$  be a graph with directed edges. Then

$$\sum_{v \in V} \deg^-(v) = \sum_{v \in V} \deg^+(v) = |E|.$$

## More Definitions ...

---

- ◇ A simple graph  $G$  is called **bipartite** if its vertex  $V$  can be partitioned into two disjoint nonempty sets  $V_1$  and  $V_2$  such that every edge in the graph connects a vertex in  $V_1$  and a vertex in  $V_2$  (so that no edge in  $G$  connects either two vertices in  $V_1$  or two vertices in  $V_2$ ).
- ◇ A **subgraph** of a graph  $G = (V, E)$  is a graph  $H = (W, F)$  where  $W \subseteq V$  and  $F \subseteq E$ .
- ◇ The **union** of two simple graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  is the simple graph with vertex set  $V_1 \cup V_2$  and edge set  $E_1 \cup E_2$ . The union of  $G_1$  and  $G_2$  is denoted by  $G_1 \cup G_2$ .
- ◇ The simple graphs  $G_1 = (V_1, E_1)$  and  $G_2 = (V_2, E_2)$  are **isomorphic** if there is a one-to-one and onto function  $f$  from  $V_1$  to  $V_2$  with the property that  $a$  and  $b$  are adjacent in  $G_1$  if and only if  $f(a)$  and  $f(b)$  are adjacent in  $G_2$ , for all  $a$  and  $b$  in  $V_1$ . Such a function  $f$  is called an **isomorphism**.