

---

# CSE 331

## Software Design & Implementation

Spring 2020

Section 3 – HW4, Abstract Data Types, and JUnit

---

# Agenda

---

- Overview of HW4
- Quick review of polynomial arithmetic
- Review abstract data types (ADTs) by example
- Unit testing with Junit – an initial tour for HW4

# HW4 – Polynomial calculator

---

A homework in 6 parts:

0. Pseudocode algorithms for polynomial arithmetic
1. Conceptual questions about `RatNum`
2. Implement `RatTerm`
3. Implement `RatPoly`
4. Implement `RatPolyStack`
5. Try out your finished calculator!
6. Run your code against our tests to make sure it works!



# The RatThings

---

- **RatNum** ADT
  - A rational number
  - Also includes a NaN (“not a number”) value
- **RatTerm** ADT
  - A polynomial term (rational coefficient w/ integer degree)
- **RatPoly** ADT
  - A polynomial expression (sum of polynomial terms)
- **RatPolyStack** ADT
  - An ordered collection of polynomial expressions



# The RatThings

---


**RatPoly** →  $\frac{1}{4}x^2 + 3x + 10$

**RatTerm** →  $\frac{2}{3}x^3$

**RatNum** →  $\frac{1}{7}$

# The RatThings

---



# Polynomial arithmetic

---

Review arithmetic operations over polynomial expressions:

1. Addition
2. Subtraction
3. Multiplication
4. Division

# Polynomial addition

---

$$(5x^4 + 4x^3 - x^2 + 5) + (3x^5 - 2x^3 + x - 5)$$



# Polynomial addition

---

$$- 3x^5 + 5x^4 + 4x^3 - x^2 + 5(2x^3 + x) - 5$$

$$\begin{array}{r}
 5x^4 + 4x^3 - 1x^2 + 5 \\
 + 3x^5 - 2x^3 + 1x - 5 \\
 \hline
 \end{array}$$

# Polynomial addition

---

$$(5x^4 + 4x^3 - x^2 + 5) + (3x^5 - 2x^3 + x - 5)$$

$$\begin{array}{r} 0x^5 \quad 5x^4 + \quad 4x^3 + \quad 1x^2 - \quad + \quad 0x \quad + \quad 5 \\ + \quad 3x^5 + \quad 0x^4 - \quad 2x^3 + \quad 0x^2 + \quad 1x - \quad 5 \\ \hline \end{array}$$

# Polynomial addition

---

$$(5x^4 + 4x^3 - x^2 + 5) + (3x^5 - 2x^3 + x - 5)$$

$$\begin{array}{r} 0x^5 + 5x^4 + 4x^3 - 1x^2 + 0x + 5 \\ + 3x^5 + 0x^4 - 2x^3 + 0x^2 + 1x - 5 \\ \hline 3x^5 + 5x^4 + 2x^3 - 1x^2 + 1x + 0 \end{array}$$

# Polynomial subtraction

---

$$(5x^4 + 4x^3 - x^2 + 5) - (3x^5 - 2x^3 + x - 5)$$

# Polynomial subtraction

---

$$(5x^4 + 4x^3 - x^2 + 5) - (3x^5 - 2x^3 + x - 5)$$

$$\begin{array}{r} 5x^4 + 4x^3 - 1x^2 + 5 \\ - 3x^5 - 2x^3 + 1x - 5 \\ \hline \end{array}$$

# Polynomial subtraction

---

$$(5x^4 + 4x^3 - x^2 + 5) - (3x^5 - 2x^3 + x - 5)$$

$$\begin{array}{r} 0x^5 + 5x^4 + 4x^3 - 1x^2 + 0x + 5 \\ - 3x^5 + 0x^4 - 2x^3 + 0x^2 + 1x - 5 \\ \hline \end{array}$$

# Polynomial subtraction

---

$$(5x^4 + 4x^3 - x^2 + 5) - (3x^5 - 2x^3 + x - 5)$$

$$\begin{array}{r} 0x^5 + 5x^4 + 4x^3 - 1x^2 + 0x + 5 \\ - 3x^5 + 0x^4 - 2x^3 + 0x^2 + 1x - 5 \\ \hline -3x^5 + 5x^4 + 6x^3 - 1x^2 - 1x + 10 \end{array}$$

# Polynomial multiplication

---

$$(4x^3 - x^2 + 5) \times (x - 5)$$



# Polynomial multiplication

---

$$(4x^3 - x^2 + 5) \times (x - 5)$$

$$\begin{array}{r} 4x^3 - x^2 + 5 \\ \times \phantom{4x^3 - x^2 + 5} \\ \hline 1x - 5 \end{array}$$

# Polynomial multiplication

---

$$(4x^3 - x^2 + 5) \times (x - 5)$$

$$\begin{array}{r} \phantom{\times} \phantom{4x^3} - \phantom{1x^2} \phantom{+} \phantom{5} \\ \times \phantom{4x^3} \phantom{-} \phantom{1x^2} \phantom{+} \phantom{5} \\ \hline \phantom{4x^3} - 20x^3 + 5x^2 \phantom{+} \phantom{5} \\ \phantom{4x^3} \phantom{-} \phantom{1x^2} \phantom{+} \phantom{5} \phantom{1x} - 5 \\ \phantom{4x^3} \phantom{-} \phantom{1x^2} \phantom{+} \phantom{5} \phantom{1x} \phantom{-} 25 \end{array}$$

# Polynomial multiplication

---

$$(4x^3 - x^2 + 5) \times (x - 5)$$

$$\begin{array}{r} \phantom{\times} \phantom{4x^3} - \phantom{1x^2} \phantom{+} \phantom{5} \\ \times \phantom{4x^3} - \phantom{1x^2} \phantom{+} \phantom{5} \\ \hline \phantom{4x^4} - \phantom{1x^3} \phantom{+} \phantom{5x} \phantom{-} \phantom{25} \\ \phantom{4x^4} - \phantom{1x^3} \phantom{+} \phantom{5x} \phantom{-} \phantom{25} \\ \phantom{4x^4} - \phantom{1x^3} \phantom{+} \phantom{5x} \phantom{-} \phantom{25} \end{array}$$

The diagram shows the multiplication of  $(4x^3 - x^2 + 5)$  by  $(x - 5)$ . The terms  $4x^3$  and  $-1x^2$  are in red, and  $1x$  and  $-5$  are in blue. A green circle highlights the  $1x$  term in the second polynomial. The partial products are shown below a horizontal line:  $-20x^3 + 5x^2$  (from  $4x^3 \times -5$  and  $-1x^2 \times -5$ ) and  $4x^4 - 1x^3 + 5x$  (from  $4x^3 \times x$  and  $-1x^2 \times x$  and  $5 \times x$ ). The final result is  $4x^4 - 1x^3 - 20x^2 + 5x + 25$ .



# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r|l} 1x^3 & 5x^6 \\ -2x & +4x^4 \\ -5 & -1x^3 \\ \hline & +5 \end{array}$$

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r|l} 1x^3 + 0x^2 - 2x - 5 & 5x^6 + 0x^5 + 4x^4 - 1x^3 + 0x^2 + 0x + 5 \end{array}$$

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} 5x^3 \\ 1x^3 + 0x^2 - 2x - 5 \overline{) 5x^6 + 0x^5 + 4x^4 - 1x^3 + 0x^2 + 0x + 5} \end{array}$$



# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} 5x^3 \\ 1x^3 + 0x^2 - 2x - 5 \overline{) 5x^6 + 0x^5 + 4x^4 - 1x^3 + 0x^2 + 0x + 5} \\ \underline{5x^6 + 0x^5 - 10x^4 - 25x^3} \end{array}$$

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} 5x^3 \\ 1x^3 + 0x^2 - 2x - 5 \overline{) 5x^6 + 0x^5 + 4x^4 - 1x^3 + 0x^2 + 0x + 5} \\ - 5x^6 + 0x^5 - 10x^4 - 25x^3 \\ \hline 0x^6 + 0x^5 + 14x^4 + 24x^3 \end{array}$$

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} 5x^3 \\ 1x^3 + 0x^2 - 2x - 5 \overline{) 5x^6 + 0x^5 + 4x^4 - 1x^3 + 0x^2 + 0x + 5} \\ \underline{- 5x^6 + 0x^5 - 10x^4 - 25x^3} \\ 0x^6 + 0x^5 + 14x^4 + 24x^3 \end{array}$$

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$\begin{array}{r} 5x^3 \\ 1x^3 + 0x^2 - 2x - 5 \overline{) 5x^6 + 0x^5 + 4x^4 - 1x^3 + 0x^2 + 0x + 5} \\ - 5x^6 + 0x^5 - 10x^4 - 25x^3 \\ \hline 0x^6 + 0x^5 + 14x^4 + 24x^3 + 0x^2 + 0x + 5 \end{array}$$

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

				$5x^3$	$+0x^2$					
$1x^3$	$+0x^2$	$-2x$	$-5$	$5x^6$	$+0x^5$	$+4x^4$	$-1x^3$	$+0x^2$	$+0x$	$+5$
			$-$	$5x^6$	$+0x^5$	$-10x^4$	$-25x^3$			
				$0x^6$	$+0x^5$	$+14x^4$	$+24x^3$	$+0x^2$		

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

				$5x^3$	$+0x^2$					
$1x^3$	$+0x^2$	$-2x$	$-5$	$5x^6$	$+0x^5$	$+4x^4$	$-1x^3$	$+0x^2$	$+0x$	$+5$
			$-$	$5x^6$	$+0x^5$	$-10x^4$	$-25x^3$			
				$0x^6$	$+0x^5$	$+14x^4$	$+24x^3$	$+0x^2$	$+0x$	

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

				$5x^3$	$+0x^2$	$+14x$		
$1x^3 + 0x^2 - 2x - 5$	-	$5x^6$	$+0x^5$	$+4x^4$	$-1x^3$	$+0x^2$	$+0x$	$+5$
		-	$5x^6$	$+0x^5$	$-10x^4$	$-25x^3$		
		$0x^6$	$+0x^5$	$+14x^4$	$+24x^3$	$+0x^2$	$+0x$	

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

				$5x^3$	$+0x^2$	$+14x$	
$1x^3 + 0x^2 - 2x - 5$	$5x^6$	$+0x^5$	$+4x^4$	$-1x^3$	$+0x^2$	$+0x$	$+5$
	$- 5x^6$	$+0x^5$	$-10x^4$	$-25x^3$			
	$0x^6$	$+0x^5$	$+14x^4$	$+24x^3$	$+0x^2$	$+0x$	
			$14x^4$	$+0x^3$	$-28x^2$	$-70x$	



# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

				$5x^3$	$+0x^2$	$+14x$	
$1x^3 + 0x^2 - 2x - 5$	$5x^6$	$+0x^5$	$+4x^4$	$-1x^3$	$+0x^2$	$+0x$	$+5$
	$- 5x^6$	$+0x^5$	$-10x^4$	$-25x^3$			
	$0x^6$	$+0x^5$	$+14x^4$	$+24x^3$	$+0x^2$	$+0x$	
		$- 14x^4$	$+0x^3$	$-28x^2$	$-70x$		
		$0x^4$	$+24x^3$	$+28x^2$	$+70x$		

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

				$5x^3$	$+0x^2$	$+14x$	
$1x^3 + 0x^2 - 2x - 5$	$5x^6$	$+0x^5$	$+4x^4$	$-1x^3$	$+0x^2$	$+0x$	$+5$
	$- 5x^6$	$+0x^5$	$-10x^4$	$-25x^3$			
	$0x^6$	$+0x^5$	$+14x^4$	$+24x^3$	$+0x^2$	$+0x$	
			$- 14x^4$	$+0x^3$	$-28x^2$	$-70x$	
			$0x^4$	$+24x^3$	$+28x^2$	$+70x$	

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

				$5x^3$	$+0x^2$	$+14x$		
$1x^3 + 0x^2 - 2x - 5$	$5x^6$	$+0x^5$	$+4x^4$	$-1x^3$	$+0x^2$	$+0x$	$+5$	
	$-$	$5x^6$	$+0x^5$	$-10x^4$	$-25x^3$			
		$0x^6$	$+0x^5$	$+14x^4$	$+24x^3$	$+0x^2$	$+0x$	
			$-$	$14x^4$	$+0x^3$	$-28x^2$	$-70x$	
				$0x^4$	$+24x^3$	$+28x^2$	$+70x$	$+5$

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

				$5x^3$	$+0x^2$	$+14x$	$+24$
$1x^3 + 0x^2 - 2x - 5$	5x <sup>6</sup>	+0x <sup>5</sup>	+4x <sup>4</sup>	-1x <sup>3</sup>	+0x <sup>2</sup>	+0x	+5
	-	5x <sup>6</sup>	+0x <sup>5</sup>	-10x <sup>4</sup>	-25x <sup>3</sup>		
		0x <sup>6</sup>	+0x <sup>5</sup>	+14x <sup>4</sup>	+24x <sup>3</sup>	+0x <sup>2</sup>	+0x
			-	14x <sup>4</sup>	+0x <sup>3</sup>	-28x <sup>2</sup>	-70x
				0x <sup>4</sup>	$+24x^3$	$+28x^2$	$+70x$
					$+5$		

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

				$5x^3$	$+0x^2$	$+14x$	$+24$
$1x^3 + 0x^2 - 2x - 5$	$5x^6$	$+0x^5$	$+4x^4$	$-1x^3$	$+0x^2$	$+0x$	$+5$
-	$5x^6$	$+0x^5$	$-10x^4$	$-25x^3$			
	$0x^6$	$+0x^5$	$+14x^4$	$+24x^3$	$+0x^2$	$+0x$	
		-	$14x^4$	$+0x^3$	$-28x^2$	$-70x$	
			$0x^4$	$+24x^3$	$+28x^2$	$+70x$	$+5$
				$24x^3$	$+0x^2$	$-48x$	$-120$

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

				<b>5x<sup>3</sup></b>	<b>+0x<sup>2</sup></b>	<b>+14x</b>	<b>+24</b>
<b>1x<sup>3</sup> +0x<sup>2</sup> -2x -5</b>	5x <sup>6</sup>	+0x <sup>5</sup>	+4x <sup>4</sup>	-1x <sup>3</sup>	+0x <sup>2</sup>	+0x	+5
	-	5x <sup>6</sup>	+0x <sup>5</sup>	-10x <sup>4</sup>	-25x <sup>3</sup>		
		0x <sup>6</sup>	+0x <sup>5</sup>	+14x <sup>4</sup>	+24x <sup>3</sup>	+0x <sup>2</sup>	+0x
			-	14x <sup>4</sup>	+0x <sup>3</sup>	-28x <sup>2</sup>	-70x
				0x <sup>4</sup>	+24x <sup>3</sup>	+28x <sup>2</sup>	+70x +5
				-	24x <sup>3</sup>	+0x <sup>2</sup>	-48x -120
					<b>0x<sup>3</sup></b>	<b>+28x<sup>2</sup></b>	<b>+118x +125</b>

# Polynomial division

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

					quotient	
				5x <sup>3</sup>	+0x <sup>2</sup>	+14x
				+24		
1x <sup>3</sup> + 0x <sup>2</sup> - 2x - 5	5x <sup>6</sup>	+0x <sup>5</sup>	+4x <sup>4</sup>	-1x <sup>3</sup>	+0x <sup>2</sup>	+0x
	- 5x <sup>6</sup>	+0x <sup>5</sup>	-10x <sup>4</sup>	-25x <sup>3</sup>		
	0x <sup>6</sup>	+0x <sup>5</sup>	+14x <sup>4</sup>	+24x <sup>3</sup>	+0x <sup>2</sup>	+0x
			- 14x <sup>4</sup>	+0x <sup>3</sup>	-28x <sup>2</sup>	-70x
				0x <sup>4</sup>	+24x <sup>3</sup>	+28x <sup>2</sup>
					+70x	+5
					- 24x <sup>3</sup>	+0x <sup>2</sup>
					-48x	-120
					0x <sup>3</sup>	+28x <sup>2</sup>
						+118x
						+125
						remainder

# Polynomial division

---

$$(5x^6 + 4x^4 - x^3 + 5) / (x^3 - 2x - 5)$$

$$5x^3 + 14x + 24 + \frac{28x^2 + 118x + 125}{x^3 - 2x - 5}$$



# Abstract data types by example

---

Review ADT concepts through two examples:

- A **Line** ADT
- A **Rectangle** ADT


On the course website, see “Resources” → “Class and Method Specifications” for a handy guide with full details.

We won't cover representation invariants today (see Friday's lecture).

# Line ADT

---


Concept: A line segment in the Cartesian co-ordinate plane



# Line ADT: Representation #1

---

```
/**
 * A Line is a mutable 2D line segment with endpoints
 * p1 and p2.
 */
public class Line {
    // AF(this) = line with endpoints p1 and p2
    private Point p1, p2;
}
```




# Line ADT: Representation #2

---

```
/**
 * A Line is a mutable 2D line segment with endpoints
 * p1 and p2.
 */
public class Line {
    // AF(this) = line with endpoints (x1, y1) and (x2, y2)
    private int x1, x2;
    private int y1, y2;
}
```

Does this representation have any advantages over #1?




# Line ADT: Representation #3

---

```
/**
 * A Line is a mutable 2D line segment with endpoints
 * p1 and p2.
 */
public class Line {
    // AF(this) = line with endpoints (x1, y1) and
    // (x1 + len * cos(angle), y1 + len * sin(angle))
    private int x1, y1;
    private double angle;
    private double len;
}
```

Does this representation have any advantages over #1?



# Try it yourself!

---

Write your own specification of a Rectangle ADT on the handout.

Then give two different possible representations for your Rectangle ADT and write abstraction functions for them

# Testing: A quick introduction

---

- For HW 4, you'll be running our test suite to verify your RatThings work.
- Let's do a quick walkthrough of our test suite
  - Just know how it works; don't need to know how to write tests!

# JUnit

---

- Industry-standard Java toolkit for unit testing
  - We're using JUnit 4
- A unit test is a test for one “component” by itself
  - “Component” typically a class or a method
- Each unit test written as a method
  - We'll see the particulars in a moment...
- Closely related unit tests should be grouped into a class
  - For example, all unit tests for the same ADT implementation



# Writing tests with JUnit

---

Annotate a method with `@Test` to flag it as a JUnit test

```
import org.junit.*;
import static org.junit.Assert.*;

/** Unit tests for my Foo ADT implementation */
public class FooTests {
    @Test
    public void testBar() {
        ... /* use JUnit assertions in here */
    }
}
```

# Using JUnit assertions

---

- JUnit assertions establish success or failure of the test method
  - *Note: JUnit assertions are different from Java's **assert** statement*
- Use to check that an actual result matches the expected value
  - Example: `assertEquals(42, meaningOfLife());`
  - Example: `assertTrue(list.isEmpty());`
- A test method stops immediately after the first assertion failure
  - If no assertion fails, then the test method passes
  - Other test methods still run either way
- JUnit results show details of any test failures

# Common JUnit assertions

---

JUnit's documentation has a full list, but these are the most common assertions.

Assertion	Failure condition
<code>assertTrue(<i>test</i>)</code>	<code><i>test</i> == false</code>
<code>assertFalse(<i>test</i>)</code>	<code><i>test</i> == true</code>
<code>assertEquals(<i>expected</i>, <i>actual</i>)</code>	<code><i>expected</i> and <i>actual</i> are not equal</code>
<code>assertSame(<i>expected</i>, <i>actual</i>)</code>	<code><i>expected</i> != <i>actual</i></code>
<code>assertNotSame(<i>expected</i>, <i>actual</i>)</code>	<code><i>expected</i> == <i>actual</i></code>
<code>assertNotNull(<i>value</i>)</code>	<code><i>value</i> != null</code>
<code>assertNotNull(<i>value</i>)</code>	<code><i>value</i> == null</code>

Any JUnit assertion can also take a string to show in case of failure, e.g., `assertEquals("helpful message", expected, actual)`.

# Checking for a thrown exception

---

- Should test that your code throws exceptions as specified
- This kind of test method fails if its body does *not* throw an exception of the named class
  - May not need any JUnit assertions inside the test method unlike our previous guideline

```
@Test(expected=IndexOutOfBoundsException.class)
public void testGetEmptyList() {
    List<String> list = new ArrayList<String>();
    list.get(0);
}
```

# Test ordering, setup, clean-up

---

JUnit does not promise to run tests in any particular order.

However, JUnit can run helper methods for common setup/cleanup

- Run before/after *each* test method in the class:

```
@Before
```

```
public void m() { ... }
```

```
@After
```

```
public void m() { ... }
```

- Run once before/after running *all* test methods in the class:

```
@BeforeClass
```

```
public static void m() { ... }
```

```
@AfterClass
```

```
public static void m() { ... }
```