

# CSE 373: Asymptotic Analysis

---

Michael Lee

Wednesday, Jan 10, 2018

# Warmup

**Warmup:** construct a *mathematical function* modeling the worst-case runtime of this method. Your model should be written in terms of  $q$ , the provided input integer.

Assume each `println` takes some constant  $c$  time to run.

```
public void mystery(int q) {
    for (int i = 0; i < q; i++) {
        for (int j = 0; j < q * q; j++) {
            System.out.println("Hello");
        }

        for (int j = 0; j < 10; j++) {
            System.out.println("World");
        }
    }
}
```

## Warmup

**Warmup:** construct a *mathematical function* modeling the worst-case runtime of this method. Your model should be written in terms of  $q$ , the provided input integer.

Assume each `println` takes some constant  $c$  time to run.

```
public void mystery(int q) {  
    for (int i = 0; i < q; i++) {  
        for (int j = 0; j < q * q; j++) {  
            System.out.println("Hello");  
        }  
  
        for (int j = 0; j < 10; j++) {  
            System.out.println("World");  
        }  
    }  
}
```

Answer:  $T(q) = q(cq^2 + 10c) = cq^3 + 10cq$

Two step process:

1. **Model** what we care about as a mathematical function
2. **Analyze** that function using asymptotic analysis

Two step process:


1. **Model** what we care about as a mathematical function
2. **Analyze** that function using asymptotic analysis  
Specifically: have a way to **compare** two functions

Two step process:

1. **Model** what we care about as a mathematical function
2. **Analyze** that function using asymptotic analysis  
Specifically: have a way to **compare** two functions  
Even more specifically: define a “less than or equal to” operator for functions


## Analysis: comparing functions

Question: Should we treat these two functions the same?



## Analysis: comparing functions


What about now?





## Analysis: comparing functions


What about now?



Intuition: our quadratic function is **dominating** the linear ones

## Analysis: comparing functions

What about now?




Intuition: our quadratic function is **dominating** the linear ones

Intuition: our linear functions (eventually) look the same


## Analysis: comparing functions

Let's zoom in...



## Analysis: comparing functions

Let's zoom in...



Intuition: quadratic function **eventually** dominates the linear ones

## Analysis: comparing functions

Our goal:

- ▶ We want a way to say  $n^2$  **eventually dominates**  $n$

# Analysis: comparing functions

Our goal:

- ▶ We want a way to say  $n^2$  **eventually dominates**  $n$
- ▶ We want a way to treat  $n$  and  $4n$  the same way

Intuition:

- ▶ Model made simplifying assumptions about constant factors
- ▶ Can usually improve constant-factor differences by being clever

# Analysis: comparing functions

Our goal:

- ▶ We want a way to say  $n^2$  **eventually dominates**  $n$
- ▶ We want a way to treat  $n$  and  $4n$  the same way

Intuition:

- ▶ Model made simplifying assumptions about constant factors
- ▶ Can usually improve constant-factor differences by being clever
- ▶ We want a way to do this rigorously!

## Function comparison: exercise

True or false?

- ▶ Is  $n$  “less then or equal to”  $5n + 3$ ?
- ▶ Is  $5n + 3$  “less then or equal to”  $n$ ?
- ▶ Is  $5n + 3$  “less then or equal to”  $1$ ?
- ▶ Is  $5n + 3$  “less then or equal to”  $n^2$ ?
- ▶ Is  $n^2 + 3n + 2$  “less then or equal to”  $n^3$ ?
- ▶ Is  $n^3$  “less then or equal to”  $n^2 + 3n + 2$ ?



## Function comparison: exercise

True or false?

- ▶ Is  $n$  “less then or equal to”  $5n + 3$ ? True
- ▶ Is  $5n + 3$  “less then or equal to”  $n$ ? True
- ▶ Is  $5n + 3$  “less then or equal to”  $1$ ? False
- ▶ Is  $5n + 3$  “less then or equal to”  $n^2$ ? True
- ▶ Is  $n^2 + 3n + 2$  “less then or equal to”  $n^3$ ? True
- ▶ Is  $n^3$  “less then or equal to”  $n^2 + 3n + 2$ ? False

# Analysis: comparing functions

Our goal:

- ▶ We want a way to say  $n^2$  **eventually dominates**  $n$
- ▶ We want a way to treat  $n$  and  $4n$  the same way

## Let's formalize this...

### Idea 1

A function  $f(n)$  is “less than or equal to”  $g(n)$  when  $f(n) \leq g(n)$  is true for all values of  $n \geq 0$ .


Does this work?

## Let's formalize this...

### Idea 1

A function  $f(n)$  is “less than or equal to”  $g(n)$  when  $f(n) \leq g(n)$  is true for all values of  $n \geq 0$ .

Does this work? Remember this?




## Let's formalize this...

### Idea 1

A function  $f(n)$  is "less than or equal to"  $g(n)$  when  $f(n) \leq g(n)$  is true for all values of  $n \geq 0$ .

Does this work? Remember this?



**Problem:** incorrectly handles the quadratic function!

## Let's formalize this...

### Idea 2

A function  $f(n)$  is “less than or equal to”  $g(n)$  when  $f(n) \leq g(n)$  is true for all values of  $n \geq n_0$ .

...where  $n_0 > 0$  is some constant value.

Does it work now?

## Let's formalize this...

### Idea 2

A function  $f(n)$  is “less than or equal to”  $g(n)$  when  $f(n) \leq g(n)$  is true for all values of  $n \geq n_0$ .

...where  $n_0 > 0$  is some constant value.

Does it work now?

We previously said we want to treat  $n$  and  $4n$  as being the “same”.

Do we?

## Let's formalize this...

### Idea 2

A function  $f(n)$  is “less than or equal to”  $g(n)$  when  $f(n) \leq g(n)$  is true for all values of  $n \geq n_0$ .

...where  $n_0 > 0$  is some constant value.

Does it work now?

We previously said we want to treat  $n$  and  $4n$  as being the “same”.

Do we?

**Problem:** No, we don't!



## Let's formalize this...

### Idea 3

A function  $f(n)$  is “less than or equal to”  $g(n)$  when  $f(n) \leq c \cdot g(n)$  is true for all values of  $n \geq n_0$ .

...where  $n_0 > 0$  is some constant value.

...where  $c > 0$  is some constant value.

Does it work now?

## Let's formalize this...

### Idea 3

A function  $f(n)$  is “less than or equal to”  $g(n)$  when  $f(n) \leq c \cdot g(n)$  is true for all values of  $n \geq n_0$ .

...where  $n_0 > 0$  is some constant value.

...where  $c > 0$  is some constant value.

Does it work now?

Yes!

## Definition: Dominated by

### Definition: Dominated by

A function  $f(n)$  is **dominated by**  $g(n)$  when...

## Definition: Dominated by

### Definition: Dominated by

A function  $f(n)$  is **dominated by**  $g(n)$  when...

- ▶ There exists two constants  $c > 0$  and  $n_0 > 0$ ...

## Definition: Dominated by

### Definition: Dominated by

A function  $f(n)$  is **dominated by**  $g(n)$  when...

- ▶ There exists two constants  $c > 0$  and  $n_0 > 0$ ...
- ▶ Such that for all values of  $n \geq n_0$ ...

## Definition: Dominated by

### Definition: Dominated by

A function  $f(n)$  is **dominated by**  $g(n)$  when...

- ▶ There exists two constants  $c > 0$  and  $n_0 > 0$ ...
- ▶ Such that for all values of  $n \geq n_0$ ...
- ▶  $f(n) \leq c \cdot g(n)$  is true

## Definition: Dominated by

### Definition: Dominated by

A function  $f(n)$  is **dominated by**  $g(n)$  when...

- ▶ There exists two constants  $c > 0$  and  $n_0 > 0$ ...
- ▶ Such that for all values of  $n \geq n_0$ ...
- ▶  $f(n) \leq c \cdot g(n)$  is true

The formal definition (not necessary to know this):

### Formal definition: Dominated by

A function  $f(n)$  is **dominated by**  $g(n)$  when

$$\exists(c > 0, n_0 > 0). \forall(n \geq n_0). (f(n) \leq cg(n))$$

...is true.

## Exercise

Demonstrate that  $5n^2 + 3n + 6$  is dominated by  $n^3$  by finding a  $c$  and  $n_0$  that satisfy the above definition.



## Exercise

Demonstrate that  $5n^2 + 3n + 6$  is dominated by  $n^3$  by finding a  $c$  and  $n_0$  that satisfy the above definition.

**Idea:** pick  $c = 10000$  and  $n_0 = 10000$ . (It probably works  $\neg \setminus (\setminus) \setminus / \setminus$ )

## Exercise

Demonstrate that  $5n^2 + 3n + 6$  is dominated by  $n^3$  by finding a  $c$  and  $n_0$  that satisfy the above definition.

**Idea:** pick  $c = 10000$  and  $n_0 = 10000$ . (It probably works  $\neg \setminus \_ ( \_ ) \_ / \_$ )

**Better idea:** show that  $5n^2 + 3n + 6$  is dominated by an easier function to analyze. E.g. note that:

$$\begin{aligned} 5n^2 + 3n + 6 &\leq 5n^2 + 3n^2 + 6n^2 && \text{for all } n \geq 1 \\ &= 14n^2 \\ &\leq 14n^3 \end{aligned}$$

## Exercise

Demonstrate that  $5n^2 + 3n + 6$  is dominated by  $n^3$  by finding a  $c$  and  $n_0$  that satisfy the above definition.

**Idea:** pick  $c = 10000$  and  $n_0 = 10000$ . (It probably works  $\setminus\_(\u201c)\_/$ )

**Better idea:** show that  $5n^2 + 3n + 6$  is dominated by an easier function to analyze. E.g. note that:

$$\begin{aligned}5n^2 + 3n + 6 &\leq 5n^2 + 3n^2 + 6n^2 && \text{for all } n \geq 1 \\ &= 14n^2 \\ &\leq 14n^3\end{aligned}$$

So, what value of  $c$  makes  $14n^3 \leq cn^3$  true (when  $n \geq 1$ )?

One possible choice:  $n_0 = 1$  and  $c = 14$ .

## Exercise

Demonstrate that  $5n^2 + 3n + 6$  is dominated by  $n^3$  by finding a  $c$  and  $n_0$  that satisfy the above definition.

**Idea:** pick  $c = 10000$  and  $n_0 = 10000$ . (It probably works  $\neg \setminus (\cup) \_ / \neg$ )

**Better idea:** show that  $5n^2 + 3n + 6$  is dominated by an easier function to analyze. E.g. note that:

$$\begin{aligned}5n^2 + 3n + 6 &\leq 5n^2 + 3n^2 + 6n^2 && \text{for all } n \geq 1 \\ &= 14n^2 \\ &\leq 14n^3\end{aligned}$$

So, what value of  $c$  makes  $14n^3 \leq cn^3$  true (when  $n \geq 1$ )?

One possible choice:  $n_0 = 1$  and  $c = 14$ .

So, since we know  $5n^2 + 3n + 6 \leq 14n^3$  for  $n \geq n_0$  and also know  $14n^3 \leq cn^3$ , we conclude  $5n^2 + 3n + 6 \leq cn^3$ .

## Exercise

Demonstrate that  $2n^3 - 3 + 9n^2 + \sqrt{n}$  is dominated by  $n^3$  (again by finding a  $c$  and  $n_0$ ).

## Exercise

Demonstrate that  $2n^3 - 3 + 9n^2 + \sqrt{n}$  is dominated by  $n^3$  (again by finding a  $c$  and  $n_0$ ).

Do the same thing. Note that:

$$\begin{aligned} 2n^3 - 3 + 9n^2 + \sqrt{n} &\leq 2n^3 + 9n^2 + n && \text{for all } n \geq 1 \\ &\leq 2n^3 + 9n^3 + n^3 \\ &= 12n^3 \end{aligned}$$

So, one possible choice of  $n_0$  and  $c$  is  $n_0 = 1$  and  $c = 12$ .

## Families of functions

Observation:

- ▶  $n$ ,  $5n + 3$ ,  $100n$ , etc... all dominate each other
- ▶ These three functions are the “same”

## Families of functions

Observation:

- ▶  $n$ ,  $5n + 3$ ,  $100n$ , etc... all dominate each other
- ▶ These three functions are the “same”

Idea: can we give a name to this “family” of functions?



# Families of functions

Observation:

- ▶  $n$ ,  $5n + 3$ ,  $100n$ , etc... all dominate each other
- ▶ These three functions are the “same”

Idea: can we give a name to this “family” of functions?

**Definition: Big- $\mathcal{O}$**

$\mathcal{O}(f(n))$  is the “family” or “set” of **all** functions that are **dominated by**  $f(n)$

# Families of functions

Observation:

- ▶  $n$ ,  $5n + 3$ ,  $100n$ , etc... all dominate each other
- ▶ These three functions are the “same”

Idea: can we give a name to this “family” of functions?

**Definition: Big- $\mathcal{O}$**

$\mathcal{O}(f(n))$  is the “family” or “set” of **all** functions that are **dominated by**  $f(n)$

Question: are  $\mathcal{O}(n)$ ,  $\mathcal{O}(5n + 3)$ , and  $\mathcal{O}(100n)$  all the same thing?

# Families of functions

Observation:

- ▶  $n$ ,  $5n + 3$ ,  $100n$ , etc... all dominate each other
- ▶ These three functions are the “same”

Idea: can we give a name to this “family” of functions?

**Definition: Big- $\mathcal{O}$**

$\mathcal{O}(f(n))$  is the “family” or “set” of **all** functions that are **dominated by**  $f(n)$

Question: are  $\mathcal{O}(n)$ ,  $\mathcal{O}(5n + 3)$ , and  $\mathcal{O}(100n)$  all the same thing?

Yes! By convention, we pick the “simplest” way of writing this and refer to this “family” as  $\mathcal{O}(n)$ .

A question: Do the following two sentences mean the same thing?

- ▶  $f(n)$  is dominated by  $g(n)$
- ▶  $f(n)$  is contained inside  $\mathcal{O}(g(n))$

A question: Do the following two sentences mean the same thing?

- ▶  $f(n)$  is dominated by  $g(n)$
- ▶  $f(n)$  is contained inside  $\mathcal{O}(g(n))$

Yes!

We can write this more concisely as  $f(n) \in \mathcal{O}(g(n))$ .

A question: Do the following two sentences mean the same thing?

- ▶  $f(n)$  is dominated by  $g(n)$
- ▶  $f(n)$  is contained inside  $\mathcal{O}(g(n))$

Yes!

We can write this more concisely as  $f(n) \in \mathcal{O}(g(n))$ .

An aside: some people write this as  $f(n) = \mathcal{O}(g(n))$

This is **wrong** (but common, so we reluctantly accept this)

## A few more questions

True or false:

- ▶  $5n + 3 \in \mathcal{O}(n)$
- ▶  $n \in \mathcal{O}(5n + 3)$
- ▶  $5n + 3 = \mathcal{O}(n)$
- ▶  $\mathcal{O}(5n + 3) = \mathcal{O}(n)$
- ▶  $\mathcal{O}(n^2) = \mathcal{O}(n)$
- ▶  $n^2 \in \mathcal{O}(1)$
- ▶  $n^2 \in \mathcal{O}(n)$
- ▶  $n^2 \in \mathcal{O}(n^2)$
- ▶  $n^2 \in \mathcal{O}(n^3)$
- ▶  $n^2 \in \mathcal{O}(n^{100})$

## A few more questions

True or false:

- ▶  $5n + 3 \in \mathcal{O}(n)$  True
- ▶  $n \in \mathcal{O}(5n + 3)$  True
- ▶  $5n + 3 = \mathcal{O}(n)$  True (by convention)
- ▶  $\mathcal{O}(5n + 3) = \mathcal{O}(n)$  True
- ▶  $\mathcal{O}(n^2) = \mathcal{O}(n)$  False
- ▶  $n^2 \in \mathcal{O}(1)$  False
- ▶  $n^2 \in \mathcal{O}(n)$  False
- ▶  $n^2 \in \mathcal{O}(n^2)$  True
- ▶  $n^2 \in \mathcal{O}(n^3)$  True
- ▶  $n^2 \in \mathcal{O}(n^{100})$  True



## Definitions: Dominates

$f(n) \in \mathcal{O}(g(n))$  is like saying “ $f(n)$  is less than or equal to  $g(n)$ ”.

Is there a way to say “greater than or equal to”?

## Definitions: Dominates

$f(n) \in \mathcal{O}(g(n))$  is like saying “ $f(n)$  is less than or equal to  $g(n)$ ”.

Is there a way to say “greater than or equal to”? Yes!

### Definition: Dominates

We say  $f(n)$  **dominates**  $g(n)$  when:

- ▶ There exists two constants  $c > 0$  and  $n_0 > 0$ ...
- ▶ Such that for all values of  $n \geq n_0$ ...
- ▶  $f(n) \geq c \cdot g(n)$  is true

## Definitions: Dominates

$f(n) \in \mathcal{O}(g(n))$  is like saying “ $f(n)$  is less than or equal to  $g(n)$ ”.

Is there a way to say “greater than or equal to”? Yes!

### Definition: Dominates

We say  $f(n)$  **dominates**  $g(n)$  when:

- ▶ There exists two constants  $c > 0$  and  $n_0 > 0$ ...
- ▶ Such that for all values of  $n \geq n_0$ ...
- ▶  $f(n) \geq c \cdot g(n)$  is true

### Definition: Big- $\Omega$

$\Omega(f(n))$  is the family of all functions that **dominates**  $f(n)$ .

## A few more questions...

True or false?

▶  $4n^2 \in \Omega(1)$

▶  $4n^2 \in \Omega(n)$

▶  $4n^2 \in \Omega(n^2)$

▶  $4n^2 \in \Omega(n^3)$

▶  $4n^2 \in \Omega(n^4)$

▶  $4n^2 \in \mathcal{O}(1)$

▶  $4n^2 \in \mathcal{O}(n)$

▶  $4n^2 \in \mathcal{O}(n^2)$

▶  $4n^2 \in \mathcal{O}(n^3)$

▶  $4n^2 \in \mathcal{O}(n^4)$

## A few more questions...

True or false?

▶  $4n^2 \in \Omega(1)$  True

▶  $4n^2 \in \Omega(n)$  True

▶  $4n^2 \in \Omega(n^2)$  True

▶  $4n^2 \in \Omega(n^3)$  False

▶  $4n^2 \in \Omega(n^4)$  False

▶  $4n^2 \in \mathcal{O}(1)$  False

▶  $4n^2 \in \mathcal{O}(n)$  False

▶  $4n^2 \in \mathcal{O}(n^2)$  True

▶  $4n^2 \in \mathcal{O}(n^3)$  True

▶  $4n^2 \in \mathcal{O}(n^4)$  True

## Definition: Big- $\Theta$

### Definition: Big- $\Theta$

We say  $f(n) \in \Theta(g(n))$  when both:

- ▶  $f(n) \in \mathcal{O}(g(n))$  and...
- ▶  $f(n) \in \Omega(g(n))$

...are true.

## Definition: Big- $\Theta$

### Definition: Big- $\Theta$

We say  $f(n) \in \Theta(g(n))$  when both:

- ▶  $f(n) \in \mathcal{O}(g(n))$  and...
- ▶  $f(n) \in \Omega(g(n))$

...are true.

Note: in industry, it's common for many people to ask for the big- $\mathcal{O}$  when they really want the big- $\Theta$ !

Important things to know:

- ▶ Intuition behind the definitions of “dominated by” and big- $\mathcal{O}$



Important things to know:

- ▶ Intuition behind the definitions of “dominated by” and  $\text{big-}\mathcal{O}$
- ▶ The precise definitions of:
  - ▶ “Dominated by” and  $\text{big-}\mathcal{O}$
  - ▶ “Dominates” and  $\text{big-}\Omega$
  - ▶  $\text{Big-}\Theta$

Important things to know:

- ▶ Intuition behind the definitions of “dominated by” and  $\text{big-}\mathcal{O}$
- ▶ The precise definitions of:
  - ▶ “Dominated by” and  $\text{big-}\mathcal{O}$
  - ▶ “Dominates” and  $\text{big-}\Omega$
  - ▶  $\text{Big-}\Theta$
- ▶ How to demonstrate that one function is dominated by another by finding  $c$  and  $n_0$  and applying the correct definition