

CSE 390Z: Mathematics for Computation Workshop

Week 1 Workshop Problems

Conceptual Review

- (a) What is the difference between an atomic and compound proposition?
- (b) What does it mean for two propositions to be *equivalent* ($p \equiv q$)?
- (c) What does it mean for two propositions to be *biconditional* ($p \leftrightarrow q$)?
- (d) What are two different methods to show that two propositions are equivalent?

1. Translation: Running from my problems

Define a set of three atomic propositions, and use them to translate the following sentences.

- (i) I am going for a run and it is snowing, or it is not snowing.
- (ii) If it's snowing and it's Friday, I am not going for a run.
- (iii) I am going for a run only if it is not Friday.

2. Translation: Age is just a number

Define a set of two atomic propositions, and use them to translate the following sentences.

- (i) If Kai is older than thirty, then Kai is older than twenty.
- (ii) Kai is older than thirty only if Kai is older than twenty.
- (iii) Whenever Kai is older than thirty, Kai is older than twenty.
- (iv) Kai being older than twenty is necessary for Kai to be older than thirty.

3. Truth Table

Draw a truth table for $(p \rightarrow \neg q) \rightarrow (r \oplus q)$

4. Proving Equivalences

We wish to prove that the following is true for all values of the propositions p and q (i.e. that it is a tautology):

$$\neg p \vee ((q \wedge p) \vee (\neg q \wedge p))$$

The following chain of equivalences does this, but it is missing citations for which rules are used. Fill in the blanks with names of the logic equivalences used at each step.

Hint: Reference the Logical Equivalences sheet under the Resources tab on the CSE 311 Course Website.

$\neg p \vee ((q \wedge p) \vee (\neg q \wedge p))$	$\equiv \neg p \vee ((p \wedge q) \vee (\neg q \wedge p))$	_____
	$\equiv \neg p \vee ((p \wedge q) \vee (p \wedge \neg q))$	_____
	$\equiv \neg p \vee (p \wedge (q \vee \neg q))$	_____
	$\equiv \neg p \vee (p \wedge T)$	_____
	$\equiv \neg p \vee p$	_____
	$\equiv p \vee \neg p$	_____
	$\equiv T$	_____

5. To Be or Not to Be Equivalent

For each of the following pairs of propositions, determine if the two propositions are equivalent. If they are, prove it using a chain of logical equivalences. If they are not, find an assignment of p and q on which their truth value differ.

(a) $p \wedge q$ vs. $p \vee q$

(b) $p \rightarrow q$ vs. $\neg q \rightarrow \neg p$

Note: You may not use the Law of Contrapositive in your justification!

(c) $p \rightarrow q$ vs. $\neg(p \wedge \neg q)$

6. Conditional Logic

You've already seen how implications translate to English and back, but we can also translate them to code. In English, $p \rightarrow q$ translates to "If p , then q ". We might also translate this (in Java) to a conditional:

```
if (p) {  
    q  
}
```

Lets say we have propositions p , q , and r . It may be helpful in the following problem to think of p and q as booleans, and r as a print statement.

(a) Convert each of the following propositions to Java (or pseudo-) code:

$$p \rightarrow (q \rightarrow r)$$

$$(p \wedge q) \rightarrow r$$

(b) Based on this, do you think $p \rightarrow (q \rightarrow r)$ and $(p \wedge q) \rightarrow r$ are logically equivalent? Why or why not?

7. Propositions in the wild

Give a real-life example of statements p , q , and r such that p and q together imply r , but neither p nor q alone imply r .