

# CSE 390Z: Mathematics for Computation Workshop

---

## Week 6 Workshop

### 0. Induction: Warm-Up

Prove by induction that  $5 \mid (6^n - 1)$  for all  $n \in \mathbb{N}$ .

### 1. Induction: Equality

Prove by induction that for every  $n \in \mathbb{N}$ , the following equality is true:

$$0 \cdot 2^0 + 1 \cdot 2^1 + 2 \cdot 2^2 + \cdots + n \cdot 2^n = (n - 1)2^{n+1} + 2.$$

## 2. Induction: Inequality

Prove by induction on  $n$  that for all integers  $n \geq 0$  the inequality  $(3 + \pi)^n \geq 3^n + n\pi 3^{n-1}$  is true.

## 3. Inductively Odd

An 123 student learning recursion wrote a recursive Java method to determine if a number is odd or not, and needs your help proving that it is correct.

```
public static boolean oddr(int n) {  
    if (n == 0)  
        return False;  
    else  
        return !oddr(n-1);  
}
```

Help the student by writing an inductive proof to prove that for all integers  $n \geq 0$ , the method `oddr` returns `True` if  $n$  is an odd number, and `False` if  $n$  is not an odd number (i.e.  $n$  is even). You may recall the definitions  $\text{Odd}(n) := \exists x \in \mathbb{Z}(n = 2x + 1)$  and  $\text{Even}(n) := \exists x \in \mathbb{Z}(n = 2x)$ ; `!True = False` and `!False = True`.

#### 4. Strong Induction: Stamp Collection

A store sells 3 cent and 5 cent stamps. Use strong induction to prove that you can make exactly  $n$  cents worth of stamps for all  $n \geq 10$ .

**Hint:** you'll need multiple base cases for this - think about how many steps back you need to go for your inductive step.

#### 5. Strong Induction: Recursively Defined Functions

Consider the function  $f(n)$  defined for integers  $n \geq 1$  as follows:

$$f(1) = 1 \text{ for } n = 1$$

$$f(2) = 4 \text{ for } n = 2$$

$$f(3) = 9 \text{ for } n = 3$$

$$f(n) = f(n-1) - f(n-2) + f(n-3) + 2(2n-3) \text{ for } n \geq 4$$

Prove by strong induction that for all  $n \geq 1$ ,  $f(n) = n^2$ .

**Complete the induction proof below.**

## 6. Strong Induction: A Variation of the Stamp Problem

A store sells candy in packs of 4 and packs of 7. Let  $P(n)$  be defined as "You are able to buy  $n$  packs of candy". For example,  $P(3)$  is not true, because you cannot buy exactly 3 packs of candy from the store. However, it turns out that  $P(n)$  is true for any  $n \geq 18$ . Use strong induction on  $n$  to prove this.

**Hint:** you'll need multiple base cases for this - think about how many steps back you need to go for your inductive step.