


CSE 417

Algorithms and Complexity

Autumn 2020
Lecture 29
NP-Completeness



Announcements


- Homework 9, Deadline, Sunday December 13
- Exam practice problems on course homepage
- Final Exam: Monday, December 14
 - 24 hour take home exam
 - Target: 2 to 4 hours of work time
- Possibly some extra office hours
- Approximate grade weighting
 - 75% HW, 25% Final

Today

REDUCIBILITY AMONG COMBINATORIAL PROBLEMS[†]

Richard M. Karp
University of California at Berkeley

There are a whole bunch of other important problems which are NP-Complete



ABSTRACT: A large class of computational problems involve the determination of properties of graphs, digraphs, integers, arrays of integers, finite families of finite sets, boolean formulas and elements of other countable domains. Through simple encodings from such domains into the set of words over a finite alphabet these problems can be converted into language recognition problems, and so can inquire into their computational complexity. It is reasonable to consider such a problem satisfactorily solved when an algorithm for its solution is found which terminates within a number of steps bounded by a polynomial in the length of the input. We show that a large number of classic combinatorial problems of covering, matching, packing, sorting, assignment and sequencing are equivalent, in the sense that either each of them possesses a polynomially-bounded algorithm or none of them does.

1. INTRODUCTION

All the general methods presently known for computing the chromatic number of a graph, deciding whether a graph has a Hamiltonian circuit, or solving a system of linear inequalities in which the variables are constrained to be 0 or 1, require a combinatorial search for which the worst case time requirements grow exponentially with the length of the input. In this paper we give theorems which strongly suggest, but do not imply, that these problems, as well as many others, will remain intractable permanently.

This research was partially supported by National Science Foundation Grant GJ-434.

Reducibility Among Combinatorial Problems





FIGURE 1 - Complete Problems

Populating the NP-Completeness Universe



- Circuit Sat $<_p$ 3-SAT
- 3-SAT $<_p$ Independent Set
- 3-SAT $<_p$ Vertex Cover
- Independent Set $<_p$ Clique
- 3-SAT $<_p$ Hamiltonian Circuit
- Hamiltonian Circuit $<_p$ Traveling Salesman
- 3-SAT $<_p$ Integer Linear Programming
- 3-SAT $<_p$ Graph Coloring
- 3-SAT $<_p$ Subset Sum
- Subset Sum $<_p$ Scheduling with Release times and deadlines

NP-Completeness Proofs

- Prove that problem X is NP-Complete
- Show that X is in NP (usually easy)
- Pick a known NP complete problem Y
- Show $Y <_p X$
- Types of NP completeness proofs
 - Easy modifications from other NP complete problems
 - Complicated gadget constructions from 3-SAT

Satisfiability

Literal: A Boolean variable or its negation.

$$x_i \text{ or } \bar{x}_i$$

Clause: A disjunction of literals.

$$C_j = x_1 \vee \bar{x}_2 \vee x_3$$

Conjunctive normal form: A propositional formula Φ that is the conjunction of clauses.

$$\Phi = C_1 \wedge C_2 \wedge C_3 \wedge C_4$$

SAT: Given CNF formula Φ , does it have a satisfying truth assignment?

3-SAT: SAT where each clause contains exactly 3 literals.

Ex: $(\bar{x}_1 \vee x_2 \vee x_3) \wedge (x_1 \vee \bar{x}_2 \vee x_3) \wedge (x_2 \vee x_3) \wedge (\bar{x}_1 \vee \bar{x}_2 \vee \bar{x}_3)$


Yes: $x_1 = \text{true}, x_2 = \text{true}, x_3 = \text{false}$.

3-SAT is NP-Complete

Theorem. 3-SAT is NP-complete.

Pf. Suffices to show that CIRCUIT-SAT \leq_p 3-SAT since 3-SAT is in NP.


- Let K be any circuit.
- Create a 3-SAT variable x_i for each circuit element i .
- Make circuit compute correct values at each node:
 - $x_5 = \neg x_3 \Rightarrow$ add 2 clauses: $x_2 \vee x_3, \bar{x}_2 \vee \bar{x}_3$
 - $x_1 = x_4 \vee x_6 \Rightarrow$ add 3 clauses: $x_1 \vee \bar{x}_4, x_1 \vee \bar{x}_5, x_1 \vee x_4 \vee x_5$
 - $x_0 = x_1 \wedge x_2 \Rightarrow$ add 3 clauses: $\bar{x}_0 \vee x_1, \bar{x}_0 \vee x_2, x_0 \vee \bar{x}_1 \vee \bar{x}_2$
- Hard-coded input values and output value.
 - $x_5 = 0 \Rightarrow$ add 1 clause: \bar{x}_5
 - $x_6 = 1 \Rightarrow$ add 1 clause: x_6
- Final step: turn clauses of length < 3 into clauses of length exactly 3. •



Independent Set

Independent Set

- Graph $G = (V, E)$, a subset S of the vertices is independent if there are no edges between vertices in S




3 Satisfiability Reduces to Independent Set

Claim. 3-SAT \leq_p INDEPENDENT-SET.

Pf. Given an instance Φ of 3-SAT, we construct an instance (G, k) of INDEPENDENT-SET that has an independent set of size k iff Φ is satisfiable.

Construction.

- G contains 3 vertices for each clause, one for each literal.
- Connect 3 literals in a clause in a triangle.
- Connect literal to each of its negations.




3 Satisfiability Reduces to Independent Set

Claim. G contains independent set of size $k = |\Phi|$ iff Φ is satisfiable.

Pf. \Rightarrow Let S be independent set of size k .

- S must contain exactly one vertex in each triangle.
- Set these literals to true. \leftarrow and any other variables in a consistent way
- Truth assignment is consistent and all clauses are satisfied.


Pf. \Leftarrow Given satisfying assignment, select one true literal from each triangle. This is an independent set of size k . •



Vertex Cover

Vertex Cover

- Graph $G = (V, E)$, a subset S of the vertices is a vertex cover if every edge in E has at least one endpoint in S




IS \leq_p VC


- Lemma: A set S is independent iff V-S is a vertex cover
- To reduce IS to VC, we show that we can determine if a graph has an independent set of size K by testing for a Vertex cover of size n - K

IS \leq_p VC

Find a maximum independent set S




Show that V-S is a vertex cover





Clique

- Clique
 - Graph $G = (V, E)$, a subset S of the vertices is a clique if there is an edge between every pair of vertices in S



Complement of a Graph

- Defn: $G'=(V,E')$ is the complement of $G=(V,E)$ if (u,v) is in E' iff (u,v) is not in E




IS \leq_p Clique

- Lemma: S is Independent in G iff S is a Clique in the complement of G
- To reduce IS to Clique, we compute the complement of the graph. The complement has a clique of size K iff the original graph has an independent set of size K


Hamiltonian Circuit Problem

- Hamiltonian Circuit – a simple cycle including all the vertices of the graph



Thm: Hamiltonian Circuit is NP Complete


- Reduction from 3-SAT



Page 475 in text


Clause Gadget

$$\overline{x_1} \vee x_2 \vee x_3$$




Reduce Hamiltonian Circuit to Hamiltonian Path

G_2 has a Hamiltonian Path iff G_1 has a Hamiltonian Circuit




Traveling Salesman Problem

- Given a complete graph with edge weights, determine the shortest tour that includes all of the vertices (visit each vertex exactly once, and get back to the starting point)



Find the minimum cost tour


Thm: $HC <_p TSP$



Matching




Two dimensional matching




Three dimensional matching (3DM)

3-SAT \leq_p 3DM



Truth Setting Gadget

3-SAT \leq_p 3DM



Clause gadget for $(\bar{X} \text{ OR } Y \text{ OR } Z)$

Garbage Collection Gadget
(Many copies)

Graph Coloring

- NP-Complete
 - Graph K-coloring
 - Graph 3-coloring
- Polynomial
 - Graph 2-Coloring

