


Another divide and conquer


Maximum contiguous subarray sum

Given: an array of integers (positive and negative), find the indices that give the maximum contiguous subarray sum.

Find i, j that maximize $A[i] + A[i + 1] + \dots + A[j]$



Conquer



Subarrays that cross have to be handled

Do we have to check all pairs i, j ?

Activity

Write a recurrence to describe the running time of this code. What's the big-O?

```
DivConqExponentiation(int a, int b, int n)
  if(n==0) return 1
  if(n==1) return a%b
  int k = divConqExponentiation(a,b,n/2) /*int div*/
  if(n%2==1) return (k*k*a)%b
  return (k*k)%b
```

Go to pollev.com/robbie so I know how long to explain

Master Theorem

Given a recurrence of the following form, where a , b , c , and d are constants:

$$T(n) = \begin{cases} d & \text{if } n \text{ is at most some constant} \\ aT\left(\frac{n}{b}\right) + f(n) & \text{otherwise} \end{cases}$$

Where $f(n)$ is $\Theta(n^c)$

If $\log_b a < c$ then $T(n) \in \Theta(n^c)$

If $\log_b a = c$ then $T(n) \in \Theta(n^c \log n)$

If $\log_b a > c$ then $T(n) \in \Theta(n^{\log_b a})$