


CSE 421 Introduction to Algorithms Winter 2001

The Network Flow Problem

CSE 421,W '01, Ruzzo

1

The Network Flow Problem



How much stuff can flow from s to t?

CSE 421,W '01, Ruzzo

2

Net Flow: Formal Definition

Given:

A digraph $G = (V, E)$

Two vertices $s, t \in V$
(source & sink)

A capacity $c(u, v) \geq 0$
for each $(u, v) \in E$
(and $c(u, v) = 0$ for all non-edges (u, v))

Find:

A flow function $f: V \times V \rightarrow \mathbb{R}$ s.t.,
for all u, v :

- $f(u, v) \leq c(u, v)$ [Capacity Constraint]
- $f(u, v) = -f(v, u)$ [Skew Symmetry]
- if $u \neq s, t$, $\sum_v f(u, v) = 0$ [Flow Conservation]

Maximizing total flow $|f| = f(s, V)$


Notation:

$$f(X, Y) = \sum_{x \in X} \sum_{y \in Y} f(x, y)$$

CSE 421,W '01, Ruzzo

3

Example: A Flow Function



flow/capacity, not .66...

$$f(s, u) = f(u, t) = 2$$

$$f(u, s) = f(t, u) = -2 \text{ (Why?)}$$


$$f(s, t) = -f(t, s) = 0 \text{ (In every flow function for this G. Why?)}$$

$$f(u, V) = \sum_{v \in V} f(u, v) = f(u, s) + f(u, t) = -2 + 2 = 0$$

CSE 421,W '01, Ruzzo

4

Example: A Flow Function




- Not shown: $f(u, v)$ if $u, v \in V$ and $f(u, v) \leq 0$
- Note: max flow ≥ 4 since f is a flow function, with $|f| = 4$

CSE 421,W '01, Ruzzo

5

Max Flow via a Greedy Alg?

While there is an $s \rightarrow t$ path in G
Pick such a path, p
Find c_p , the min capacity of any edge in p
Subtract c_p from all capacities on p
Delete edges of capacity 0




CSE 421,W '01, Ruzzo

6

Max Flow via a Greedy Alg?

This does **NOT** always find a max flow:
If you pick $s \rightarrow b \rightarrow a \rightarrow t$ first,



Flow stuck at 2. But flow 3 possible.

CSE 421, W '01, Ruzzo

7

A Brief History of Flow


#	year	discoverer(s)	bound
1	1951	Dantzig	$O(n^2 m U)$
2	1955	Ford & Fulkerson	$O(n m U)$
3	1970	Dinitz	$O(n m^2)$
4	1970	Edmonds & Karp	$O(n^2 m)$
5	1972	Edmonds & Karp	$O(n^3 \log U)$
6	1972	Dinitz	$O(n m \log U)$
7	1974	Gabow	$O(n^3)$
8	1977	Cherkassky	$O(n^2 \sqrt{m})$
9	1980	Gall & Naamad	$O(n m \log^2 n)$
10	1980	Goldberg & Tarjan	$O(n m \log^2(m))$
11	1986	Goldberg & Tarjan	$O(n^2 m \log^2(m))$
12	1987	Aluja & Orlin	$O(n m + n^2 \log U)$
13	1987	Aluja et al.	$O(n m \log(n \log U / (m+2)))$
14	1989	Cheriyan & Hagerup	$O(n m + n^2 \log^2 n)$
15	1990	Cheriyan et al.	$O(n m + n^2 \log n)$
16	1990	Alex	$O(n m + n^{2/3} \log n)$
17	1992	King et al.	$O(n m + n^{2/3})$
18	1994	Phillips & Westbrook	$O(n(m \log_{m/n} n + \log^{2/3} n))$
19	1994	King et al.	$O(n m \log_{m/n} (n \log n))$
20	1997	Goldberg & Rao	$O(n^{2/3} m \log(n/m) \log U)$ $O(n^{2/3} m \log(n/m) \log U)$

$n = \#$ of vertices
 $m = \#$ of edges
 $U = \text{Max capacity}$

Source: Goldberg & Rao,
FOCS '97

8

Greed Revisited




CSE 421, W '01, Ruzzo

9

Residual Capacity

- The **residual capacity** (w.r.t. f) of (u,v) is $c_f(u,v) = c(u,v) - f(u,v)$
- E.g.:
 - $c_f(s,b) = 7$;
 - $c_f(a,x) = 1$;
 - $c_f(x,a) = 3$;
 - $c_f(x,t) = 0$ (a **saturated edge**)



CSE 421, W '01, Ruzzo

10

Residual Networks & Augmenting Paths

- The **residual network** (w.r.t. f) is the graph $G_f = (V, E_f)$, where


$$E_f = \{(u,v) \mid c_f(u,v) > 0\}$$

- An **augmenting path** (w.r.t. f) is a simple $s \rightarrow t$ path in G_f .

CSE 421, W '01, Ruzzo

11


A Residual Network



CSE 421, W '01, Ruzzo

12

An Augmenting Path



CSE 421, W '01, Ruzzo

13

Lemma 1


If f admits an augmenting path p , then f is not maximal.

Proof: "obvious" -- augment along p by c_p , the min residual capacity of p 's edges.

CSE 421, W '01, Ruzzo

14

Augmenting A Flow



CSE 421, W '01, Ruzzo


15

Proof of Lemma 1—Case 1


Let (u,v) be any edge in augmenting path. Note

$$c_r(u,v) = c(u,v) - f(u,v) \geq c_p > 0$$

Case 1: $f(u,v) \geq 0$:



Add forward flow



CSE 421, W '01, Ruzzo




16

Proof of Lemma 1—Case 2

Let (u,v) be any edge in augmenting path. Note
 $c_r(u,v) = c(u,v) - f(u,v) \geq c_p > 0$

Case 2: $f(u,v) \leq -c_p$:
 $f(v,u) = -f(u,v) \geq c_p$

Cancel/redirect reverse flow



CSE 421, W '01, Ruzzo




17

Proof of Lemma 1—Case 3

Let (u,v) be any edge in augmenting path. Note
 $c_r(u,v) = c(u,v) - f(u,v) \geq c_p > 0$

Case 3: $-c_p \leq f(u,v) < 0$:

???



CSE 421, W '01, Ruzzo



18

Proof of Lemma 1—Case 3

Let (u,v) be any edge in augmenting path. Note
 $c_f(u,v) = c(u,v) - f(u,v) \geq c_p > 0$

Case 3: $-c_p \leq f(u,v) < 0$
 $c_p \geq f(v,u) > 0$:

Both:
cancel/redirect
reverse flow
and
add forward flow



CSE 421, W '01, Ruzzo

19

Ford-Fulkerson Method

While G_f has an augmenting path,
augment

Questions:


- » Does it halt?
- » Does it find a maximum flow?
- » How fast?

CSE 421, W '01, Ruzzo

20

Cuts

- A partition S, T of V is a **cut** if $s \in S, t \in T$
- **Capacity** of cut S, T is $c(S, T) = \sum_{u \in S} \sum_{v \in T} c(u, v)$




CSE 421, W '01, Ruzzo

21

Lemma 2

- For any flow f and any cut S, T ,
 - » the net flow across the cut equals the total flow, i.e., $|f| = f(S, T)$, and
 - » the net flow across the cut cannot exceed the capacity of the cut, i.e. $f(S, T) \leq c(S, T)$
- Corollary:
 $\text{Max flow} \leq \text{Min cut}$



CSE 421, W '01, Ruzzo

22

Max Flow / Min Cut Theorem

For any flow f , the following are equivalent

- (1) $|f| = c(S, T)$ for some cut S, T (a min cut)
- (2) f is a maximum flow
- (3) f admits no augmenting path


Proof:

- (1) \Rightarrow (2): corollary to lemma 2
- (2) \Rightarrow (3): contrapositive of lemma 1

CSE 421, W '01, Ruzzo

23

(3) \Rightarrow (1)



$S = \{ u \mid \exists \text{ an augmenting path from } s \text{ to } u \}$

$T = V - S; s \in S, t \in T$

For any (u, v) in $S \times T$, \exists an augmenting path from s to u , but **not** to v .

$\therefore (u, v)$ has 0 residual capacity:

$$(u, v) \in E \Rightarrow \text{saturated} \quad f(u, v) = c(u, v) \\ (v, u) \in E \Rightarrow \text{no flow} \quad f(v, u) = 0 = -f(u, v)$$


This is true for every edge crossing the cut, i.e.

$$|f| = f(S, T) = \sum_{u \in S} \sum_{v \in T} f(u, v) = \sum_{u \in S, v \in T, (u, v) \in E} f(u, v) = \sum_{u \in S, v \in T, (u, v) \in E} c(u, v) = c(S, T)$$

24

Corollaries & Facts

- If Ford-Fulkerson terminates, then it's found a max flow.
- It will terminate if $c(e)$ integer or rational (but may not if they're irrational).
- However, may take exponential time, even with integer capacities:



$c = 10^{99}$, say

CSE 421, W '01, Ruzzo

25

Edmonds-Karp Algorithm

- Use a shortest augmenting path (via Breadth First Search in residual graph)
- Time: $O(n m^2)$


CSE 421, W '01, Ruzzo

26

BFS/Shortest Path Lemmas

Distance from s is never reduced by:

- Deleting an edge
proof: no new (hence no shorter) path created
- Adding an edge (u,v) , provided v is nearer than u
proof: BFS is unchanged, since v visited before (u,v) examined



CSE 421, W '01, Ruzzo

27

Lemma 3


Let f be a flow, G_f the residual graph, and p a shortest augmenting path. Then no vertex is closer to s after augmentation along p .

Proof: Augmentation only deletes edges, adds back edges

CSE 421, W '01, Ruzzo

28

Augmentation vs BFS



CSE 421, W '01, Ruzzo

29

Theorem 2

The Edmonds-Karp Algorithm performs $O(mn)$ flow augmentations


Proof:

$\{u,v\}$ is critical on augmenting path p if it's closest to s having min residual capacity. Won't be critical again until farther from s . So each edge critical at most n times.

CSE 421, W '01, Ruzzo

30

Augmentation vs BFS Level



Corollary

Edmonds-Karp runs in $O(nm^2)$


CSE 421, W '01, Ruzzo

32

Flow Integrity Theorem

If all capacities are integers


- » The max flow has an integer value
- » Ford-Fulkerson method finds a max flow in which $f(u,v)$ is an integer for all edges (u,v)



CSE 421, W '01, Ruzzo

33

Bipartite Maximum Matching



Bipartite Graphs:

- $G = (V, E)$
- $V = L \cup R$ ($L \cap R = \emptyset$)
- $E \subseteq L \times R$

Matching:


- A set of edges $M \subseteq E$ such that no two edges touch a common vertex

Problem:

- Find a matching M of maximum size

34


Reducing Matching to Flow



35

Reducing Matching to Flow

Theorem: Max matching size = max flow value



36

Notes on Matching

- Max Flow Algorithm is probably overly general here
- But most direct matching algorithms use "augmenting path" type ideas similar to that in max flow – See text & homework
- Time $mn^{1/2}$ possible via Edmonds-Karp