


**CSE 421  
Algorithms**


Richard Anderson  
Lecture 26  
NP-Completeness

## Complexity Classes

- P: Class of problems that can be solved in polynomial time
- NP: Class of problems that can be solved in non-deterministic polynomial time
  - Corresponds to problems where we can verify a solution in polynomial time



instance s



certificate t

## Polynomial time reductions

- Y is Polynomial Time Reducible to X
  - Solve problem Y with a polynomial number of computation steps and a polynomial number of calls to a black box that solves X
  - Notations:  $Y <_p X$

## Lemmas



- Suppose  $Y <_p X$ . If X can be solved in polynomial time, then Y can be solved in polynomial time.
- Suppose  $Y <_p X$ . If Y cannot be solved in polynomial time, then X cannot be solved in polynomial time.

## NP-Completeness


- A problem X is NP-complete if
  - X is in NP
  - For every Y in NP,  $Y <_p X$
- X is a “hardest” problem in NP
- If X is NP-Complete, Z is in NP and  $X <_p Z$ 
  - Then Z is NP-Complete

## Cook’s Theorem


- The Circuit Satisfiability Problem is NP-Complete



- ### History
- Jack Edmonds
    - Identified NP
  - Steve Cook
    - Cook's Theorem – NP-Completeness
  - Dick Karp
    - Identified "standard" collection of NP-Complete Problems
  - Leonid Levin
    - Independent discovery of NP-Completeness in USSR




- ### Populating the NP-Completeness Universe
- 
- Circuit Sat  $\leq_p$  3-SAT
  - 3-SAT  $\leq_p$  Independent Set
  - 3-SAT  $\leq_p$  Vertex Cover
  - Independent Set  $\leq_p$  Clique
  - 3-SAT  $\leq_p$  Hamiltonian Circuit
  - Hamiltonian Circuit  $\leq_p$  Traveling Salesman
  - 3-SAT  $\leq_p$  Integer Linear Programming
  - 3-SAT  $\leq_p$  Graph Coloring
  - 3-SAT  $\leq_p$  Subset Sum
  - Subset Sum  $\leq_p$  Scheduling with Release times and deadlines



## Vertex Cover

- Vertex Cover
  - Graph  $G = (V, E)$ , a subset  $S$  of the vertices is a vertex cover if every edge in  $E$  has at least one endpoint in  $S$




## Cook's Theorem

- The Circuit Satisfiability Problem is NP-Complete
- Circuit Satisfiability
  - Given a boolean circuit, determine if there is an assignment of boolean values to the input to make the output true

## Circuit SAT

Find a satisfying assignment




## Proof of Cook's Theorem

- Reduce an arbitrary problem  $Y$  in NP to  $X$
- Let  $A$  be a non-deterministic polynomial time algorithm for  $Y$
- Convert  $A$  to a circuit, so that  $Y$  is a Yes instance iff and only if the circuit is satisfiable

## Populating the NP-Completeness Universe

- Circuit Sat  $\leq_p$  3-SAT
- 3-SAT  $\leq_p$  Independent Set
- 3-SAT  $\leq_p$  Vertex Cover
- Independent Set  $\leq_p$  Clique
- 3-SAT  $\leq_p$  Hamiltonian Circuit
- Hamiltonian Circuit  $\leq_p$  Traveling Salesman
- 3-SAT  $\leq_p$  Integer Linear Programming
- 3-SAT  $\leq_p$  Graph Coloring
- 3-SAT  $\leq_p$  Subset Sum
- Subset Sum  $\leq_p$  Scheduling with Release times and deadlines



## Satisfiability

Literal: A Boolean variable or its negation.

$$x_i \text{ or } \bar{x}_i$$

Clause: A disjunction of literals.

$$C_j = x_1 \vee \bar{x}_2 \vee x_3$$

Conjunctive normal form: A propositional formula  $\Phi$  that is the conjunction of clauses.

$$\Phi = C_1 \wedge C_2 \wedge C_3 \wedge C_4$$

SAT: Given CNF formula  $\Phi$ , does it have a satisfying truth assignment?

3-SAT: SAT where each clause contains exactly 3 literals.


Ex:  $(\bar{x}_1 \vee x_2 \vee x_3) \wedge (x_1 \vee \bar{x}_2 \vee x_3) \wedge (x_2 \vee x_3) \wedge (\bar{x}_1 \vee \bar{x}_2 \vee \bar{x}_3)$   
 Yes:  $x_1 = \text{true}, x_2 = \text{true}, x_3 = \text{false}$ .

# 3-SAT is NP-Complete

Theorem. 3-SAT is NP-complete.

Pf. Suffices to show that CIRCUIT-SAT  $\leq_p$  3-SAT since 3-SAT is in NP.

- Let  $K$  be any circuit.
- Create a 3-SAT variable  $x_i$  for each circuit element  $i$ .
- Make circuit compute correct values at each node:
  - $x_2 = \neg x_3 \Rightarrow$  add 2 clauses:  $x_2 \vee x_3, \overline{x_2} \vee \overline{x_3}$
  - $x_1 = x_4 \vee x_5 \Rightarrow$  add 3 clauses:  $\overline{x_1} \vee x_4, \overline{x_1} \vee x_5, x_1 \vee \overline{x_4} \vee \overline{x_5}$
  - $x_0 = x_1 \wedge x_2 \Rightarrow$  add 3 clauses:  $\overline{x_0} \vee x_1, \overline{x_0} \vee x_2, x_0 \vee \overline{x_1} \vee \overline{x_2}$
- Hard-coded input values and output value.
  - $x_5 = 0 \Rightarrow$  add 1 clause:  $\overline{x_5}$
  - $x_0 = 1 \Rightarrow$  add 1 clause:  $x_0$
- Final step: turn clauses of length  $< 3$  into clauses of length exactly 3. •




# 3 Satisfiability Reduces to Independent Set

Claim. 3-SAT  $\leq_p$  INDEPENDENT-SET.

Pf. Given an instance  $\Phi$  of 3-SAT, we construct an instance  $(G, k)$  of INDEPENDENT-SET that has an independent set of size  $k$  iff  $\Phi$  is satisfiable.

Construction.

- $G$  contains 3 vertices for each clause, one for each literal.
- Connect 3 literals in a clause in a triangle.
- Connect literal to each of its negations.




# 3 Satisfiability Reduces to Independent Set

Claim.  $G$  contains independent set of size  $k = |\Phi|$  iff  $\Phi$  is satisfiable.

Pf.  $\Rightarrow$  Let  $S$  be independent set of size  $k$ .

- $S$  must contain exactly one vertex in each triangle.
- Set these literals to true.  $\leftarrow$  and any other variables in a consistent way
- Truth assignment is consistent and all clauses are satisfied.

Pf.  $\Leftarrow$  Given satisfying assignment, select one true literal from each triangle. This is an independent set of size  $k$ . •




# IS $\leq_p$ VC


- Lemma: A set  $S$  is independent iff  $V-S$  is a vertex cover
- To reduce IS to VC, we show that we can determine if a graph has an independent set of size  $K$  by testing for a Vertex cover of size  $n - K$

# IS $\leq_p$ VC

Find a maximum independent set  $S$




Show that  $V-S$  is a vertex cover



# Clique


• Clique

- Graph  $G = (V, E)$ , a subset  $S$  of the vertices is a clique if there is an edge between every pair of vertices in  $S$



## Complement of a Graph

- Defn:  $G'=(V,E')$  is the complement of  $G=(V,E)$  if  $(u,v)$  is in  $E'$  iff  $(u,v)$  is not in  $E$




## IS $<_p$ Clique

- Lemma: S is Independent in G iff S is a Clique in the complement of G
- To reduce IS to Clique, we compute the complement of the graph. The complement has a clique of size K iff the original graph has an independent set of size K

## Hamiltonian Circuit Problem

- Hamiltonian Circuit – a simple cycle including all the vertices of the graph




## Thm: Hamiltonian Circuit is NP Complete


- Reduction from 3-SAT

## Traveling Salesman Problem

- Given a complete graph with edge weights, determine the shortest tour that includes all of the vertices (visit each vertex exactly once, and get back to the starting point)



## Thm: HC $<_p$ TSP



## Graph Coloring

- NP-Complete
  - Graph K-coloring
  - Graph 3-coloring
- Polynomial
  - Graph 2-Coloring

