

CSE 421 Algorithms

Richard Anderson
Lecture 13
Recurrences, Part 2

Announcements


- Midterm
 - Monday, Oct 31, in class, closed book
 - Through section 5.2
 - Midterm review
- Homework 5 available

Recurrence Examples

- $T(n) = 2 T(n/2) + cn$
 - $O(n \log n)$
- $T(n) = T(n/2) + cn$
 - $O(n)$
- More useful facts:
 - $\log_k n = \log_2 n / \log_2 k$
 - $k^{\log n} = n^{\log k}$

$$\sum_{i=0}^n x^i = \frac{1 - x^{n+1}}{1 - x}$$

Unrolling the recurrence



Recursive Matrix Multiplication

Multiply 2 x 2 Matrices:
 $\begin{vmatrix} r & s \\ t & u \end{vmatrix} = \begin{vmatrix} a & b \\ c & d \end{vmatrix} \begin{vmatrix} e & g \\ f & h \end{vmatrix}$

$$\begin{aligned} r &= ae + bf \\ s &= ag + bh \\ t &= ce + df \\ u &= cg + dh \end{aligned}$$

A $N \times N$ matrix can be viewed as a 2×2 matrix with entries that are $(N/2) \times (N/2)$ matrices.

The recursive matrix multiplication algorithm recursively multiplies the $(N/2) \times (N/2)$ matrices and combines them using the equations for multiplying 2×2 matrices

Recursive Matrix Multiplication

- How many recursive calls are made at each level?
- How much work in combining the results?
- What is the recurrence?


What is the run time for the recursive Matrix Multiplication Algorithm?

- Recurrence:


$$T(n) = 4T(n/2) + n$$

Total Work


$$\sum_{k=0}^{\log n} 2^k n = (2n - 1)n$$



$$T(n) = 2T(n/2) + n^2$$



$$T(n) = 2T(n/2) + n^{1/2}$$



Recurrences

- Three basic behaviors
 - Dominated by initial case
 - Dominated by base case
 - All cases equal – we care about the depth

What you really need to know about recurrences

- Work per level changes geometrically with the level
- Geometrically increasing ($x > 1$)
 - The bottom level wins
- Geometrically decreasing ($x < 1$)
 - The top level wins
- Balanced ($x = 1$)
 - Equal contribution

Classify the following recurrences (Increasing, Decreasing, Balanced)

- $T(n) = n + 5T(n/8)$
- $T(n) = n + 9T(n/8)$
- $T(n) = n^2 + 4T(n/2)$
- $T(n) = n^3 + 7T(n/2)$
- $T(n) = n^{1/2} + 3T(n/4)$

Strassen's Algorithm

Multiply 2 x 2 Matrices:

$$\begin{bmatrix} r & s \\ t & u \end{bmatrix} = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} e & g \\ f & h \end{bmatrix}$$

Where:

$$p_1 = (b + d)(f + g)$$

$$p_2 = (c + d)e$$

$$p_3 = a(g - h)$$

$$p_4 = d(f - e)$$

$$p_5 = (a - b)h$$

$$p_6 = (c - d)(e + g)$$

$$p_7 = (b - d)(f + h)$$

$$r = p_1 + p_4 - p_5 + p_7$$

$$s = p_3 + p_5$$

$$t = p_2 + p_6$$

$$u = p_1 + p_3 - p_2 + p_7$$

Recurrence for Strassen's Algorithms

- $T(n) = 7 T(n/2) + cn^2$
- What is the runtime?

BFPRT Recurrence

- $T(n) \leq T(3n/4) + T(n/5) + 20n$

What bound do you expect?