


CSE 421
Algorithms
 Richard Anderson
 Lecture 25
 NP-Completeness

NP Completeness




Algorithms vs. Lower bounds

- Algorithmic Theory
 - What we can compute
 - I can solve problem X with resources R
 - Proofs are almost always to give an algorithm that meets the resource bounds
- Lower bounds
 - How do we show that something can't be done?

Theory of NP Completeness

The Universe



Polynomial Time

- P: Class of problems that can be solved in polynomial time
 - Corresponds with problems that can be solved efficiently in practice
 - Right class to work with “theoretically”

Decision Problems

- Theory developed in terms of yes/no problems
 - Independent set
 - Given a graph G and an integer K, does G have an independent set of size at least K
 - Network Flow
 - Given a graph G with edge capacities, a source vertex s, and sink vertex t, and an integer K, does the graph have flow function with value at least K

Definition of P

Decision problems for which there is a polynomial time algorithm

Problem	Description	Algorithm	Yes	No
MULTIPLE	Is x a multiple of y?	Grade school division	51, 17	51, 16
RELPRIME	Are x and y relatively prime?	Euclid's algorithm	34, 39	34, 51
PRIMES	Is x prime?	Agrawal, Kayal, Saxena (2002)	53	51
EDIT-DISTANCE	Is the edit distance between x and y less than 5?	Dynamic programming	niether neither	acgggt ttttta
LSOLVE	Is there a vector x that satisfies Ax = b?	Gaussian elimination	$\begin{bmatrix} 0 & 1 & 1 & 4 \\ 2 & 4 & -2 & 2 \\ 0 & 3 & 15 & 36 \end{bmatrix}$	$\begin{bmatrix} 1 & 0 & 0 & 2 \\ 1 & 1 & 1 & 1 \\ 0 & 1 & 1 & 1 \end{bmatrix}$

What is NP?

- Problems solvable in non-deterministic polynomial time . . .
- Problems where “yes” instances have polynomial time checkable certificates

Certificate examples

- Independent set of size K
 - The Independent Set
- Satisfiable formula
 - Truth assignment to the variables
- Hamiltonian Circuit Problem
 - A cycle including all of the vertices
- K-coloring a graph
 - Assignment of colors to the vertices

Certifiers and Certificates: 3-Satisfiability

SAT: Does a given CNF formula have a satisfying formula

Certificate: An assignment of truth values to the n boolean variables

Certifier: Check that each clause has at least one true literal,

instance s

$$(\bar{x}_1 \vee x_2 \vee x_3) \wedge (x_1 \vee \bar{x}_2 \vee x_3) \wedge (x_1 \vee x_2 \vee x_4) \wedge (\bar{x}_1 \vee \bar{x}_3 \vee \bar{x}_4)$$

certificate t


$$x_1 = 1, x_2 = 1, x_3 = 0, x_4 = 1$$

Certifiers and Certificates: Hamiltonian Cycle

HAM-CYCLE. Given an undirected graph $G = (V, E)$, does there exist a simple cycle C that visits every node?

Certificate. A permutation of the n nodes.

Certifier. Check that the permutation contains each node in V exactly once, and that there is an edge between each pair of adjacent nodes in the permutation.



Polynomial time reductions

- Y is Polynomial Time Reducible to X
 - Solve problem Y with a polynomial number of computation steps and a polynomial number of calls to a black box that solves X
 - Notations: $Y <_P X$

Lemmas

- Suppose $Y <_P X$. If X can be solved in polynomial time, then Y can be solved in polynomial time.
- Suppose $Y <_P X$. If Y cannot be solved in polynomial time, then X cannot be solved in polynomial time.

NP-Completeness


- A problem X is NP-complete if
 - X is in NP
 - For every Y in NP, $Y <_P X$
- X is a “hardest” problem in NP
- If X is NP-Complete, Z is in NP and $X <_P Z$
 - Then Z is NP-Complete

Cook’s Theorem


- The Circuit Satisfiability Problem is NP-Complete

Circuit SAT

Find a satisfying assignment




Garey and Johnson



History


- Jack Edmonds
 - Identified NP
- Steve Cook
 - Cook's Theorem – NP-Completeness
- Dick Karp
 - Identified "standard" collection of NP-Complete Problems
- Leonid Levin
 - Independent discovery of NP-Completeness in USSR

P vs. NP Question




Populating the NP-Completeness Universe

- Circuit Sat \leq_p 3-SAT
- 3-SAT \leq_p Independent Set
- 3-SAT \leq_p Vertex Cover
- Independent Set \leq_p Clique
- 3-SAT \leq_p Hamiltonian Circuit
- Hamiltonian Circuit \leq_p Traveling Salesman
- 3-SAT \leq_p Integer Linear Programming
- 3-SAT \leq_p Graph Coloring
- 3-SAT \leq_p Subset Sum
- Subset Sum \leq_p Scheduling with Release times and deadlines




Sample Problems

- Independent Set
 - Graph $G = (V, E)$, a subset S of the vertices is independent if there are no edges between vertices in S




Vertex Cover

- Vertex Cover
 - Graph $G = (V, E)$, a subset S of the vertices is a vertex cover if every edge in E has at least one endpoint in S



Cook's Theorem

- The Circuit Satisfiability Problem is NP-Complete
- Circuit Satisfiability
 - Given a boolean circuit, determine if there is an assignment of boolean values to the input to make the output true



Proof of Cook's Theorem

- Reduce an arbitrary problem Y in NP to X
- Let A be a non-deterministic polynomial time algorithm for Y
- Convert A to a circuit, so that Y is a Yes instance iff and only if the circuit is satisfiable

Populating the NP-Completeness Universe

- Circuit Sat \leq_p 3-SAT
- 3-SAT \leq_p Independent Set
- 3-SAT \leq_p Vertex Cover
- Independent Set \leq_p Clique
- 3-SAT \leq_p Hamiltonian Circuit
- Hamiltonian Circuit \leq_p Traveling Salesman
- 3-SAT \leq_p Integer Linear Programming
- 3-SAT \leq_p Graph Coloring
- 3-SAT \leq_p Subset Sum
- Subset Sum \leq_p Scheduling with Release times and deadlines

Satisfiability

Literal: A Boolean variable or its negation. x_i or \bar{x}_i

Clause: A disjunction of literals. $C_j = x_1 \vee \bar{x}_2 \vee x_3$

Conjunctive normal form: A propositional formula Φ that is the conjunction of clauses. $\Phi = C_1 \wedge C_2 \wedge C_3 \wedge C_4$

SAT: Given CNF formula Φ , does it have a satisfying truth assignment?

3-SAT: SAT where each clause contains exactly 3 literals.


Ex: $(\bar{x}_1 \vee \bar{x}_2 \vee x_3) \wedge (x_1 \vee \bar{x}_2 \vee x_3) \wedge (x_2 \vee x_3) \wedge (\bar{x}_1 \vee \bar{x}_2 \vee \bar{x}_3)$
 Yes: $x_1 = \text{true}, x_2 = \text{true}, x_3 = \text{false}$.

3-SAT is NP-Complete

Theorem. 3-SAT is NP-complete.

Pf. Suffices to show that CIRCUIT-SAT \leq_p 3-SAT since 3-SAT is in NP.

- Let K be any circuit.
- Create a 3-SAT variable x_i for each circuit element i .
- Make circuit compute correct values at each node:
 - $x_2 = \neg x_3 \Rightarrow$ add 2 clauses: $x_2 \vee \bar{x}_3, \bar{x}_2 \vee x_3$
 - $x_1 = x_4 \vee x_5 \Rightarrow$ add 3 clauses: $x_1 \vee \bar{x}_4, x_1 \vee \bar{x}_5, \bar{x}_1 \vee x_4 \vee x_5$
 - $x_0 = x_1 \wedge x_2 \Rightarrow$ add 3 clauses: $\bar{x}_0 \vee x_1, \bar{x}_0 \vee x_2, x_0 \vee \bar{x}_1 \vee \bar{x}_2$
- Hard-coded input values and output value.
 - $x_6 = 0 \Rightarrow$ add 1 clause: \bar{x}_6
 - $x_0 = 1 \Rightarrow$ add 1 clause: x_0
- Final step: turn clauses of length < 3 into clauses of length exactly 3.



3 Satisfiability Reduces to Independent Set




Claim. G contains independent set of size $k = |\Phi|$ iff Φ is satisfiable.

Pf. \Rightarrow Let S be independent set of size k.

- S must contain exactly one vertex in each triangle.
- Set these literals to true. \leftarrow and any other variables in a consistent way
- Truth assignment is consistent and all clauses are satisfied.

Pf. \Leftarrow Given satisfying assignment, select one true literal from each triangle. This is an independent set of size k. *



k = 3


$$\Phi = (\bar{x}_1 \vee x_2 \vee x_3) \wedge (\bar{x}_1 \vee \bar{x}_2 \vee x_3) \wedge (\bar{x}_1 \vee x_2 \vee x_4)$$

IS \leq_p VC


- Lemma: A set S is independent iff V-S is a vertex cover
- To reduce IS to VC, we show that we can determine if a graph has an independent set of size K by testing for a Vertex cover of size $n - K$

IS \leq_p VC

Find a maximum independent set S




Show that V-S is a vertex cover




Clique

- Clique
 - Graph $G = (V, E)$, a subset S of the vertices is a clique if there is an edge between every pair of vertices in S



Complement of a Graph

- Defn: $G' = (V, E')$ is the complement of $G = (V, E)$ if (u, v) is in E' iff (u, v) is not in E




IS \leq_p Clique

- Lemma: S is Independent in G iff S is a Clique in the complement of G
- To reduce IS to Clique, we compute the complement of the graph. The complement has a clique of size K iff the original graph has an independent set of size K

Hamiltonian Circuit Problem

- Hamiltonian Circuit – a simple cycle including all the vertices of the graph




Thm: Hamiltonian Circuit is NP Complete

- Reduction from 3-SAT


Traveling Salesman Problem

- Given a complete graph with edge weights, determine the shortest tour that includes all of the vertices (visit each vertex exactly once, and get back to the starting point)



Find the minimum cost tour

Thm: $HC <_P TSP$



Graph Coloring

- NP-Complete
 - Graph K-coloring
 - Graph 3-coloring
- Polynomial
 - Graph 2-Coloring

