

## 1 Asymptotics

Some properties of asymptotics:

- If  $f \leq O(g)$  and  $g \leq O(h)$  then  $f \leq O(h)$ .
- If  $f \geq \Omega(g)$  and  $g \geq \Omega(h)$  then  $f \geq \Omega(h)$ .
- If  $f = \Theta(g)$  and  $g = \Theta(h)$  then  $f = \Theta(h)$ .
- If  $f = O(h)$ ,  $g = O(h)$  then  $f + g = O(h)$ .

Some common running times:

- Polynomial:  $O(n^d)$ . Exponential  $2^{O(n)}$ , Logarithmic  $O(\log n)$ .
- For every positive  $\epsilon$  (no matter how small),  $\log n \leq O(n^\epsilon)$ . For every positive  $d$  (no matter how large),  $n^d \leq O(2^n)$ .

## 2 In class exercise

Arrange in increasing order of asymptotic growth. All logs are in base 2.

- $n^{5/3}$
- $2^{\sqrt{\log n}}$
- $\sqrt{n^n}$
- $\frac{n^2}{\log n}$
- $2^n$ .

**Hint:** Recall rules of logarithm

- $\log(a \cdot b) = \log a + \log b$ ,
- $\log(a/b) = \log a - \log b$ .
- $\log a^b = b \log a$ .

Always keep in mind  $n = 2^{\log_2 n}$ . For example,  $n^{1.5} = 2^{1.5 \log_2 n}$ . Also recall and that  $(2^a)^b = 2^{a \cdot b}$ .

### 3 Solution

In this part I will discuss the solution to the exercise. In many cases it might be difficult to directly compare two function  $f(n), g(n)$  asymptotically. An idea that usually helps out is to compare  $\log f(n)$  with  $\log g(n)$ . Recall that logarithm is an increasing function, so if  $f(n) > g(n)$  for some  $n > N$  then  $\log f(n) > \log g(n)$  for  $n > N$ . Here are two important rules when comparing the logs.

**When comparing logarithms ignore additive constants:** We said in class that  $n, 3n$  are asymptotically the same. If you take the log, then you are comparing  $\log n$  with  $\log n + \log 3$ . So, you can ignore the additive  $\log 3$ .

**When comparing logarithms, multiplicative constants matter:** Consider the two functions  $n, n^2$ . Obviously  $n^2$  grows asymptotically faster. When comparing the log, we have  $\log n, 2 \log n$ . So, the 2 multiplicative constant matters and shows that  $n^2$  grows faster.

Having said these, I will write down the solution to exercise. First, we calculate the logs:

a)  $\log n^{5/3} = \frac{5}{3} \log n$ .

b)  $\log 2^{\sqrt{\log n}} = \sqrt{\log n} \log 2 = \sqrt{\log n}$ .

c)  $\log \sqrt{n^n} = \frac{1}{2} \log n^n = \frac{n}{2} \log n$ .

d)  $\log \frac{n^2}{\log n} = \log n^2 - \log \log n = 2 \log n - \log \log n$ .

e)  $\log 2^n = n \log 2 = n$ .

Therefore,  $\sqrt{\log n} < \frac{5}{3} \log n < 2 \log n - \log \log n < n < \frac{n}{2} \log n$ . Note that when comparing  $\frac{5}{3} \log n$  and  $2 \log n - \log \log n$ ,  $\log \log n$  is a lower order term. So, first we compare the dominating terms  $\frac{5}{3} \log n$  and  $2 \log n$ . In this case the latter is bigger. If the dominating term were the same then we would have compared to lower order terms.