



CSE446: Linear Regression Regularization Bias / Variance Tradeoff Winter 2015

Luke Zettlemoyer

Slides adapted from Carlos Guestrin

Prediction of continuous variables

- Billionaire says: Wait, that's not what I meant!
- You say: Chill out, dude.
- He says: I want to predict a continuous variable for continuous inputs: I want to predict salaries from GPA.
- You say: **I can regress that...**





Linear Regression

Prediction

$$\hat{y} = w_0 + w_1 f_1(x)$$


Prediction

$$\hat{y}_i = w_0 + w_1 f_1(x) + w_2 f_2(x)$$



Ordinary Least Squares (OLS)

$$\text{total error} = \sum_i (y_i - \hat{y}_i)^2 = \sum_i \left(y_i - \sum_k w_k f_k(x_i) \right)^2$$




The regression problem


- Instances: $\langle \mathbf{x}_j, t_j \rangle$
- Learn: Mapping from \mathbf{x} to $t(\mathbf{x})$ $H = \{h_1, \dots, h_K\}$
- Hypothesis space:
 - Given, basis functions $\{h_1, \dots, h_k\}$
 - $h_i(\mathbf{x}) \in \mathbb{R}$
 - Find coeffs $\mathbf{w} = \{w_1, \dots, w_k\}$
 - Why is this usually called *linear regression*?
 - model is linear in the parameters
 - Can we estimate functions that are not lines???

Linear Basis: 1D input

Need a bias term: $\{h_1(x) = x, h_2(x)=1\}$



- Parabola: $\{h_1(\mathbf{x}) = \mathbf{x}^2, h_2(\mathbf{x})=\mathbf{x}, h_3(\mathbf{x})=1\}$



- 2D: $\{h_1(\mathbf{x}) = \mathbf{x}_1^2, h_2(\mathbf{x})=\mathbf{x}_2^2, h_3(\mathbf{x})=\mathbf{x}_1\mathbf{x}_2, \dots\}$
- Can define any basis functions $h_i(\mathbf{x})$ for n-dimensional input $\mathbf{x}=\langle x_1, \dots, x_n \rangle$

The regression problem

- Instances: $\langle \mathbf{x}_j, t_j \rangle$
- Learn: Mapping from x to $t(x)$ $H = \{h_1, \dots, h_K\}$

- Hypothesis space:

- Given, basis functions $\{h_1, \dots, h_k\}$
- $h_i(\mathbf{x}) \in \mathbb{R}$
- Find coeffs $\mathbf{w} = \{w_1, \dots, w_k\}$

$$t(\mathbf{x}) \underbrace{\approx \hat{f}(\mathbf{x})}_{\text{data}} = \sum_i w_i h_i(\mathbf{x})$$

- Why is this usually called *linear regression*?
 - model is linear in the parameters
 - Can we estimate functions that are not lines???

- Precisely, minimize the **residual squared error**:


$$\mathbf{w}^* = \arg \min_{\mathbf{w}} \sum_j \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$


Regression: matrix notation

$$\mathbf{w}^* = \arg \min_{\mathbf{w}} \sum_j \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$


$$\mathbf{w}^* = \arg \min_{\mathbf{w}} \underbrace{(\mathbf{H}\mathbf{w} - \mathbf{t})^T (\mathbf{H}\mathbf{w} - \mathbf{t})}_{\text{residual error}}$$

$$\mathbf{H} = \begin{bmatrix} h_1 & \dots & h_K \end{bmatrix}$$


K basis functions



N data points

$$\mathbf{w} = \begin{bmatrix} w_1 \\ \vdots \\ w_K \end{bmatrix}$$


K basis func

weights

$$\mathbf{t} = \begin{bmatrix} t_1 \\ \vdots \\ t_N \end{bmatrix}$$


N observed outputs

measurements

Regression: closed form solution

$$\mathbf{w}^* = \arg \min_{\mathbf{w}} \sum_j \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

$$\mathbf{w}^* = \arg \min_{\mathbf{w}} (\mathbf{H}\mathbf{w} - \mathbf{t})^T (\mathbf{H}\mathbf{w} - \mathbf{t})$$

$$\mathbf{F}(\mathbf{w}) = (\mathbf{H}\mathbf{w} - \mathbf{t})^T (\mathbf{H}\mathbf{w} - \mathbf{t})$$

$$\nabla_{\mathbf{w}} \mathbf{F}(\mathbf{w}) = \mathbf{0}$$

$$2\mathbf{H}^T (\mathbf{H}\mathbf{w} - \mathbf{t}) = \mathbf{0}$$

$$\mathbf{H}^T \mathbf{H}\mathbf{w} - \mathbf{H}^T \mathbf{t} = \mathbf{0}$$

$$\boxed{\mathbf{w}^* = (\mathbf{H}^T \mathbf{H})^{-1} \mathbf{H}^T \mathbf{t}}$$

Regression solution: simple matrix math

$$\mathbf{w}^* = \arg \min_{\mathbf{w}} \underbrace{(\mathbf{H}\mathbf{w} - \mathbf{t})^T (\mathbf{H}\mathbf{w} - \mathbf{t})}_{\text{residual error}}$$

$$\text{solution: } \mathbf{w}^* = \underbrace{(\mathbf{H}^T \mathbf{H})^{-1}}_{\mathbf{A}^{-1}} \underbrace{\mathbf{H}^T \mathbf{t}}_{\mathbf{b}} = \mathbf{A}^{-1} \mathbf{b}$$

where $\mathbf{A} = \mathbf{H}^T \mathbf{H} = \begin{bmatrix} & & \\ & & \\ & & \\ & & \\ & & \end{bmatrix}$ $\mathbf{b} = \mathbf{H}^T \mathbf{t} = \begin{bmatrix} & \\ & \\ & \\ & \end{bmatrix}$

$\underbrace{\hspace{10em}}$ $\underbrace{\hspace{10em}}$

$k \times k$ matrix
for k basis functions $k \times 1$ vector

But, why?

- Billionaire (again) says: Why sum squared error???
- You say: Gaussians, Dr. Gateson, Gaussians...
- Model: prediction is linear function plus Gaussian noise
 - $t(\mathbf{x}) = \sum_i w_i h_i(\mathbf{x}) + \varepsilon$
- Learn \mathbf{w} using MLE:

$$P(t \mid \mathbf{x}, \mathbf{w}, \sigma) = \frac{1}{\sigma \sqrt{2\pi}} e^{\frac{-[t - \sum_i w_i h_i(\mathbf{x})]^2}{2\sigma^2}}$$

Maximizing log-likelihood



Maximize wrt w:

$$\begin{aligned}\ln P(\mathcal{D} \mid \mathbf{w}, \sigma) &= \ln \left(\frac{1}{\sigma \sqrt{2\pi}} \right)^N \prod_{j=1}^N e^{\frac{-[t_j - \sum_i w_i h_i(\mathbf{x}_j)]^2}{2\sigma^2}} \\ \arg \max_w \ln \left(\frac{1}{\sigma \sqrt{2\pi}} \right)^N &+ \sum_{j=1}^N \frac{-[t_j - \sum_i w_i h_i(x_j)]^2}{2\sigma^2} \\ = \arg \max_w \sum_{j=1}^N &\frac{-[t_j - \sum_i w_i h_i(x_j)]^2}{2\sigma^2} \\ = \arg \min_w \sum_{j=1}^N &[t_j - \sum_i w_i h_i(x_j)]^2\end{aligned}$$

Least-squares Linear Regression is MLE for Gaussians!!!

Regularization in Linear Regression

- One sign of overfitting: large parameter values!



- *Regularized or penalized* regressions modified learning object to penalize large parameters

Ridge Regression

- Introduce a new objective function:

$$\hat{w}_{ridge} = \arg \min_w \sum_{j=1}^N \left(t(x_j) - (w_0 + \sum_{i=1}^k w_i h_i(x_j)) \right)^2 + \lambda \sum_{i=1}^k w_i^2$$

- Prefer low error but also add a squared penalty for large weights
- λ is hyperparameter that balances tradeoff
- Explicitly writing out bias feature (essentially $h_0=1$), which is not penalized

Ridge Regression: matrix notation

$$\hat{w}_{ridge} = \arg \min_w \sum_{j=1}^N \left(t(x_j) - (w_0 + \sum_{i=1}^k w_i h_i(x_j)) \right)^2 + \lambda \sum_{i=1}^k w_i^2$$

$$= \underbrace{\arg \min_{\mathbf{w}} (\mathbf{H}\mathbf{w} - \mathbf{t})^T (\mathbf{H}\mathbf{w} - \mathbf{t})}_{\text{residual error}} + \lambda \mathbf{w}^T I_{0+k} \mathbf{w}$$

$$\mathbf{H} = \begin{bmatrix} h_1 & \dots & h_K \\ \vdots & & \vdots \\ 1 & & 1 \end{bmatrix}$$

N data points

$\brace{ }$

bias column and k basis functions

$$\mathbf{w} = \begin{bmatrix} \vdots \\ w_k \\ \vdots \\ w_0 \end{bmatrix}$$

$\brace{ }$

k basis functs plus bias

weights

$$\mathbf{t} = \begin{bmatrix} t \\ \vdots \\ t \end{bmatrix}$$

$\brace{ }$

N observed outputs

measurements

$$I_{0+k} = \begin{bmatrix} 0 & 0 & 0 & \dots \\ 0 & 1 & 0 & \dots \\ 0 & 0 & 1 & \dots \\ \dots & \dots & \dots & \dots \end{bmatrix}_{k+1 \times k+1}$$

k+1 x k+1 identity matrix, but with 0 in upper left

Ridge Regression: closed form solution

$$\hat{w}_{ridge} = \arg \min_w \sum_{j=1}^N \left(t(x_j) - (w_0 + \sum_{i=1}^k w_i h_i(x_j)) \right)^2 + \lambda \sum_{i=1}^k w_i^2$$
$$= \arg \min_{\mathbf{w}} \underbrace{(\mathbf{H}\mathbf{w} - \mathbf{t})^T (\mathbf{H}\mathbf{w} - \mathbf{t})}_{\text{residual error}} + \lambda \mathbf{w}^T I_{0+k} \mathbf{w}$$

$$\mathbf{F}(\mathbf{w}) = (\mathbf{H}\mathbf{w} - \mathbf{t})^T (\mathbf{H}\mathbf{w} - \mathbf{t}) + \lambda \mathbf{w}^T I_{0+k} \mathbf{w}$$

$$\nabla_{\mathbf{w}} \mathbf{F}(\mathbf{w}) = \mathbf{0}$$

$$2\mathbf{H}^T (\mathbf{H}\mathbf{w} - \mathbf{t}) + 2\lambda I_{0+k} \mathbf{w} = \mathbf{0}$$

$$\mathbf{w}_{ridge}^* = (\mathbf{H}^T \mathbf{H} + \lambda I_{0+k})^{-1} \mathbf{H}^T \mathbf{t}$$

Regression solution: simple matrix math

$$\begin{aligned}\hat{w}_{ridge} &= \arg \min_w \sum_{j=1}^N \left(t(x_j) - (w_0 + \sum_{i=1}^k w_i h_i(x_j)) \right)^2 + \lambda \sum_{i=1}^k w_i^2 \\ &= \arg \min_{\mathbf{w}} \underbrace{(\mathbf{H}\mathbf{w} - \mathbf{t})^T (\mathbf{H}\mathbf{w} - \mathbf{t})}_{\text{residual error}} + \lambda \mathbf{w}^T I_{0+k} \mathbf{w}\end{aligned}$$

$$w_{ridge}^* = (\mathbf{H}^T \mathbf{H} + \lambda I_{0+k})^{-1} \mathbf{H}^T \mathbf{t}$$

Compare to un-regularized regression:

$$\mathbf{w}^* = (\mathbf{H}^T \mathbf{H})^{-1} \mathbf{H}^T \mathbf{t}$$


Ridge Regression

How does varying lambda change w?

$$\hat{w}_{ridge} = \arg \min_w \sum_{j=1}^N \left(t(x_j) - (w_0 + \sum_{i=1}^k w_i h_i(x_j)) \right)^2 + \lambda \sum_{i=1}^k w_i^2$$

- Larger λ ? Smaller λ ?
- As $\lambda \rightarrow 0$?
 - Becomes same a MLE, unregularized
- As $\lambda \rightarrow \infty$?
 - All weights will be 0!


Ridge Coefficient Path




From
Kevin Murphy
textbook

How to pick lambda?

- **Experimentation cycle**
 - Select a hypothesis f to best match training set
 - Tune hyperparameters on held-out set
 - Try many different values of lambda, pick best one
- **Or, can do k-fold cross validation**
 - No held-out set
 - Divide training set into k subsets
 - Repeatedly train on $k-1$ and test on remaining one
 - Average the results



Why squared regularization?



- Ridge:


$$\hat{w}_{ridge} = \arg \min_w \sum_{j=1}^N \left(t(x_j) - (w_0 + \sum_{i=1}^k w_i h_i(x_j)) \right)^2 + \lambda \sum_{i=1}^k w_i^2$$

- LASSO:


$$\hat{w}_{LASSO} = \arg \min_w \sum_{j=1}^N \left(t(x_j) - (w_0 + \sum_{i=1}^k w_i h_i(x_j)) \right)^2 + \lambda \sum_{i=1}^k |w_i|$$

- Linear penalty pushes more weights to zero
- Allows for a type of *feature selection*
- But, not differentiable and no closed form solution....

Geometric Intuition




Ridge Regression



Lasso

From
Rob
Tibshirani
slides


LASSO Coefficient Path




From
Kevin Murphy
textbook

Bias-Variance tradeoff – Intuition

- Model too simple: does not fit the data well
 - A *biased* solution



- Model too complex: small changes to the data, solution changes a lot
 - A *high-variance* solution



(Squared) Bias of learner

- Given: dataset D with m samples
- Learn: for different data D , you will learn different $h_D(x)$
- Expected prediction (averaged over hypotheses):

$$\bar{h}(x) = E_D[h_D(x)]$$

- Bias: expected difference between expected prediction and truth (here we square it)

$$E_x[(t(x) - \bar{h}(x))^2]$$

- Measures how well you expect to represent true solution
- Decreases with more complex model
- Zero bias typically means good predictions as $m \rightarrow \infty$

Variance of learner

- Given: dataset D with m samples $\bar{h}(x) = E_D[h_D(x)]$
- Learn: for different datasets D , you will learn different $h_D(x)$
- Variance: difference between what you expect to learn and what you learn from a particular dataset

$$E_D[E_x[(h_D(x) - \bar{h}(x))^2]]$$


- Measures how sensitive learner is to specific dataset
- Decreases with simpler model

Bias–Variance decomposition of error

$$\begin{aligned}MSE &= E_D[E_x[(t(x) - h_D(x))^2]] \\&= E_D[E_x[(t(x) - \bar{h}(x) + \bar{h}(x) - h_D(x))^2]] \\&= E_D[E_x[(t(x) - \bar{h}(x))^2]] + E_D[E_x[(\bar{h}(x) - h_D(x))^2]] \\&\quad + \underbrace{2E_D[E_x[(t(x) - \bar{h}(x))(\bar{h}(x) - h_D(x))]]}_{=0, \text{ because } \bar{h}(x) = E_D[h_D(x)]} \\&= \underbrace{E_x[(t(x) - \bar{h}(x))^2]}_{\text{Bias}} + \underbrace{E_D[E_x[(h_D(x) - \bar{h}(x))^2]]]}_{\text{Variance}}\end{aligned}$$

Bias-Variance Tradeoff

- Choice of hypothesis class introduces learning bias
 - More complex class → less bias
 - More complex class → more variance



Select points by clicking on the graph or press

[Example](#)

Degree of polynomial: Fit Y to X
 Fit X to Y

[Calculate](#) [View Polynomial](#) [Reset](#)



Select points by clicking on the graph or press

[Example](#)

Degree of polynomial: Fit Y to X
 Fit X to Y

[Calculate](#) [View Polynomial](#) [Reset](#)



Select points by clicking on the graph or press

[Example](#)

Degree of polynomial: Fit Y to X
 Fit X to Y

[Calculate](#) [View Polynomial](#) [Reset](#)

Training set error



$$\mathbf{w}^* = \arg \min_{\mathbf{w}} \sum_j \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

- Given a dataset (Training data)
- Choose a loss function
 - e.g., squared error (L_2) for regression
- **Training error:** For a particular set of parameters, loss function on training data:

$$error_{train}(\mathbf{w}) = \frac{1}{N_{train}} \sum_{j=1}^{N_{train}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

Training error as a function of model complexity


$$error_{train}(\mathbf{w}) = \frac{1}{N_{train}} \sum_{j=1}^{N_{train}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$



Select points by clicking on the graph or press [Example](#)

Degree of polynomial: Fit Y to X
 Fit X to Y


[Calculate](#) [View Polynomial](#) [Reset](#)



Select points by clicking on the graph or press [Example](#)

Degree of polynomial: Fit Y to X
 Fit X to Y

[Calculate](#) [View Polynomial](#) [Reset](#)



Prediction error




- Training set error can be poor measure of “quality” of solution
- Prediction error (true error): We really care about error over all possibilities:

$$\begin{aligned}
 error_{true}(\mathbf{w}) &= E_{\mathbf{x}} \left[\left(t(\mathbf{x}) - \sum_i w_i h_i(\mathbf{x}) \right)^2 \right] \\
 &= \int_{\mathbf{x}} \left(t(\mathbf{x}) - \sum_i w_i h_i(\mathbf{x}) \right)^2 p(\mathbf{x}) d\mathbf{x}
 \end{aligned}$$

Prediction error as a function of model complexity

$$error_{train}(\mathbf{w}) = \frac{1}{N_{train}} \sum_{j=1}^{N_{train}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

$$error_{true}(\mathbf{w}) = \int_{\mathbf{x}} \left(t(\mathbf{x}) - \sum_i w_i h_i(\mathbf{x}) \right)^2 p(\mathbf{x}) d\mathbf{x}$$



Computing prediction error

- To correctly predict error
 - Hard integral!
 - May not know $t(\mathbf{x})$ for every \mathbf{x} , may not know $p(\mathbf{x})$

$$error_{true}(\mathbf{w}) = \int_{\mathbf{x}} \left(t(\mathbf{x}) - \sum_i w_i h_i(\mathbf{x}) \right)^2 p(\mathbf{x}) d\mathbf{x}$$

- Monte Carlo integration (sampling approximation)
 - Sample a set of i.i.d. points $\{\mathbf{x}_1, \dots, \mathbf{x}_M\}$ from $p(\mathbf{x})$
 - Approximate integral with sample average

$$error_{true}(\mathbf{w}) \approx \frac{1}{M} \sum_{j=1}^M \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

Why training set error doesn't approximate prediction error?

- Sampling approximation of prediction error:

$$error_{true}(\mathbf{w}) \approx \frac{1}{M} \sum_{j=1}^M \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

- Training error :

$$error_{train}(\mathbf{w}) = \frac{1}{N_{train}} \sum_{j=1}^{N_{train}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

- Very similar equations!!!

- Why is training set a bad measure of prediction error???

Why training set error doesn't approximate prediction error?

- Same samples
 - Training error
 - Very similar equations!!!
 - Why is training set a bad measure of prediction error???
- Because you cheated!!!**

Training error good estimate for a single w ,
But you optimized w with respect to the training error,
and found w that is good for this set of samples

Training error is a (optimistically) biased estimate of prediction error

Test set error

$$\mathbf{w}^* = \arg \min_{\mathbf{w}} \sum_j \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

- Given a dataset, **randomly** split it into two parts:
 - Training data – $\{\mathbf{x}_1, \dots, \mathbf{x}_{N_{train}}\}$
 - Test data – $\{\mathbf{x}_1, \dots, \mathbf{x}_{N_{test}}\}$
- Use training data to optimize parameters \mathbf{w}
- **Test set error:** For the *final solution* \mathbf{w}^* , evaluate the error using:



$$error_{test}(\mathbf{w}) = \frac{1}{N_{test}} \sum_{j=1}^{N_{test}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

Test set error as a function of model complexity

$$error_{train}(\mathbf{w}) = \frac{1}{N_{train}} \sum_{j=1}^{N_{train}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

$$error_{true}(\mathbf{w}) = \int_{\mathbf{x}} \left(t(\mathbf{x}) - \sum_i w_i h_i(\mathbf{x}) \right)^2 p(\mathbf{x}) d\mathbf{x}$$

$$error_{test}(\mathbf{w}) = \frac{1}{N_{test}} \sum_{j=1}^{N_{test}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$



Degree of polynomial: 1 Fit Y to X
 Fit X to Y



Degree of polynomial: 13 Fit Y to X
 Fit X to Y

Overfitting: this slide is so important we are looking at it again!

- Assume:
 - Data generated from distribution $D(X, Y)$
 - A hypothesis space H
- Define: errors for hypothesis $h \in H$
 - Training error: $error_{train}(h)$
 - Data (true) error: $error_{true}(h)$
- We say h **overfits** the training data if there exists an $h' \in H$ such that:

$$error_{train}(h) < error_{train}(h')$$

and

$$error_{true}(h) > error_{true}(h')$$

Summary: error estimators

- Gold Standard:

$$error_{true}(\mathbf{w}) = \int_{\mathbf{x}} \left(t(\mathbf{x}) - \sum_i w_i h_i(\mathbf{x}) \right)^2 p(\mathbf{x}) d\mathbf{x}$$


- Training: optimistically biased

$$error_{train}(\mathbf{w}) = \frac{1}{N_{train}} \sum_{j=1}^{N_{train}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

- Test: our final measure

$$error_{test}(\mathbf{w}) = \frac{1}{N_{test}} \sum_{j=1}^{N_{test}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$


Error as a function of number of training examples for a fixed model complexity



$$error_{train}(\mathbf{w}) = \frac{1}{N_{train}} \sum_{j=1}^{N_{train}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

$$error_{true}(\mathbf{w}) = \int_{\mathbf{x}} \left(t(\mathbf{x}) - \sum_i w_i h_i(\mathbf{x}) \right)^2 p(\mathbf{x}) d\mathbf{x}$$

$$error_{test}(\mathbf{w}) = \frac{1}{N_{test}} \sum_{j=1}^{N_{test}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$




little data

infinite data

Error as function of regularization parameter, fixed model complexity

$$error_{train}(\mathbf{w}) = \frac{1}{N_{train}} \sum_{j=1}^{N_{train}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

$$error_{true}(\mathbf{w}) = \int_{\mathbf{x}} \left(t(\mathbf{x}) - \sum_i w_i h_i(\mathbf{x}) \right)^2 p(\mathbf{x}) d\mathbf{x}$$



Summary: error estimators

- Good news: test set unbiased
- Be careful!!!
Test set only unbiased if you never never ever ever do any any any any learning on the test data
- Test: our final measure
For example, if you use the test set to select the degree of the polynomial... no longer unbiased!!!
(We will address this problem later in the semester)
- Test: our final measure

$$error_{test}(\mathbf{w}) = \frac{1}{N_{test}} \sum_{j=1}^{N_{test}} \left(t(\mathbf{x}_j) - \sum_i w_i h_i(\mathbf{x}_j) \right)^2$$

What you need to know

- Regression
 - Basis function = features
 - Optimizing sum squared error
 - Relationship between regression and Gaussians
- Regularization
 - Ridge regression math
 - LASSO Formulation
 - How to set lambda
- Bias-Variance trade-off