

CSE 446: Machine Learning

Homework 0

Due: 1/13/20 9:30 AM

This assignment is meant to be a review of prerequisite knowledge for this course. Your work will be graded based on completion. If an honest attempt was made on each problem, you may receive full marks.

Collaboration: Make certain that you understand the course collaboration policy, described on the course website. You must complete this assignment **individually**; you are **not** allowed to collaborate with anyone else. You may *discuss* the homework to understand the problems and the mathematics behind the various learning algorithms, but **you are not allowed to share problem solutions with any other students**. You must also not consult code on the internet that is directly related to the programming exercise.

You are also prohibited from posting any part of your solution to the internet, even after the course is complete. Similarly, please don't post this PDF file or the homework skeleton code to the internet.

Formatting: You must submit your solutions as a single PDF document. Your work can be either handwritten or typeset, but keep in mind that illegible submissions (as determined by the TA's subjective opinion) will be given no credit. For this reason, we **strongly** encourage you to write up your responses using L^AT_EX.

Submission: You will submit your assignment on Gradescope. Instructions to submit on Gradescope can be found here: https://courses.cs.washington.edu/courses/csep521/19wi/gradescope_guide.pdf.

Probability and Statistics

- [2 points]* (Bayes Rule, from Murphy exercise 2.4.) After your yearly checkup, the doctor has bad news and good news. The bad news is that you tested positive for a serious disease, and that the test is 99% accurate (i.e., the probability of testing positive given that you have the disease is 0.99, as is the probability of testing negative given that you don't have the disease). The good news is that this is a rare disease, striking only one in 10,000 people. What are the chances that you actually have the disease? (Show your calculations as well as giving the final result.)
- For any two random variables X, Y the *covariance* is defined as $\text{Cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}[X])(Y - \mathbb{E}[Y])]$. You may assume X and Y take on a discrete values if you find that is easier to work with.
 - [1 points]* If $\mathbb{E}[Y|X = x] = x$ show that $\text{Cov}(X, Y) = \mathbb{E}[(X - \mathbb{E}[X])^2]$.
 - [1 points]* If X, Y are independent show that $\text{Cov}(X, Y) = 0$.
- Let X and Y be independent random variables with PDFs given by f and g , respectively. Let h be the PDF of the random variable $Z = X + Y$.
 - [2 points]* Show that $h(z) = \int_{-\infty}^{\infty} f(x)g(z-x)dx$. (If you are more comfortable with discrete probabilities, you can instead derive an analogous expression for the discrete case, and then you should give a one sentence explanation as to why your expression is analogous to the continuous case.)
 - [1 points]* If X and Y are both independent and uniformly distributed on $[0, 1]$ (i.e. $f(x) = g(x) = 1$ for $x \in [0, 1]$ and 0 otherwise) what is h , the PDF of $Z = X + Y$?

4. [1 points] A random variable $X \sim \mathcal{N}(\mu, \sigma^2)$ is Gaussian distributed with mean μ and variance σ^2 . Given that for any $a, b \in \mathbb{R}$, we have that $Y = aX + b$ is also Gaussian, find a, b such that $Y \sim \mathcal{N}(0, 1)$.

5. [2 points] For a random variable Z , its mean and variance are defined as $\mathbb{E}[Z]$ and $\mathbb{E}[(Z - \mathbb{E}[Z])^2]$, respectively. Let X_1, \dots, X_n be independent and identically distributed random variables, each with mean μ and variance σ^2 . If we define $\hat{\mu}_n = \frac{1}{n} \sum_{i=1}^n X_i$, what is the mean and variance of $\sqrt{n}(\hat{\mu}_n - \mu)$?

6. If $f(x)$ is a PDF, the cumulative distribution function (CDF) is defined as $F(x) = \int_{-\infty}^x f(y)dy$. For any function $g : \mathbb{R} \rightarrow \mathbb{R}$ and random variable X with PDF $f(x)$, recall that the expected value of $g(X)$ is defined as $\mathbb{E}[g(X)] = \int_{-\infty}^{\infty} g(y)f(y)dy$. For a boolean event A , define $\mathbf{1}\{A\}$ as 1 if A is true, and 0 otherwise. Thus, $\mathbf{1}\{x \leq a\}$ is 1 whenever $x \leq a$ and 0 whenever $x > a$. Note that $F(x) = \mathbb{E}[\mathbf{1}\{X \leq x\}]$. Let X_1, \dots, X_n be independent and identically distributed random variables with CDF $F(x)$. Define $\hat{F}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbf{1}\{X_i \leq x\}$. Note, for every x , that $\hat{F}_n(x)$ is an empirical estimate of $F(x)$. You may use your answers to the previous problem.

a. [1 points] For any x , what is $\mathbb{E}[\hat{F}_n(x)]$?

b. [1 points] For any x , the variance of $\hat{F}_n(x)$ is $\mathbb{E}[(\hat{F}_n(x) - F(x))^2]$. Show that $\text{Variance}(\hat{F}_n(x)) = \frac{F(x)(1-F(x))}{n}$.

c. [1 points] Using your answer to b, show that for all $x \in \mathbb{R}$, we have $\mathbb{E}[(\hat{F}_n(x) - F(x))^2] \leq \frac{1}{4n}$.

Linear Algebra and Vector Calculus

7. (Rank) Let $A = \begin{bmatrix} 1 & 2 & 1 \\ 1 & 0 & 3 \\ 1 & 1 & 2 \end{bmatrix}$ and $B = \begin{bmatrix} 1 & 2 & 3 \\ 1 & 0 & 1 \\ 1 & 1 & 2 \end{bmatrix}$. For each matrix A and B ,

a. [2 points] what is its rank?

b. [2 points] what is a (minimal size) basis for its column span?

8. (Linear equations) Let $A = \begin{bmatrix} 0 & 2 & 4 \\ 2 & 4 & 2 \\ 3 & 3 & 1 \end{bmatrix}$, $b = [-2 \quad -2 \quad -4]^T$, and $c = [1 \quad 1 \quad 1]^T$.

a. [1 points] What is Ac ?

b. [2 points] What is the solution to the linear system $Ax = b$? (Show your work).

9. (Hyperplanes) Assume w is an n -dimensional vector and b is a scalar. A hyperplane in \mathbb{R}^n is the set $\{x : x \in \mathbb{R}^n, \text{ s.t. } w^T x + b = 0\}$.

a. [1 points] ($n = 2$ example) Draw the hyperplane for $w = [-1, 2]^T$, $b = 2$? Label your axes.

b. [1 points] ($n = 3$ example) Draw the hyperplane for $w = [1, 1, 1]^T$, $b = 0$? Label your axes.

c. [2 points] Given some $x_0 \in \mathbb{R}^n$, find the squared distance to the hyperplane defined by $w^T x + b = 0$. In other words, solve the following optimization problem:

$$\begin{aligned} \min_x & \|x_0 - x\|^2 \\ \text{s.t.} & w^T x + b = 0 \end{aligned}$$

(Hint: if \tilde{x}_0 is the minimizer of the above problem, note that $\|x_0 - \tilde{x}_0\| = \left| \frac{w^T(x_0 - \tilde{x}_0)}{\|w\|} \right|$. What is $w^T \tilde{x}_0$?)

10. For possibly non-symmetric $\mathbf{A}, \mathbf{B} \in \mathbb{R}^{n \times n}$ and $c \in \mathbb{R}$, let $f(x, y) = x^T \mathbf{A}x + y^T \mathbf{B}y + c$. Define $\nabla_z f(x, y) = \left[\frac{\partial f(x, y)}{\partial z_1} \quad \frac{\partial f(x, y)}{\partial z_2} \quad \dots \quad \frac{\partial f(x, y)}{\partial z_n} \right]^T$.

- a. [2 points] Explicitly write out the function $f(x, y)$ in terms of the components $A_{i,j}$ and $B_{i,j}$ using appropriate summations over the indices.
- b. [2 points] What is $\nabla_x f(x, y)$ in terms of the summations over indices *and* vector notation?
- c. [2 points] What is $\nabla_y f(x, y)$ in terms of the summations over indices *and* vector notation?

Programming

11. For the A, b, c as defined in Problem 8, use NumPy to compute (take a screen shot of your answer):

- a. [2 points] What is A^{-1} ?
- b. [1 points] What is $A^{-1}b$? What is Ac ?

12. [4 points] Two random variables X and Y have equal distributions if their CDFs, F_X and F_Y , respectively, are equal, i.e. for all x , $|F_X(x) - F_Y(x)| = 0$. The central limit theorem says that the sum of k independent, zero-mean, variance-1/ k random variables converges to a (standard) Normal distribution as k goes off to infinity. We will study this phenomenon empirically (you will use the Python packages Numpy and Matplotlib). Define $Y^{(k)} = \frac{1}{\sqrt{k}} \sum_{i=1}^k B_i$ where each B_i is equal to -1 and 1 with equal probability. From your solution to problem 5, we know that $\frac{1}{\sqrt{k}} B_i$ is zero-mean and has variance $1/k$.

- a. For $i = 1, \dots, n$ let $Z_i \sim \mathcal{N}(0, 1)$. If $F(x)$ is the true CDF from which each Z_i is drawn (i.e., Gaussian) and $\hat{F}_n(x) = \frac{1}{n} \sum_{i=1}^n \mathbf{1}\{Z_i \leq x\}$, use the answer to problem 1.5 above to choose n large enough such that, for all $x \in \mathbb{R}$, $\sqrt{\mathbb{E}[(\hat{F}_n(x) - F(x))^2]} \leq 0.0025$, and plot $\hat{F}_n(x)$ from -3 to 3 .
(Hint: use `Z=npumpy.random.randn(n)` to generate the random variables, and `import matplotlib.pyplot as plt;`
`plt.step(sorted(Z), np.arange(1,n+1)/float(n))` to plot).
- b. For each $k \in \{1, 8, 64, 512\}$ generate n independent copies $Y^{(k)}$ and plot their empirical CDF on the same plot as part a.
(Hint: `np.sum(np.sign(np.random.randn(n, k))*np.sqrt(1./k), axis=1)` generates n of the $Y^{(k)}$ random variables.)

Be sure to always label your axes. Your plot should look something like the following (Tip: checkout `seaborn` for instantly better looking plots.)

