

What are particle systems?

A **particle system** is a collection of point masses that obeys some physical laws (e.g, gravity or spring behaviors).

Particle systems can be used to simulate all sorts of physical phenomena:

- Smoke
- Snow
- Fireworks
- Hair
- Cloth
- Snakes
- Fish

3

Lecture 15: Particle Systems

Overview

1. One lousy particle
2. Particle systems
3. Forces: gravity, springs
4. Implementation

Reading

- Required:
 - Witkin, *Particle System Dynamics*, SIGGRAPH '97 course notes on Physically Based Modeling.
- Optional
 - Witkin and Baraff, *Differential Equation Basics*, SIGGRAPH '97 course notes on Physically Based Modeling.
 - Hockney and Eastwood. *Computer simulation using particles*. Adam Hilger, New York, 1988.
 - Gavin Miller. “The motion dynamics of snakes and worms.” *Computer Graphics* 22:169-178, 1988.


4

2

Diff eqs and integral curves

The equation $\dot{\mathbf{x}} = \mathbf{g}(\mathbf{x}, t)$ is actually a **first order differential equation**.

We can solve for \mathbf{x} through time by starting at an initial point and stepping along the vector field:



This is called an **initial value problem** and the solution is called an **integral curve**.


7

Particle in a flow field

We begin with a single particle with:

$$\begin{aligned} &- \text{Position, } \mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix} \\ &- \text{Velocity, } \mathbf{v} \equiv \dot{\mathbf{x}} = \frac{d\mathbf{x}}{dt} = \begin{bmatrix} dx/dt \\ dy/dt \end{bmatrix} \end{aligned}$$

Suppose the velocity is dictated by some driving function \mathbf{g} :



5

Euler's method


One simple approach is to choose a time step, Δt , and take linear steps along the flow:

$$\begin{aligned} \mathbf{x}(t+\Delta t) &= \mathbf{x}(t) + \Delta t \cdot \dot{\mathbf{x}}(t) \\ &= \mathbf{x}(t) + \Delta t \cdot \mathbf{g}(\mathbf{x}, t) \end{aligned}$$


This approach is called **Euler's method** and looks like:

Properties:

- Simplest numerical method
- Bigger steps, bigger errors



Need to take pretty small steps, so not very efficient. Better (more complicated) methods exist, e.g., "Runge-Kutta."



How does our particle move through the vector field?

8

6

Phase space

$$\begin{bmatrix} \mathbf{x} \\ \mathbf{v} \end{bmatrix}$$

Concatenate \mathbf{x} and \mathbf{v} to make a 6-vector: position
in **phase space**.

$$\begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\mathbf{v}} \end{bmatrix}$$

Taking the time derivative: another 6-vector.

$$\begin{bmatrix} \ddot{\mathbf{x}} \\ \ddot{\mathbf{v}} \end{bmatrix} = \begin{bmatrix} \mathbf{v} \\ \mathbf{f}/m \end{bmatrix}$$

A vanilla 1st-order differential equation.

- Now consider a particle in a force field \mathbf{f} .
- In this case, the particle has:
 - Mass, m
 - Acceleration, $\mathbf{a} \equiv \ddot{\mathbf{x}} = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$

- The particle obeys Newton's law: $\mathbf{f} = m\mathbf{a} = m\ddot{\mathbf{x}}$

- The force field \mathbf{f} can in general depend on the position and velocity of the particle as well as time.
- Thus, with some rearrangement, we end up with:

$$\ddot{\mathbf{x}} = \frac{\mathbf{f}(\mathbf{x}, \dot{\mathbf{x}}, t)}{m}$$

11

9

Particle in a force field

- Now consider a particle in a force field \mathbf{f} .
- In this case, the particle has:
 - Mass, m
 - Acceleration, $\mathbf{a} \equiv \ddot{\mathbf{x}} = \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$

- The particle obeys Newton's law: $\mathbf{f} = m\mathbf{a} = m\ddot{\mathbf{x}}$


- The force field \mathbf{f} can in general depend on the position and velocity of the particle as well as time.
- Thus, with some rearrangement, we end up with:

$$\ddot{\mathbf{x}} = \frac{\mathbf{f}(\mathbf{x}, \dot{\mathbf{x}}, t)}{m}$$

11

9

Particle structure



Second order equations

$$\text{This equation: } \ddot{\mathbf{x}} = \frac{\mathbf{f}(\mathbf{x}, \mathbf{v}, t)}{m}$$

is a **second order differential equation**.

Our solution method, though, worked on first order differential equations.

$$\begin{bmatrix} \dot{\mathbf{x}} = \mathbf{v} \\ \dot{\mathbf{v}} = \frac{\mathbf{f}(\mathbf{x}, \mathbf{v}, t)}{m} \end{bmatrix}$$


where we have added a new variable \mathbf{v} to get a pair of **coupled first order equations**.

12

10

Solver interface


Solver interface



15


Forces

- Constant (gravity)
- Position/time dependent (force fields)
- Velocity-dependent (drag)
- N-ary (springs)



13

Particle systems



16


Damped spring

Force law:

$$\mathbf{f}_1 = -k_s(|\Delta \mathbf{x}| - \mathbf{r}) + k_d \left(\frac{\Delta \mathbf{v}_{\Delta \mathbf{x}}}{|\Delta \mathbf{x}|} \right) \frac{\Delta \mathbf{x}}{|\Delta \mathbf{x}|}$$

$$\mathbf{f}_2 = -\mathbf{f}_1$$

\mathbf{r} = rest length



$$\Delta \mathbf{v} = \mathbf{v}_1 - \mathbf{v}_2$$

19

Gravity


Force law:

$$\mathbf{f}_{grav} = m \mathbf{G}$$

$$\mathbf{p} \rightarrow \mathbf{f} \quad + = \quad \mathbf{p} \rightarrow \mathbf{m} \quad * \quad \mathbf{F} \rightarrow \mathbf{G}$$

17

Particle systems with forces




$$\mathbf{f}_{drag} = -k_{drag} \mathbf{v}$$

$$\mathbf{p} \rightarrow \mathbf{f} \quad - = \quad \mathbf{F} \rightarrow \mathbf{k} \quad * \quad \mathbf{p} \rightarrow \mathbf{v}$$

20

Solver interface



23

derivEval loop


1. Clear forces
 - Loop over particles, zero force accumulators
2. Calculate forces
 - Sum all forces into accumulators
3. Gather
 - Loop over particles, copying \mathbf{v} and \mathbf{f}/m into destination array

21

Differential equation solver

$$\text{Euler method: } \begin{bmatrix} \mathbf{x}_1^{i+1} \\ \mathbf{v}_1^{i+1} \\ \vdots \\ \mathbf{x}_n^{i+1} \\ \mathbf{v}_n^{i+1} \end{bmatrix} = \begin{bmatrix} \mathbf{x}_1^i \\ \mathbf{v}_1^i \\ \vdots \\ \mathbf{x}_n^i \\ \mathbf{v}_n^i \end{bmatrix} + \Delta t \begin{bmatrix} \mathbf{v}_1^i / m_1 \\ \mathbf{f}_1^i / m_1 \\ \vdots \\ \mathbf{v}_n^i / m_n \\ \mathbf{f}_n^i / m_n \end{bmatrix}$$

derivEval Loop




24

22

Collision Detection


Bouncing off the walls

- $(\mathbf{X} - \mathbf{P}) \cdot \mathbf{N} < \varepsilon$ Within ε of the wall
- $\mathbf{N} \cdot \mathbf{V} < 0$ Heading in



27


- Add-on for a particle simulator
- For now, just simple point-plane collisions



25

Collision Response


Normal and tangential components



$$\mathbf{V}' = \mathbf{V}_T - k_r \mathbf{V}_N$$

after

before



28

$$\begin{aligned}\mathbf{V}_N &= (\mathbf{N} \cdot \mathbf{V}) \mathbf{N} \\ \mathbf{V}_T &= \mathbf{V} - \mathbf{V}_N\end{aligned}$$

26

Summary

What you should take away from this lecture:

- The meanings of all the **boldfaced** terms
- Euler method for solving differential equations
- Combining particles into a particle system
- Physics of a particle system
- Various forces acting on a particle
- Simple collision detection