

Particle Systems

Zoran Popovic
CSE 457
Spring 2019

Reading

- Required:
 - Witkin, *Particle System Dynamics*, SIGGRAPH '97 course notes on Physically Based Modeling.
- Optional
 - Witkin and Baraff, *Differential Equation Basics*, SIGGRAPH '97 course notes on Physically Based Modeling.
 - Hockney and Eastwood. *Computer simulation using particles*. Adam Hilger, New York, 1988.
 - Gavin Miller. "The motion dynamics of snakes and worms." *Computer Graphics* 22:169-178, 1988.

2

What are particle systems?

A **particle system** is a collection of point masses that obeys some physical laws (e.g. gravity or spring behaviors).

Particle systems can be used to simulate all sorts of physical phenomena:

- Smoke
- Snow
- Fireworks
- Hair
- Cloth
- Snakes
- Fish

3

Overview

1. One lousy particle
2. Particle systems
3. Forces: gravity, springs
4. Implementation


4

Particle in a flow field

We begin with a single particle with:

- Position, $\mathbf{x} = \begin{bmatrix} x \\ y \end{bmatrix}$
- Velocity, $\mathbf{v} \equiv \frac{d\mathbf{x}}{dt} = \begin{bmatrix} dx/dt \\ dy/dt \end{bmatrix}$


Suppose the velocity is dictated by some driving function \mathbf{g} :



5

Vector fields

At any moment in time, the function \mathbf{g} defines a vector field over \mathbf{x} :




How does our particle move through the vector field?

6

Diff eqs and integral curves

The equation $\dot{\mathbf{x}} = \mathbf{g}(\mathbf{x}, t)$ is actually a **first order differential equation**.

We can solve for \mathbf{x} through time by starting at an initial point and stepping along the vector field:



This is called an **initial value problem** and the solution is called an **integral curve**.

7

Euler's method


One simple approach is to choose a time step, Δt , and take linear steps along the flow:

$$\begin{aligned} \mathbf{x}(t+\Delta t) &= \mathbf{x}(t) + \Delta t \cdot \dot{\mathbf{x}} \\ &= \mathbf{x}(t) + \Delta t \cdot \mathbf{g}(\mathbf{x}, t) \end{aligned}$$

This approach is called **Euler's method** and looks like:

Properties:

- Simplest numerical method
- Bigger steps, bigger errors



Need to take pretty small steps, so not very efficient. Better (more complicated) methods exist, e.g., "Runge-Kutta."

8

Particle in a force field

- Now consider a particle in a force field \mathbf{f} .
- In this case, the particle has:
 - Mass, m
 - Acceleration, $\mathbf{a} \equiv \frac{d\mathbf{v}}{dt} = \frac{d^2\mathbf{x}}{dt^2}$
- The particle obeys Newton's law: $\mathbf{f} = m\mathbf{a} = m\ddot{\mathbf{x}}$
- The force field \mathbf{f} can in general depend on the position and velocity of the particle as well as time.
- Thus, with some rearrangement, we end up with:

$$\ddot{\mathbf{x}} = \frac{\mathbf{f}(\mathbf{x}, \mathbf{v}, t)}{m}$$

9

Second order equations

This equation: $\ddot{\mathbf{x}} = \frac{\mathbf{f}(\mathbf{x}, \mathbf{v}, t)}{m}$

is a **second order differential equation**.

Our solution method, though, worked on first order differential equations.

We can rewrite this as:

$$\begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\mathbf{v}} \end{bmatrix} = \begin{bmatrix} \mathbf{v} \\ \frac{\mathbf{f}(\mathbf{x}, \mathbf{v}, t)}{m} \end{bmatrix}$$

where we have added a new variable \mathbf{v} to get a pair of **coupled first order equations**.

10

Phase space

$$\begin{bmatrix} \mathbf{x} \\ \mathbf{v} \end{bmatrix}$$

Concatenate \mathbf{x} and \mathbf{v} to make a 6-vector: position in **phase space**.

$$\begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\mathbf{v}} \end{bmatrix}$$


Taking the time derivative: another 6-vector.

$$\begin{bmatrix} \dot{\mathbf{x}} \\ \dot{\mathbf{v}} \end{bmatrix} = \mathbf{f}$$

A vanilla 1st-order differential equation.


11

Particle structure




12

Solver interface



13


Particle systems



$$\begin{bmatrix} \mathbf{x}_1 \\ \mathbf{v}_1 \\ \mathbf{f}_1 \\ m_1 \end{bmatrix} \begin{bmatrix} \mathbf{x}_2 \\ \mathbf{v}_2 \\ \mathbf{f}_2 \\ m_2 \end{bmatrix} \begin{bmatrix} \mathbf{x}_3 \\ \mathbf{v}_3 \\ \mathbf{f}_3 \\ m_3 \end{bmatrix} \dots \begin{bmatrix} \mathbf{x}_n \\ \mathbf{v} \\ \mathbf{f}_n \\ m_n \end{bmatrix}$$

14

Solver interface



15

Forces

- Constant (gravity)
- Position/time dependent (force fields)
- Velocity-dependent (drag)
- N-ary (springs)

16

Gravity

Force law:

$$\mathbf{f}_{grav} = m\mathbf{G}$$

$$\mathbf{p} \rightarrow \mathbf{f} += \mathbf{p} \rightarrow \mathbf{m} * \mathbf{F} \rightarrow \mathbf{G}$$

17

Viscous drag

Force law:

$$\mathbf{f}_{drag} = -k_{drag} \mathbf{v}$$

$$\mathbf{p} \rightarrow \mathbf{f} -= \mathbf{F} \rightarrow \mathbf{k} * \mathbf{p} \rightarrow \mathbf{v}$$

18

Damped spring


Force law:

$$\mathbf{f}_1 = -k_s \left(\frac{\mathbf{x}_1 - \mathbf{x}_2}{r} - 1 \right)$$

$$\mathbf{f}_2 = -\mathbf{f}_1$$


$$k_d \left(\frac{\mathbf{v}_1 - \mathbf{v}_2}{r} \right)$$

r = rest length



19

Particle systems with forces




20

derivEval loop

1. Clear forces
 - Loop over particles, zero force accumulators
2. Calculate forces
 - Sum all forces into accumulators
3. Gather
 - Loop over particles, copying v and f/m into destination array


21

derivEval Loop



22

Solver interface



23

Differential equation solver


$$\begin{bmatrix} \dot{\mathbf{x}}_1 \\ \dot{\mathbf{v}}_1 \\ \vdots \\ \dot{\mathbf{x}}_n \\ \dot{\mathbf{v}}_n \end{bmatrix} = \mathbf{f}$$

Euler method:

$$\begin{bmatrix} \mathbf{x}_1^{i+1} \\ \mathbf{v}_1^{i+1} \\ \vdots \\ \mathbf{x}_n^{i+1} \\ \mathbf{v}_n^{i+1} \end{bmatrix} = \begin{bmatrix} \mathbf{x}_1^i \\ \mathbf{v}_1^i \\ \vdots \\ \mathbf{x}_n^i \\ \mathbf{v}_n^i \end{bmatrix} + \begin{bmatrix} \mathbf{v}_1^i \\ \mathbf{f}_1^i / m_1 \\ \vdots \\ \mathbf{v}_n^i \\ \mathbf{f}_n^i / m_n \end{bmatrix} \Delta t$$

24



Bouncing off the walls



- Add-on for a particle simulator
- For now, just simple point-plane collisions

25

Normal and tangential components




$$\mathbf{V}_N = (\mathbf{N} \cdot \mathbf{V})\mathbf{N}$$

$$\mathbf{V}_T = \mathbf{V} - \mathbf{V}_N$$

26


Collision Detection




$(\mathbf{X} - \mathbf{P}) \cdot \mathbf{N} < \varepsilon$ Within ε of the wall
 $\mathbf{N} \cdot \mathbf{V} < 0$ Heading in

27

Collision Response



before




after

$$\mathbf{V}' = \mathbf{V}_T - k_r \mathbf{V}_N$$

28

Artificial Fish



29

Related Research

- Determining dynamic parameters for cloth simulation

30

Summary

What you should take away from this lecture:

- The meanings of all the **boldfaced** terms
- Euler method for solving differential equations
- Combining particles into a particle system
- Physics of a particle system
- Various forces acting on a particle
- Simple collision detection

31