Rational decisions

Chapter 16

Outline

- ◇ Rational preferences
- ◇ Utilities
- ◇ Money
- ◇ Multiattribute utilities
- ◇ Decision networks
- ◇ Value of information

Preferences

An agent chooses among prizes (A, B, etc.) and lotteries, i.e., situations with uncertain prizes



Lottery $L = [p, A;\ (1-p), B]$

Notation:

$A \succ B$	A preferred to B
$A \sim B$	indifference between A and B
$A \succsim B$	B not preferred to A

Rational preferences

Idea: preferences of a rational agent must obey constraints.
Rational preferences $\Rightarrow$
behavior describable as maximization of expected utility

Constraints:

Orderability
$(A \succ B) \vee (B \succ A) \vee (A \sim B)$

Transitivity
$(A \succ B) \wedge (B \succ C) \Rightarrow (A \succ C)$

Continuity
$A \succ B \succ C \Rightarrow \exists p\ [p, A;\ 1-p, C] \sim B$

Substitutability
$A \sim B \Rightarrow [p, A;\ 1-p, C] \sim [p, B; 1-p, C]$

Monotonicity
$A \succ B \Rightarrow (p \geq q \Leftrightarrow [p, A;\ 1-p, B] \succsim [q, A;\ 1-q, B])$

Rational preferences contd.

Violating the constraints leads to self-evident irrationality

For example: an agent with intransitive preferences can be induced to give away all its money

If $B \succ C$, then an agent who has C would pay (say) 1 cent to get B

If $A \succ B$, then an agent who has B would pay (say) 1 cent to get A

If $C \succ A$, then an agent who has A would pay (say) 1 cent to get C



Maximizing expected utility

Theorem (Ramsey, 1931; von Neumann and Morgenstern, 1944):
Given preferences satisfying the constraints
there exists a real-valued function U such that

$U(A) \geq U(B) \quad \Leftrightarrow \quad A \succsim B$
$U([p_1, S_1;\ \ldots\ ;\ p_n, S_n]) = \Sigma_i\ p_i U(S_i)$

MEU principle:
Choose the action that maximizes expected utility

Note: an agent can be entirely rational (consistent with MEU) without ever representing or manipulating utilities and probabilities

E.g., a lookup table for perfect tictactoe

Utilities

Utilities map states to real numbers. Which numbers?

Standard approach to assessment of human utilities:
compare a given state A to a standard lottery L_p that has
"best possible prize" $u_\top$ with probability p
"worst possible catastrophe" $u_\bot$ with probability $(1-p)$
adjust lottery probability p until $A \sim L_p$



Utility scales

Normalized utilities: $u_\top = 1.0$, $u_\bot = 0.0$

Micromorts: one-millionth chance of death
useful for Russian roulette, paying to reduce product risks, etc.

QALYs: quality-adjusted life years
useful for medical decisions involving substantial risk

Note: behavior is invariant w.r.t. +ve linear transformation

$U'(x) = k_1 U(x) + k_2$ where $k_1 > 0$

With deterministic prizes only (no lottery choices), only ordinal utility can be determined, i.e., total order on prizes

Money

Money does not behave as a utility function

Given a lottery L with expected monetary value $EMV(L)$,
usually $U(L) < U(EMV(L))$, i.e., people are risk-averse

Utility curve: for what probability p am I indifferent between a fixed prize x and a lottery $[p, \$M;\ (1-p), \$0]$ for large M?

Typical empirical data, extrapolated with risk-prone behavior:



Student group utility

For each x, adjust p until half the class votes for lottery (M=10,000)



Decision networks

Add action nodes and utility nodes to belief networks
to enable rational decision making



Algorithm:
For each value of action node
compute expected value of utility node given action, evidence
Return MEU action

Multiattribute utility

How can we handle utility functions of many variables $X_1 \ldots X_n$?
E.g., what is $U(Deaths, Noise, Cost)$?

How can complex utility functions be assessed from
preference behaviour?

Idea 1: identify conditions under which decisions can be made without complete identification of $U(x_1, \ldots, x_n)$

Idea 2: identify various types of independence in preferences
and derive consequent canonical forms for $U(x_1, \ldots, x_n)$

Strict dominance

Typically define attributes such that U is monotonic in each

Strict dominance: choice B strictly dominates choice A iff
$\forall i \;\; X_i(B) \geq X_i(A)$ (and hence $U(B) \geq U(A)$)



Strict dominance seldom holds in practice

Stochastic dominance



Distribution p_1 stochastically dominates distribution p_2 iff
$\forall t \;\; \int_{-\infty}^{t} p_1(x)dx \leq \int_{-\infty}^{t} p_2(t)dt$

If U is monotonic in x, then A_1 with outcome distribution p_1 stochastically dominates A_2 with outcome distribution p_2:
$\int_{-\infty}^{\infty} p_1(x)U(x)dx \geq \int_{-\infty}^{\infty} p_2(x)U(x)dx$
Multiattribute case: stochastic dominance on all attributes $\Rightarrow$ optimal

Stochastic dominance contd.

Stochastic dominance can often be determined without exact distributions using qualitative reasoning

E.q., construction cost increases with distance from city
S_2 is further from the city than S_1
$\Rightarrow$ S_1 stochastically dominates S_2 on cost

E.g., injury increases with collision speed

Can annotate belief networks with stochastic dominance information:
$X \xrightarrow{+} Y$ (X positively influences Y) means that
For every value $\mathbf{z}$ of Y's other parents $\mathbf{Z}$
$\forall x_1, x_2 \;\; x_1 \geq x_2 \Rightarrow \mathbf{P}(Y|x_1, \mathbf{z})$ stochastically dominates $\mathbf{P}(Y|x_2, \mathbf{z})$

Example: car insurance

Which arcs are positive or negative influences?



Preference structure: Deterministic

X_1 and X_2 preferentially independent of X_3 iff
preference between $\langle x_1, x_2, x_3 \rangle$ and $\langle x_1', x_2', x_3 \rangle$
does not depend on x_3

E.g., $\langle Noise, Cost, Safety \rangle$:
⟨20,000 suffer, \$4.6 billion, 0.06 deaths/mpm⟩ vs.
⟨70,000 suffer, \$4.2 billion, 0.06 deaths/mpm⟩

Theorem (Leontief, 1947): if every pair of attributes is P.I. of its complement, then every subset of attributes is P.I of its complement: mutual P.I..

Theorem (Debreu, 1960): mutual P.I. $\Rightarrow$ $\exists$ additive value function:

$$V(S) = \Sigma_i V_i(X_i(S))$$

Hence assess n single-attribute functions; often a good approximation

Preference structure: Stochastic

Need to consider preferences over lotteries:
$\mathbf{X}$ is utility-independent of $\mathbf{Y}$ iff
preferences over lotteries $\mathbf{X}$ do not depend on $\mathbf{y}$

Mutual U.I.: each subset is U.I of its complement
$\Rightarrow$ $\exists$ multiplicative utility function:
$$U = k_1U_1 + k_2U_2 + k_3U_3 + k_1k_2U_1U_2 + k_2k_3U_2U_3 + k_3k_1U_3U_1 + k_1k_2k_3U_1U_2U_3$$

Routine procedures and software packages for generating preference tests to identify various canonical families of utility functions

Value of information

Idea: compute value of acquiring each possible piece of evidence
Can be done directly from decision network

Example: buying oil drilling rights
  Two blocks A and B, exactly one has oil, worth k
  Prior probabilities 0.5 each, mutually exclusive
  Current price of each block is $k/2$
  Consultant offers accurate survey of A. Fair price?

Solution: compute expected value of information
  = expected value of best action given the information
    minus expected value of best action without information
Survey may say "oil in A" or "no oil in A", prob. 0.5 each
  $= [0.5 \times$ value of "buy A" given "oil in A"
    $+ 0.5 \times$ value of "buy B" given "no oil in A"$]$
    − 0
  $= (0.5 \times k/2) + (0.5 \times k/2) - 0 = k/2$

General formula

Current evidence E, current best action α
Possible action outcomes S_i, potential new evidence E_j

$$EU(\alpha|E) = \max_a \Sigma_i\, U(S_i)\; P(S_i|E,a)$$

Suppose we knew $E_j = e_{jk}$, then we would choose $\alpha_{e_{jk}}$ s.t.

$$EU(\alpha_{e_{jk}}|E, E_j = e_{jk}) = \max_a \Sigma_i\, U(S_i)\; P(S_i|E,a,E_j = e_{jk})$$

E_j is a random variable whose value is *currently* unknown
⇒ must compute expected gain over all possible values:

$$VPI_E(E_j) = \left(\Sigma_k\; P(E_j = e_{jk}|E) EU(\alpha_{e_{jk}}|E, E_j = e_{jk})\right) - EU(\alpha|E)$$

(VPI = value of perfect information)

Properties of VPI

Nonnegative—in *expectation*, not *post hoc*

$$\forall j, E\;\; VPI_E(E_j) \geq 0$$

Nonadditive—consider, e.g., obtaining E_j twice

$$VPI_E(E_j, E_k) \neq VPI_E(E_j) + VPI_E(E_k)$$

Order-independent

$$VPI_E(E_j, E_k) = VPI_E(E_j) + VPI_{E,E_j}(E_k) = VPI_E(E_k) + VPI_{E,E_k}(E_j)$$

Note: when more than one piece of evidence can be gathered,
maximizing VPI for each to select one is not always optimal
⇒ evidence-gathering becomes a sequential decision problem

Qualitative behaviors

a) Choice is obvious, information worth little
b) Choice is nonobvious, information worth a lot
c) Choice is nonobvious, information worth little