

## CS 473: Artificial Intelligence

### Bayes' Nets: Independence

Dieter Fox

[These slides were created by Dan Klein and Pieter Abbeel for CS188 Intro to AI at UC Berkeley. All CS188 materials are available at <http://ai.berkeley.edu>.]

## Recap: Bayes' Nets

- A Bayes' net is an efficient encoding of a probabilistic model of a domain

- Questions we can ask:
  - Inference: given a fixed BN, what is  $P(X | e)$ ?
  - Representation: given a BN graph, what kinds of distributions can it encode?
  - Modeling: what BN is most appropriate for a given domain?

## Size of a Bayes' Net

- How big is a joint distribution over  $N$  Boolean variables?  
 $2^N$
- How big is an  $N$ -node net if nodes have up to  $k$  parents?  
 $O(N * 2^{k+1})$

*zomp!*

- Both give you the power to calculate  $P(X_1, X_2, \dots, X_n)$
- BNs: Huge space savings!
- Also easier to elicit local CPTs
- Also faster to answer queries (coming)

## Bayes' Nets

- ✓ Representation
- Conditional Independences
- Probabilistic Inference
- Learning Bayes' Nets from Data

## Conditional Independence

- $X$  and  $Y$  are **independent** if  
 $\forall x, y \ P(x, y) = P(x)P(y) \ \dashrightarrow \ X \perp\!\!\!\perp Y$
- $X$  and  $Y$  are **conditionally independent** given  $Z$   
 $\forall x, y, z \ P(x, y|z) = P(x|z)P(y|z) \ \dashrightarrow \ X \perp\!\!\!\perp Y|Z$
- (Conditional) independence is a property of a distribution
- Example:  $Alarm \perp\!\!\!\perp Fire | Smoke$

## Bayes Nets: Assumptions

- Assumptions we are required to make to define the Bayes net when given the graph:  
 $P(x_i | x_1 \dots x_{i-1}) = P(x_i | \text{parents}(X_i))$
- Beyond above "chain rule  $\rightarrow$  Bayes net" conditional independence assumptions
  - Often additional conditional independences
  - They can be read off the graph
- Important for modeling: understand assumptions made when choosing a Bayes net graph

### Independence in a BN

- Important question about a BN:
  - Are two nodes independent given certain evidence?
  - If yes, can prove using algebra (tedious in general)
  - If no, can prove with a counter example
- Example:
- Question: are X and Z necessarily independent?
  - Answer: no. Example: low pressure causes rain, which causes traffic.
  - X can influence Z, Z can influence X (via Y)
  - Addendum: they *could* be independent: how?

### D-separation: Outline

### D-separation: Outline

- Study independence properties for triples
- Analyze complex cases in terms of member triples
- D-separation: a condition / algorithm for answering such queries

### Causal Chains

- This configuration is a "causal chain"
- Guaranteed X independent of Z? **No!**
  - One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.
- Example:
  - Low pressure causes rain causes traffic, high pressure causes no rain causes no traffic
- In numbers:
 
$$P(+y | +x) = 1, P(-y | -x) = 1, P(+z | +y) = 1, P(-z | -y) = 1$$

X: Low pressure    Y: Rain    Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

### Causal Chains

- This configuration is a "causal chain"
- Guaranteed X independent of Z given Y?
  - Yes!**
  - Evidence along the chain "blocks" the influence

X: Low pressure    Y: Rain    Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

$$P(z|x, y) = \frac{P(x, y, z)}{P(x, y)} = \frac{P(x)P(y|x)P(z|y)}{P(x)P(y|x)} = P(z|y)$$

### Common Cause

- This configuration is a "common cause"
- Guaranteed X independent of Z? **No!**
  - One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.
- Example:
  - Project due causes both forums busy and lab full
- In numbers:
 
$$P(+x | +y) = 1, P(-x | -y) = 1, P(+z | +y) = 1, P(-z | -y) = 1$$

Y: Project due    X: Forums busy    Z: Lab full

$$P(x, y, z) = P(y)P(x|y)P(z|y)$$

### Common Cause

- This configuration is a "common cause"
- Guaranteed X and Z independent given Y?

Y: Project due

X: Forums busy

Z: Lab full

$P(x, y, z) = P(y)P(x|y)P(z|y)$

$$P(z|x, y) = \frac{P(x, y, z)}{P(x, y)}$$

$$= \frac{P(y)P(x|y)P(z|y)}{P(y)P(x|y)}$$

$$= P(z|y)$$

Yes!

Observing the cause blocks influence between effects.

### Common Effect

- Last configuration: two causes of one effect (v-structures)
- Are X and Y independent?
- Are X and Y independent given Z?

X: Raining

Y: Ballgame

Z: Traffic

- Yes: the ballgame and the rain cause traffic, but they are not correlated
- Still need to prove they must be (try it!)
- No: seeing traffic puts the rain and the ballgame in competition as explanation.
- This is backwards from the other cases
- Observing an effect activates influence between possible causes.

### The General Case

### The General Case

- General question: in a given BN, are two variables independent (given evidence)?
- Solution: analyze the graph
- Any complex example can be broken into repetitions of the three canonical cases

### Reachability

- Recipe: shade evidence nodes, look for paths in the resulting graph
- Attempt 1: if two nodes are connected by an undirected path not blocked by a shaded node, they are conditionally independent
- Almost works, but not quite
  - Where does it break?
  - Answer: the v-structure at T doesn't count as a link in a path unless "active"

### Active / Inactive Paths

- Question: Are X and Y conditionally independent given evidence variables [Z]?
- Yes, if X and Y "d-separated" by Z
- Consider all (undirected) paths from X to Y
- No active paths = independence!
- A path is active if each triple is active:
  - Causal chain  $A \rightarrow B \rightarrow C$  where B is unobserved (either direction)
  - Common cause  $A \leftarrow B \rightarrow C$  where B is unobserved
  - Common effect (aka v-structure)  $A \rightarrow B \leftarrow C$  where B or one of its descendants is observed
- All it takes to block a path is a single inactive segment


Active Triples

Inactive Triples

### D-Separation

- Query:  $X_i \perp\!\!\!\perp X_j | \{X_{k_1}, \dots, X_{k_n}\}$  ?
- Check all (undirected!) paths between  $X_i$  and  $X_j$ 
  - If one or more active, then independence not guaranteed
- Otherwise (i.e. if all paths are inactive), then independence is guaranteed

$X_i \not\perp\!\!\!\perp X_j | \{X_{k_1}, \dots, X_{k_n}\}$




$X_i \perp\!\!\!\perp X_j | \{X_{k_1}, \dots, X_{k_n}\}$

### Example

$R \perp\!\!\!\perp B$      **Yes**

$R \perp\!\!\!\perp B | T$

$R \perp\!\!\!\perp B | T'$



### Example


$L \perp\!\!\!\perp T' | T$      **Yes**

$L \perp\!\!\!\perp B$      **Yes**

$L \perp\!\!\!\perp B | T$

$L \perp\!\!\!\perp B | T'$

$L \perp\!\!\!\perp B | T, R$      **Yes**



### Example

**Variables:**


- R: Raining
- T: Traffic
- D: Roof drips
- S: I'm sad

**Questions:**

$T \perp\!\!\!\perp D$

$T \perp\!\!\!\perp D | R$      **Yes**


$T \perp\!\!\!\perp D | R, S$



### Structure Implications

- Given a Bayes net structure, can run d-separation algorithm to build a complete list of conditional independences that are necessarily true of the form


$X_i \perp\!\!\!\perp X_j | \{X_{k_1}, \dots, X_{k_n}\}$




- This list determines the set of probability distributions that can be represented

### Computing All Independences


**COMPUTE ALL THE INDEPENDENCES!**





## Topology Limits Distributions

- Given some graph topology  $G$ , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences
- (There might be more independence)
- Adding arcs increases the set of distributions, but has several costs
- Full conditioning can encode any distribution



## Bayes Nets Representation Summary

- Bayes nets compactly encode joint distributions
- Guaranteed independencies of distributions can be deduced from BN graph structure
- D-separation gives precise conditional independence guarantees from graph alone
- A Bayes' net's joint distribution may have further (conditional) independence that is not detectable until you inspect its specific distribution

## Bayes' Nets

- ✓ Representation
- ✓ Conditional Independences
  - Probabilistic Inference
    - Enumeration (exact, exponential complexity)
    - Variable elimination (exact, worst-case exponential complexity, often better)
    - Probabilistic inference is NP-complete
    - Sampling (approximate)
  - Learning Bayes' Nets from Data