


I COULD RESTRUCTURE  
THE PROGRAM'S FLOW  
OR USE ONE LITTLE  
'GOTO' INSTEAD.



EH, SCREW GOOD PRACTICE.  
HOW BAD CAN IT BE?


```
goto main_sub3;
```

\*COMPILE\*



# CS 473: Artificial Intelligence

## Bayes' Nets: Independence




Steve Tanimoto

[These slides were created by Dan Klein and Pieter Abbeel for CS188 Intro to AI at UC Berkeley. All CS188 materials are available at <http://ai.berkeley.edu>.]

# Recap: Bayes' Nets

- A Bayes' net is an efficient encoding of a probabilistic model of a domain



- Questions we can ask:

- Inference: given a fixed BN, what is  $P(X \mid e)$ ?
- Representation: given a BN graph, what kinds of distributions can it encode?
- Modeling: what BN is most appropriate for a given domain?

# Bayes' Nets

---

- ✓ Representation
  - Conditional Independences
  - Probabilistic Inference
  - Learning Bayes' Nets from Data

# Conditional Independence

- X and Y are **independent** if

$$\forall x, y \quad P(x, y) = P(x)P(y) \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y$$

- X and Y are **conditionally independent** given Z

$$\forall x, y, z \quad P(x, y|z) = P(x|z)P(y|z) \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y|Z$$

- (Conditional) independence is a property of a distribution

- Example:  $Alarm \perp\!\!\!\perp Fire|Smoke$



# Bayes Nets: Assumptions

- Assumptions we are required to make to define the Bayes net when given the graph:


$$P(x_i | x_1 \cdots x_{i-1}) = P(x_i | \text{parents}(X_i))$$

- Beyond above “chain rule  $\rightarrow$  Bayes net” conditional independence assumptions
  - Often additional conditional independences
  - They can be read off the graph
- Important for modeling: understand assumptions made when choosing a Bayes net graph




# Independence in a BN

- Important question about a BN:
  - Are two nodes independent given certain evidence?
  - If yes, can prove using algebra (tedious in general)
  - If no, can prove with a counter example
  - Example:



- Question: are X and Z necessarily independent?
  - Answer: no. Example: low pressure causes rain, which causes traffic.
  - X can influence Z, Z can influence X (via Y)
  - Addendum: they *could* be independent: how?

# D-separation: Outline




# D-separation: Outline

---

- Study independence properties for triples
- Analyze complex cases in terms of member triples
- D-separation: a condition / algorithm for answering such queries

# Causal Chains

- This configuration is a “causal chain”



$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Guaranteed X independent of Z ? *No!*

- One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

- Example:


- Low pressure causes rain causes traffic, high pressure causes no rain causes no traffic

- In numbers:

$$P(+y | +x) = 1, P(-y | -x) = 1, \\ P(+z | +y) = 1, P(-z | -y) = 1$$

# Causal Chains

- This configuration is a “causal chain”



- Guaranteed X independent of Z given Y?

$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} \\ &= \frac{P(x)P(y|x)P(z|y)}{P(x)P(y|x)} \\ &= P(z|y) \end{aligned}$$


*Yes!*

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Evidence along the chain “blocks” the influence

# Common Cause

- This configuration is a “common cause”



$$P(x, y, z) = P(y)P(x|y)P(z|y)$$

- Guaranteed X independent of Z ? *No!*

- One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

- Example:

- Project due causes both forums busy and lab full


- In numbers:

$$P(+x | +y) = 1, P(-x | -y) = 1,$$
$$P(+z | +y) = 1, P(-z | -y) = 1$$

# Common Cause

- This configuration is a “common cause”

- Guaranteed X and Z independent given Y?



$$P(x, y, z) = P(y)P(x|y)P(z|y)$$


$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} \\ &= \frac{P(y)P(x|y)P(z|y)}{P(y)P(x|y)} \\ &= P(z|y) \end{aligned}$$

*Yes!*

- Observing the cause blocks influence between effects.

# Common Effect


- Last configuration: two causes of one effect (v-structures)



- Are X and Y independent?
  - **Yes**: the ballgame and the rain cause traffic, but they are not correlated
  - Still need to prove they must be (try it!)
- Are X and Y independent given Z?
  - **No**: seeing traffic puts the rain and the ballgame in competition as explanation.
- **This is backwards from the other cases**
  - Observing an effect **activates** influence between possible causes.


# The General Case

---




# The General Case

- General question: in a given BN, are two variables independent (given evidence)?
- Solution: analyze the graph
- Any complex example can be broken into repetitions of the three canonical cases



# Reachability

- Recipe: shade evidence nodes, look for paths in the resulting graph
- Attempt 1: if two nodes are connected by an undirected path not blocked by a shaded node, then they are not conditionally independent
- Almost works, but not quite
  - Where does it break?
  - Answer: the v-structure at T doesn't count as a link in a path unless "active"



# Active / Inactive Paths

- Question: Are X and Y conditionally independent given evidence variables {Z}?





- Yes, if X and Y “d-separated” by Z
- Consider all (undirected) paths from X to Y
- No active paths = independence!

- A path is active if each triple is active:



- Causal chain  $A \rightarrow B \rightarrow C$  where B is unobserved (either direction)
- Common cause  $A \leftarrow B \rightarrow C$  where B is unobserved
- Common effect (aka v-structure)  
 $A \rightarrow B \leftarrow C$  where B or one of its descendants is observed

- All it takes to block a path is a single inactive segment

Active Triples



Inactive Triples




# D-Separation

- Query:  $X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\} ?$
- Check all (undirected!) paths between  $X_i$  and  $X_j$ 
  - If one or more active, then independence not guaranteed

$$X_i \not\perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$

- Otherwise (i.e. if all paths are inactive), then independence is guaranteed

$$X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$




# Example

---

$R \perp\!\!\!\perp B$       *Yes*

$R \perp\!\!\!\perp B | T$

$R \perp\!\!\!\perp B | T'$



# Example

---


$L \perp\!\!\!\perp T' | T$       *Yes*

$L \perp\!\!\!\perp B$       *Yes*

$L \perp\!\!\!\perp B | T$

$L \perp\!\!\!\perp B | T'$

$L \perp\!\!\!\perp B | T, R$       *Yes*



# Example


- Variables:
  - R: Raining
  - T: Traffic
  - D: Roof drips
  - S: I'm sad

- Questions:

$$T \perp\!\!\!\perp D$$

$$T \perp\!\!\!\perp D | R \quad \text{Yes}$$

$$T \perp\!\!\!\perp D | R, S$$




# Structure Implications

- Given a Bayes net structure, can run d-separation algorithm to build a complete list of conditional independences that are necessarily true of the form


$$X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$

- This list determines the set of probability distributions that can be represented



# Computing All Independences



COMPUTE ALL THE  
INDEPENDENCES!




# Topology Limits Distributions

- Given some graph topology  $G$ , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences
- (There might be more independence)
- Adding arcs increases the set of distributions, but has several costs
- Full conditioning can encode any distribution


$\{X \perp\!\!\!\perp Y, X \perp\!\!\!\perp Z, Y \perp\!\!\!\perp Z, X \perp\!\!\!\perp Z \mid Y, X \perp\!\!\!\perp Y \mid Z, Y \perp\!\!\!\perp Z \mid X\}$



$\{X \perp\!\!\!\perp Z \mid Y\}$



$\{\}$



# Bayes Nets Representation Summary

---

- Bayes nets compactly encode joint distributions
- Guaranteed independencies of distributions can be deduced from BN graph structure
- D-separation gives precise conditional independence guarantees from graph alone
- A Bayes' net's joint distribution may have further (conditional) independence that is not detectable until you inspect its specific distribution

# Bayes' Nets

---

✓ Representation

✓ Conditional Independences

- Probabilistic Inference

- Enumeration (exact, exponential complexity)

- Variable elimination (exact, worst-case exponential complexity, often better)

- Probabilistic inference is NP-complete

- Sampling (approximate)

- Learning Bayes' Nets from Data