

CS 473: Artificial Intelligence

Bayes' Nets: Independence

Steve Tanimoto

[These slides were created by Dan Klein and Pieter Abbeel for CS188 Intro to AI at UC Berkeley. All CS188 materials are available at <http://ai.berkeley.edu>.]

Recap: Bayes' Nets

- A Bayes' net is an efficient encoding of a probabilistic model of a domain

- Questions we can ask:
 - Inference: given a fixed BN, what is $P(X | e)$?
 - Representation: given a BN graph, what kinds of distributions can it encode?
 - Modeling: what BN is most appropriate for a given domain?

Bayes' Nets

- ✓ Representation
 - Conditional Independences
 - Probabilistic Inference
 - Learning Bayes' Nets from Data

Conditional Independence

- X and Y are **independent** if

$$\forall x, y \quad P(x, y) = P(x)P(y) \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y$$
- X and Y are **conditionally independent** given Z

$$\forall x, y, z \quad P(x, y|z) = P(x|z)P(y|z) \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y | Z$$
- (Conditional) independence is a property of a distribution
- Example: $Alarm \perp\!\!\!\perp Fire | Smoke$


Bayes Nets: Assumptions

- Assumptions we are required to make to define the Bayes net when given the graph:

$$P(x_i | x_1 \dots x_{i-1}) = P(x_i | \text{parents}(X_i))$$
- Beyond above "chain rule \rightarrow Bayes net" conditional independence assumptions
 - Often additional conditional independences
 - They can be read off the graph
- Important for modeling: understand assumptions made when choosing a Bayes net graph


Independence in a BN

- Important question about a BN:
 - Are two nodes independent given certain evidence?
 - If yes, can prove using algebra (tedious in general)
 - If no, can prove with a counter example
 - Example:



- Question: are X and Z necessarily independent?
 - Answer: no. Example: low pressure causes rain, which causes traffic.
 - X can influence Z, Z can influence X (via Y)
 - Addendum: they *could* be independent: how?

D-separation: Outline




D-separation: Outline

- Study independence properties for triples
- Analyze complex cases in terms of member triples
- D-separation: a condition / algorithm for answering such queries

Causal Chains

- This configuration is a "causal chain"
- Guaranteed X independent of Z? **No!**



X: Low pressure Y: Rain Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

Example:


- Low pressure causes rain causes traffic, high pressure causes no rain causes no traffic

In numbers:

$$P(+y | +x) = 1, P(-y | -x) = 1, \\ P(+z | +y) = 1, P(-z | -y) = 1$$

Causal Chains

- This configuration is a "causal chain"
- Guaranteed X independent of Z given Y?



X: Low pressure Y: Rain Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$


$$P(z|x, y) = \frac{P(x, y, z)}{P(x, y)} \\ = \frac{P(x)P(y|x)P(z|y)}{P(x)P(y|x)} \\ = P(z|y)$$

Yes!

- Evidence along the chain "blocks" the influence

Common Cause

- This configuration is a "common cause"
- Guaranteed X independent of Z? **No!**



X: Forums busy Y: Project due Z: Lab full

$$P(x, y, z) = P(y)P(x|y)P(z|y)$$

- One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

Example:


- Project due causes both forums busy and lab full

In numbers:

$$P(+x | +y) = 1, P(-x | -y) = 1, \\ P(+z | +y) = 1, P(-z | -y) = 1$$

Common Cause

- This configuration is a "common cause"
- Guaranteed X and Z independent given Y?



$$P(z|x, y) = \frac{P(x, y, z)}{P(x, y)}$$

$$= \frac{P(y)P(x|y)P(z|y)}{P(y)P(x|y)}$$


$$= P(z|y)$$

Yes!

- Observing the cause blocks influence between effects.


Common Effect

- Last configuration: two causes of one effect (v-structures)
- Are X and Y independent?




- Are X and Y independent?
 - Yes: the ballgame and the rain cause traffic, but they are not correlated
 - Still need to prove they must be (try it!)
- Are X and Y independent given Z?
 - No: seeing traffic puts the rain and the ballgame in competition as explanation.
- This is backwards from the other cases
 - Observing an effect **activates** influence between possible causes.

The General Case




The General Case

- General question: in a given BN, are two variables independent (given evidence)?
- Solution: analyze the graph
- Any complex example can be broken into repetitions of the three canonical cases



Reachability


- Recipe: shade evidence nodes, look for paths in the resulting graph
- Attempt 1: if two nodes are connected by an undirected path not blocked by a shaded node, then they are not conditionally independent
- Almost works, but not quite
 - Where does it break?
 - Answer: the v-structure at T doesn't count as a link in a path unless "active"



Active / Inactive Paths

- Question: Are X and Y conditionally independent given evidence variables {Z}?

- Yes, if X and Y "d-separated" by Z
- Consider all (undirected) paths from X to Y
- No active paths = independence!



- A path is active if each triple is active:
 - Causal chain $A \rightarrow B \rightarrow C$ where B is unobserved (either direction)
 - Common cause $A \leftarrow B \rightarrow C$ where B is unobserved
 - Common effect (aka v-structure) $A \rightarrow B \leftarrow C$ where B or one of its descendants is observed
- All it takes to block a path is a single inactive segment


D-Separation

- Query: $X_i \perp\!\!\!\perp X_j | \{X_{k_1}, \dots, X_{k_n}\}$?
- Check all (undirected!) paths between X_i and X_j
 - If one or more active, then independence not guaranteed

$$X_i \not\perp\!\!\!\perp X_j | \{X_{k_1}, \dots, X_{k_n}\}$$


- Otherwise (i.e. if all paths are inactive), then independence is guaranteed

$$X_i \perp\!\!\!\perp X_j | \{X_{k_1}, \dots, X_{k_n}\}$$




Example

$R \perp\!\!\!\perp B$ Yes
 $R \perp\!\!\!\perp B | T$
 $R \perp\!\!\!\perp B | T'$



Example

$L \perp\!\!\!\perp T' | T$ Yes
 $L \perp\!\!\!\perp B$ Yes
 $L \perp\!\!\!\perp B | T$
 $L \perp\!\!\!\perp B | T'$
 $L \perp\!\!\!\perp B | T, R$ Yes




Example

- Variables:
 - R: Raining
 - T: Traffic
 - D: Roof drips
 - S: I'm sad

- Questions:

$T \perp\!\!\!\perp D$
 $T \perp\!\!\!\perp D | R$ Yes
 $T \perp\!\!\!\perp D | R, S$




Structure Implications

- Given a Bayes net structure, can run d-separation algorithm to build a complete list of conditional independences that are necessarily true of the form



$$X_i \perp\!\!\!\perp X_j | \{X_{k_1}, \dots, X_{k_n}\}$$

- This list determines the set of probability distributions that can be represented




Computing All Independences

COMPUTE ALL THE INDEPENDENCES!



Topology Limits Distributions

- Given some graph topology G , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences
- (There might be more independence)
- Adding arcs increases the set of distributions, but has several costs
- Full conditioning can encode any distribution



Bayes Nets Representation Summary

- Bayes nets compactly encode joint distributions
- Guaranteed independencies of distributions can be deduced from BN graph structure
- D-separation gives precise conditional independence guarantees from graph alone
- A Bayes' net's joint distribution may have further (conditional) independence that is not detectable until you inspect its specific distribution

Bayes' Nets

- ✓ Representation
- ✓ Conditional Independences
 - Probabilistic Inference
 - Enumeration (exact, exponential complexity)
 - Variable elimination (exact, worst-case exponential complexity, often better)
 - Probabilistic inference is NP-complete
 - Sampling (approximate)
 - Learning Bayes' Nets from Data