


CSE 473: Artificial Intelligence

Bayes' Nets: Inference



Steve Tanimoto

[These slides were created by Dan Klein and Pieter Abbeel for CS188 Intro to AI at UC Berkeley. All CS188 materials are available at <http://ai.berkeley.edu>.]

Inference

▪ Inference: calculating some useful quantity from a joint probability distribution

▪ Examples:

▪ Posterior probability


$$P(Q|E_1 = e_1, \dots, E_k = e_k)$$

▪ Most likely explanation:

$$\text{argmax}_q P(Q = q|E_1 = e_1 \dots)$$



Test for Infant Metabolic Defects



Blue ovals represent chromatographic peaks, grey ovals represent 20 metabolic diseases

Inference by Enumeration

▪ General case:

▪ Evidence variables: $E_1 \dots E_k = e_1 \dots e_k$
 ▪ Query* variable: Q
 ▪ Hidden variables: $H_1 \dots H_r$

▪ We want:


$$P(Q|e_1 \dots e_k)$$

* Works fine with multiple query variables, too

▪ Step 1: Select the entries consistent with the evidence

x	prob
-3	0.08
-1	0.25
0	0.07
1	0.2
5	0.01

▪ Step 2: Sum out H to get joint of Query and evidence



▪ Step 3: Normalize

$$\times \frac{1}{Z}$$

$$Z = \sum_q P(Q, e_1 \dots e_k)$$

$$P(Q|e_1 \dots e_k) = \frac{1}{Z} P(Q, e_1 \dots e_k)$$

$$P(Q, e_1 \dots e_k) = \sum_{h_1 \dots h_r} P(Q, h_1 \dots h_r, e_1 \dots e_k)$$

Inference by Enumeration in Bayes' Net


- Given unlimited time, inference in BNs is easy
- Reminder of inference by enumeration by example:

$$P(B | +j, +m) \propto_B P(B, +j, +m)$$

$$= \sum_{e,a} P(B, e, a, +j, +m)$$

$$= \sum_{e,a} P(B)P(e)P(a|B, e)P(+j|a)P(+m|a)$$


$$= P(B)P(+e)P(+a|B, +e)P(+j|+a)P(+m|+a) + P(B)P(+e)P(-a|B, +e)P(+j|-a)P(+m|-a) + P(B)P(-e)P(+a|B, -e)P(+j|+a)P(+m|+a) + P(B)P(-e)P(-a|B, -e)P(+j|-a)P(+m|-a)$$



Inference by Enumeration vs. Variable Elimination


▪ Why is inference by enumeration so slow?

- You join up the whole joint distribution before you sum out the hidden variables



▪ Idea: interleave joining and marginalizing!

- Called "Variable Elimination"
- Still NP-hard, but usually much faster than inference by enumeration



▪ First we'll need some new notation: factors

Traffic Domain

$$\begin{matrix} \textcircled{R} \\ \downarrow \\ \textcircled{T} \\ \downarrow \\ \textcircled{L} \end{matrix}$$

$P(L) = ?$

- Inference by Enumeration
- Variable Elimination

$$= \sum_t \sum_r P(L|t) P(r) P(t|r)$$

Join on r

Join on t

Eliminate r

Eliminate t


$$= \sum_t P(L|t) \sum_r P(r) P(t|r)$$

Join on r

Eliminate r

Join on t

Eliminate t



Factor Zoo I

- Joint distribution: $P(X, Y)$
 - Entries $P(x, y)$ for all x, y
 - Sums to 1
- Selected joint: $P(x, Y)$
 - A slice of the joint distribution
 - Entries $P(x, y)$ for fixed x , all y
 - Sums to $P(x)$
- Number of capitals = dimensionality of the table

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

T	W	P
cold	sun	0.2
cold	rain	0.3

Factor Zoo I

Two dimensions

	sun	rain
hot	0.4	0.1
cold	0.2	0.3

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

One dimension

	sun	rain
cold	0.2	0.3

T	W	P
cold	sun	0.2
cold	rain	0.3

- Number of capitals = dimensionality of the table

Factor Zoo II

- Single conditional: $P(Y | x)$
 - Entries $P(y | x)$ for fixed x , all y
 - Sums to 1
- Family of conditionals: $P(X | Y)$
 - Multiple conditionals
 - Entries $P(x | y)$ for all x, y
 - Sums to $|Y|$

T	W	P
cold	sun	0.4
cold	rain	0.6

T	W	P
hot	sun	0.8
hot	rain	0.2
cold	sun	0.4
cold	rain	0.6

$\left. \begin{matrix} P(W|hot) \\ P(W|cold) \end{matrix} \right\}$

Factor Zoo III


- Specified family: $P(y | X)$
 - Entries $P(y | x)$ for fixed y , but for all x
 - Sums to ... who knows!

T	W	P
hot	rain	0.2
cold	rain	0.6

$\left. \begin{matrix} P(rain|hot) \\ P(rain|cold) \end{matrix} \right\}$

Factor Zoo Summary

- In general, when we write $P(Y_1 \dots Y_N \mid X_1 \dots X_M)$
 - It is a "factor," a multi-dimensional array
 - Its values are $P(y_1 \dots y_N \mid x_1 \dots x_M)$
 - Any assigned (=lower-case) X or Y is a dimension missing (selected) from the array



Example: Traffic Domain

Random Variables

- R: Raining
- T: Traffic
- L: Late for class!



$$P(R)$$

+r	0.1
-r	0.9

$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$$P(L|T)$$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

$$P(L) = ?$$

$$= \sum_{r,t} P(r,t,L)$$

$$= \sum_{r,t} P(r)P(t|r)P(L|t)$$

Inference by Enumeration: Procedural Outline

- Track objects called **factors**
- Initial factors are local CPTs (one per node)
- Any known values are selected
 - E.g. if we know $L = +l$, the initial factors are

$$P(R)$$

+r	0.1
-r	0.9

$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$$P(L|T)$$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

$$P(R)$$


+r	0.1
-r	0.9

$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$$P(+l|T)$$



+t	+l	0.3
+t	-l	0.1



- Procedure: Join all factors, then eliminate all hidden variables


Operation 1: Join Factors

- First basic operation: **joining factors**
- Combining factors:
 - Just like a database join
 - Get all factors over the joining variable
 - Build a new factor over the union of the variables involved
- Example: Join on R




- Computation for each entry: pointwise products $\forall r, t: P(r, t) = P(r) \cdot P(t|r)$

Example: Multiple Joins



Example: Multiple Joins



$$P(R)$$

+r	0.1
-r	0.9

$$P(T|R)$$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$$P(L|T)$$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

$$P(R, T)$$

+r	+t	0.08
+r	-t	0.02
-r	+t	0.09
-r	-t	0.81

$$P(R, T, L)$$

+r	+t	+l	0.024
+r	+t	-l	0.056
+r	-t	+l	0.002
+r	-t	-l	0.018
-r	+t	+l	0.027
-r	+t	-l	0.063
-r	-t	+l	0.081
-r	-t	-l	0.729

Operation 2: Eliminate

- Second basic operation: **marginalization**
- Take a factor and sum out a variable
 - Shrinks a factor to a smaller one
 - An **aggregate/project** operation
- Example:

$P(R, T)$

+r	+t	0.08
+r	-t	0.02
-r	+t	0.09
-r	-t	0.81

sum R

$P(T)$

+t	0.17
-t	0.83

Multiple Elimination

$P(R, T, L)$

+r	+t	+l	0.024
+r	+t	-l	0.056
+r	-t	+l	0.002
+r	-t	-l	0.018
-r	+t	+l	0.027
-r	+t	-l	0.063
-r	-t	+l	0.081
-r	-t	-l	0.729

Sum out R

$P(T, L)$

+t	+l	0.051
+t	-l	0.119
-t	+l	0.083
-t	-l	0.747

Sum out T

$P(L)$

+l	0.134
-l	0.866

Thus Far: Multiple Join, Multiple Eliminate (= Inference by Enumeration)

Marginalizing Early (= Variable Elimination)

Traffic Domain

$P(L) = ?$

Inference by Enumeration

$$= \sum_t \sum_r P(L|t) P(r) P(t|r)$$

Variable Elimination

$$= \sum_t P(L|t) \sum_r P(r) P(t|r)$$

Marginalizing Early! (aka VE)

$P(R)$

+r	0.1
-r	0.9

Join R

$P(R, T)$

+r	+t	0.08
+r	-t	0.02
-r	+t	0.09
-r	-t	0.81

Sum out R

$P(T)$

+t	0.17
-t	0.83

Join T

$P(T, L)$

+t	+l	0.051
+t	-l	0.119
-t	+l	0.083
-t	-l	0.747

Sum out T

$P(L)$

+l	0.134
-l	0.866

Evidence

- If evidence, start with factors that select that evidence

- If there is no evidence, then use these initial factors:


$P(R)$	$P(T R)$	$P(L T)$
+r 0.1	+r +t 0.8	+t +l 0.3
-r 0.9	+r -t 0.2	+t -l 0.7
	-r +t 0.1	-t +l 0.1
	-r -t 0.9	-t -l 0.9

- But if given some evidence, eg +r, then select for it...

- Computing $P(L|+r)$ the initial factors become:

$P(+r)$	$P(T +r)$	$P(L T)$
+r 0.1	+r +t 0.8	+t +l 0.3
	+r -t 0.2	+t -l 0.7
	-r +t 0.1	-t +l 0.1
	-r -t 0.9	-t -l 0.9

- Next do joins & eliminate, removing all vars other than query + evidence



Evidence II


- Result will be a selected joint of query and evidence

- E.g. for $P(L|+r)$, we would end up with:

$P(+r, L)$	Normalize	$P(L +r)$
+r +l 0.026	→	+l 0.26
+r -l 0.074		-l 0.74

- To get our answer, just normalize this!

- That's it!



General Variable Elimination



- Query: $P(Q|E_1 = e_1, \dots, E_k = e_k)$

- Start with initial factors:
 - Local CPTs (but instantiated by evidence)

- While there are still hidden variables (not Q or evidence):

- Choose a hidden variable H
- Join all factors mentioning H
- Eliminate (sum out) H

- Join all remaining factors and normalize




$$\text{Factors} \times \frac{1}{Z}$$

Example: Alarm Network

$$P(B|j, m) = ?$$

B	P(B)
+b	0.001
-b	0.999



E	P(E)
+e	0.002
-e	0.998



A	J	P(J A)
+a	+j	0.9
+a	-j	0.1
-a	+j	0.05
-a	-j	0.95


A	M	P(M A)
+a	+m	0.7
+a	-m	0.3
-a	+m	0.01
-a	-m	0.99

B	E	A	P(A B,E)
+b	+e	+a	0.95
+b	+e	-a	0.05
+b	-e	+a	0.94
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999

Example

$$P(B|j, m) \propto P(B, j, m)$$

$P(B)$	$P(E)$	$P(A B, E)$	$P(j A)$	$P(m A)$
--------	--------	-------------	----------	----------



Choose A

$$P(A|B, E) P(j|A) P(m|A) \xrightarrow{\times} P(j, m, A|B, E) \xrightarrow{\sum} P(j, m|B, E)$$

$P(B)$	$P(E)$	$P(j, m B, E)$
--------	--------	----------------

Example

$P(B)$	$P(E)$	$P(j, m B, E)$
--------	--------	----------------


Choose E

$$P(E) P(j, m|B, E) \xrightarrow{\times} P(j, m, E|B) \xrightarrow{\sum} P(j, m|B)$$

$P(B)$	$P(j, m B)$
--------	-------------

Finish with B

$$P(B) P(j, m|B) \xrightarrow{\times} P(j, m, B) \xrightarrow{\text{Normalize}} P(B|j, m)$$



Same Example in Equations

$$P(B|j, m) \propto P(B, j, m)$$

$P(B)$	$P(E)$	$P(A B, E)$	$P(j A)$	$P(m A)$
--------	--------	-------------	----------	----------

$$P(B|j, m) \propto \sum_{e,a} P(B, j, m, e, a)$$

$$= \sum_{e,a} P(B)P(e)P(a|B, e)P(j|a)P(m|a)$$

use Bayes' net joint distribution expression

$$= \sum_{e,a} P(B)P(e) \sum_a P(a|B, e)P(j|a)P(m|a)$$

use $xy + xz = x*(y+z)$ **do sum first**

$$= \sum_e P(B)P(e) f_1(B, e, j, m)$$

joining on a, and then summing out gives f_1

$$= P(B) \sum_e P(e) f_1(B, e, j, m)$$

use $xy + xz = x*(y+z)$ **do sum first**

$$= P(B) f_2(B, j, m)$$

joining on e, and then summing out gives f_2

Simple! Exploiting $uwy + uwz + uxy + uxz + vwy + vwz + vxy + vxz = (u+v)(w+x)(y+z)$ to reduce computation

Variable Elimination

$$P(b|j, m) = \sum_e P(b) \sum_a P(e) \sum_a P(a|b, e) P(j|a) P(m, a)$$

Repeated computations \rightarrow Dynamic Programming

Choices during Variable Elimination

- Query: $P(Q|E_1 = e_1, \dots, E_k = e_k)$
- Start with initial factors:
 - Local CPTs (but instantiated by evidence)
- While there are still hidden variables (not Q or evidence):
 - **Choose a hidden variable H**
 - Join all factors mentioning H
 - Eliminate (sum out) H
- Join all remaining factors and normalize

Another Variable Elimination Example

Query: $P(X_3|Y_1 = y_1, Y_2 = y_2, Y_3 = y_3)$

Start by inserting evidence, which gives the following initial factors:

$$p(Z)p(X_1|Z)p(X_2|Z)p(X_3|Z)p(y_1|X_1)p(y_2|X_2)p(y_3|X_3)$$

What variables could we eliminate?

Another Variable Elimination Example

Query: $P(X_3|Y_1 = y_1, Y_2 = y_2, Y_3 = y_3)$

Start by inserting evidence, which gives the following initial factors:

$$p(Z)p(X_1|Z)p(X_2|Z)p(X_3|Z)p(y_1|X_1)p(y_2|X_2)p(y_3|X_3)$$

Eliminate X_1 , this introduces the factor $f_1(Z, y_1) = \sum_{x_1} p(x_1|Z)p(y_1|x_1)$, and we are left with:

$$p(Z) f_1(Z, y_1) p(X_2|Z) p(X_3|Z) p(y_2|X_2) p(y_3|X_3)$$

Eliminate X_2 , this introduces the factor $f_2(Z, y_2) = \sum_{x_2} p(x_2|Z)p(y_2|x_2)$, and we are left with:

$$p(Z) f_1(Z, y_1) f_2(Z, y_2) p(X_3|Z) p(y_3|X_3)$$

Eliminate Z, this introduces the factor $f_3(y_1, y_2, y_3) = \sum_z p(z) f_1(z, y_1) f_2(z, y_2) p(X_3|z) p(y_3|X_3)$, and we are left with:

$$p(y_3|X_3) f_3(y_1, y_2, y_3)$$

No hidden variables left. Join the remaining factors to get:

$$f_4(y_1, y_2, y_3, X_3) = P(y_3|X_3) f_3(y_1, y_2, y_3)$$

Normalizing over X_3 gives $P(X_3|y_1, y_2, y_3)$.

1

What dimension are f_1, f_2 & f_3 ?

Another Variable Elimination Example

Query: $P(X_3|Y_1 = y_1, Y_2 = y_2, Y_3 = y_3)$

Start by inserting evidence, which gives the following initial factors:

$$p(Z)p(X_1|Z)p(X_2|Z)p(X_3|Z)p(y_1|X_1)p(y_2|X_2)p(y_3|X_3)$$

Alternatively, suppose we start by eliminating Z:

$P(X_1 | Z)$
 $P(X_2 | Z)$
 $P(X_3 | Z)$

\times

$f_Z(X_1, X_2, X_3)$
 $p(y_1 | X_1)$
 $p(y_2 | X_2)$
 $p(y_3 | X_3)$

What is the resulting factor? 3

What dimension is it? 3

How many entries? k^3

Another Variable Elimination Example

Query: $P(X_3|Y_1 = y_1, Y_2 = y_2, Y_3 = y_3)$

Start by inserting evidence, which gives the following initial factors:

$$p(Z)p(X_1|Z)p(X_2|Z)p(X_3|Z)p(y_1|X_1)p(y_2|X_2)p(y_3|X_3)$$

Eliminate X_1 , this introduces the factor $f_1(Z, y_1) = \sum_{x_1} p(x_1|Z)p(y_1|x_1)$, and we are left with:

$$p(Z)f_1(Z, y_1)p(X_2|Z)p(X_3|Z)p(y_2|X_2)p(y_3|X_3)$$

Eliminate X_2 , this introduces the factor $f_2(Z, y_2) = \sum_{x_2} p(x_2|Z)p(y_2|x_2)$, and we are left with:

$$p(Z)f_1(Z, y_1)f_2(Z, y_2)p(X_3|Z)p(y_3|X_3)$$


Eliminate Z , this introduces the factor $f_3(y_1, y_2, X_3) = \sum_z p(z)f_1(z, y_1)f_2(z, y_2)p(X_3|z)$, and we are left:

$$p(y_3|X_3) \cdot f_3(y_1, y_2, X_3)$$

No hidden variables left. Join the remaining factors to get:

$$f_4(y_1, y_2, y_3, X_3) = P(y_3|X_3)f_3(y_1, y_2, X_3)$$


Normalizing over X_3 gives $P(X_3|y_1, y_2, y_3)$.



Computational complexity depends on the **largest factor** generated by the process.
Size of factor = number of entries in table.

Variable Elimination Ordering

- For the query $P(X_n|Y_1, \dots, Y_n)$ work through the following two different orderings as done in previous slide: $Z, X_{12}, \dots, X_{n-3}$ and $X_{12}, \dots, X_{n-1}, Z$. What is the size of the maximum factor generated for each of the orderings?



- Answer: 2^{n+1} versus 2^2 (assuming binary)
- In general: the ordering can **greatly** affect efficiency.

VE: Computational and Space Complexity

- The computational and space complexity of variable elimination is determined by the largest factor
- The elimination ordering can greatly affect the size of the largest factor.
 - E.g., previous slide's example 2^n vs. 2
- Does there always exist an ordering that only results in small factors?
 - No!**

Worst Case Complexity?

- CSP:

$$(x_1 \vee x_2 \vee \neg x_3) \wedge (\neg x_1 \vee x_3 \vee \neg x_4) \wedge (x_2 \vee \neg x_2 \vee x_4) \wedge (\neg x_3 \vee \neg x_4 \vee \neg x_5) \wedge (x_2 \vee x_5 \vee x_7) \wedge (x_4 \vee x_5 \vee x_6) \wedge (\neg x_5 \vee x_6 \vee \neg x_7) \wedge (\neg x_5 \vee \neg x_6 \vee x_7)$$

$$P(X_1 = 0) = P(X_1 = 1) = 0.5$$

$$Y_1 = X_1 \vee X_2 \vee \neg X_3$$

$$Y_6 = \neg X_5 \vee X_6 \vee X_7$$


$$Y_{1,2} = Y_1 \wedge Y_2$$

$$Y_{7,8} = Y_7 \wedge Y_8$$

$$Y_{1,2,3,4} = Y_{1,2} \wedge Y_{3,4}$$

$$Y_{5,6,7,8} = Y_{5,6} \wedge Y_{7,8}$$

$$Z = Y_{1,2,3,4} \wedge Y_{5,6,7,8}$$



- If we can answer $P(z)$ equal to zero or not, we answered whether the 3-SAT problem has a solution.
- Hence inference in Bayes' nets is NP-hard. No known efficient probabilistic inference in general.

Bayes' Nets

- Representation
- Conditional Independences
- Probabilistic Inference
 - Enumeration (exact, exponential complexity)
 - Variable elimination (exact, worst-case exponential complexity, often better)
 - Inference is NP-complete
 - Sampling (approximate)
- Learning Bayes' Nets from Data