

---

# CSE 473: Introduction to Artificial Intelligence

Hanna Hajishirzi  
Hidden Markov Models

slides adapted from  
Dan Klein, Pieter Abbeel [ai.berkeley.edu](http://ai.berkeley.edu)  
And Dan Weld, Luke Zettlemoyer



# Probability Summary

---

- **Conditional probability**  $P(x|y) = \frac{P(x, y)}{P(y)}$
- **Product rule**  $P(x, y) = P(x|y)P(y)$
- **Chain rule** 
$$\begin{aligned} P(X_1, X_2, \dots, X_n) &= P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots \\ &= \prod_{i=1}^n P(X_i|X_1, \dots, X_{i-1}) \end{aligned}$$
- **X, Y independent if and only if:**  $\forall x, y : P(x, y) = P(x)P(y)$
- **X and Y are conditionally independent given Z if and only if:**  $X \perp\!\!\!\perp Y|Z$   
 $\forall x, y, z : P(x, y|z) = P(x|z)P(y|z)$


# Reasoning over Time or Space

---

- Often, we want to **reason about a sequence** of observations
  - Speech recognition
  - Robot localization
  - User attention
  - Medical monitoring
- Need to introduce time (or space) into our models

# Markov Models


- Value of  $X$  at a given time is called the **state**



- Parameters: called **transition probabilities** or dynamics, specify how the state evolves over time (also, initial state probabilities)
- Stationarity assumption: transition probabilities the same at all times
- Same as MDP transition model, but no choice of action
- A (growable) BN: We can always use generic BN reasoning on it if we truncate the chain at a fixed length

# Markov Assumption: Conditional Independence

---



- Basic conditional independence:
  - Past and future independent given the present
  - Each time step only depends on the previous
  - This is called the (first order) Markov property


# Example Markov Chain: Weather

- States:  $X = \{\text{rain, sun}\}$



- Initial distribution: 1.0 sun

- CPT  $P(X_t | X_{t-1})$ :

$X_{t-1}$	$X_t$	$P(X_t   X_{t-1})$
sun	sun	0.9
sun	rain	0.1
rain	sun	0.3
rain	rain	0.7




Two new ways of representing the same CPT



# Bayes Nets -- Independence

---




- Bayes Net
- Chain Rule

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$
$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | x_1 \dots x_{i-1})$$

# Markov Models (Markov Chains)

---



- A **Markov model** defines
  - a joint probability distribution:

$$P(X_1, X_2, X_3, X_4) =$$

- More generally:

$$P(X_1, X_2, \dots, X_T) = P(X_1)P(X_2|X_1)P(X_3|X_2) \dots P(X_T|X_{T-1})$$


$$P(X_1, \dots, X_n) = P(X_1) \prod_{t=2}^n P(X_t|X_{t-1})$$

- Why?
- Chain Rule, Indep. Assumption?
- One common inference problem:
  - Compute marginals  $P(X_t)$  for all time steps  $t$



# Example Markov Chain: Weather

- Initial distribution: 1.0 sun



- What is the probability distribution after one step?


$$P(X_2 = sun) = \sum_{x_1} P(x_1, X_2 = sun) = \sum_{x_1} P(X_2 = sun|x_1)P(x_1)$$

$$P(X_2 = sun) = P(X_2 = sun|X_1 = sun)P(X_1 = sun) + P(X_2 = sun|X_1 = rain)P(X_1 = rain)$$

$$0.9 \cdot 1.0 + 0.3 \cdot 0.0 = 0.9$$

# Mini-Forward Algorithm


- Question: What's  $P(X)$  on some day  $t$ ?



$$P(x_1) = \text{known}$$

$$P(x_t) = \sum_{x_{t-1}} P(x_{t-1}, x_t)$$
$$= \sum_{x_{t-1}} P(x_t \mid x_{t-1}) P(x_{t-1})$$

← *Forward simulation*



# Example Run of Mini-Forward Algorithm

- From initial observation of sun

$$\begin{array}{ccccc}
 \left\langle \begin{array}{c} 1.0 \\ 0.0 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.9 \\ 0.1 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.84 \\ 0.16 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.804 \\ 0.196 \end{array} \right\rangle & \longrightarrow & \left\langle \begin{array}{c} 0.75 \\ 0.25 \end{array} \right\rangle \\
 P(X_1) & P(X_2) & P(X_3) & P(X_4) & & P(X_\infty)
 \end{array}$$

- From initial observation of rain

$$\begin{array}{ccccc}
 \left\langle \begin{array}{c} 0.0 \\ 1.0 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.3 \\ 0.7 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.48 \\ 0.52 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.588 \\ 0.412 \end{array} \right\rangle & \longrightarrow & \left\langle \begin{array}{c} 0.75 \\ 0.25 \end{array} \right\rangle \\
 P(X_1) & P(X_2) & P(X_3) & P(X_4) & & P(X_\infty)
 \end{array}$$


- From yet another initial distribution  $P(X_1)$ :

$$\begin{array}{ccc}
 \left\langle \begin{array}{c} p \\ 1-p \end{array} \right\rangle & \dots & \longrightarrow & \left\langle \begin{array}{c} 0.75 \\ 0.25 \end{array} \right\rangle \\
 P(X_1) & & & P(X_\infty)
 \end{array}$$

# Pac-man Markov Chain

---

Pac-man knows the ghost's initial position, but gets no observations!



# Video of Demo Ghostbusters Circular Dynamics

---



# Stationary Distributions

- For most chains:

- Influence of the initial distribution gets less and less over time.
- The distribution we end up in is independent of the initial distribution

- Stationary distribution:

- The distribution we end up with is called the **stationary distribution**  $P_\infty$  of the chain


- It satisfies

$$P_\infty(X) = P_{\infty+1}(X) = \sum_x P(X|x)P_\infty(x)$$



# Example: Stationary Distributions

- Question: What's  $P(X)$  at time  $t = \text{infinity}$ ?



$$P_{\infty}(\text{sun}) = P(\text{sun}|\text{sun})P_{\infty}(\text{sun}) + P(\text{sun}|\text{rain})P_{\infty}(\text{rain})$$

$$P_{\infty}(\text{rain}) = P(\text{rain}|\text{sun})P_{\infty}(\text{sun}) + P(\text{rain}|\text{rain})P_{\infty}(\text{rain})$$


$$P_{\infty}(\text{sun}) = 0.9P_{\infty}(\text{sun}) + 0.3P_{\infty}(\text{rain})$$

$$P_{\infty}(\text{rain}) = 0.1P_{\infty}(\text{sun}) + 0.7P_{\infty}(\text{rain})$$

$$P_{\infty}(\text{sun}) = 3P_{\infty}(\text{rain})$$


$$P_{\infty}(\text{rain}) = 1/3P_{\infty}(\text{sun})$$

Also:  $P_{\infty}(\text{sun}) + P_{\infty}(\text{rain}) = 1$



$$P_{\infty}(\text{sun}) = 3/4$$

$$P_{\infty}(\text{rain}) = 1/4$$




$X_{t-1}$	$X_t$	$P(X_t X_{t-1})$
sun	sun	0.9
sun	rain	0.1
rain	sun	0.3
rain	rain	0.7

# Application of Stationary Distribution: Web Link Analysis


- PageRank over a web graph

- Each web page is a possible value of a state
- Initial distribution: uniform over pages
- Transitions:
  - With prob.  $c$ , uniform jump to a random page (dotted lines, not all shown)
  - With prob.  $1-c$ , follow a random outlink (solid lines)



- Stationary distribution

- Will spend more time on highly reachable pages
- E.g. many ways to get to the Acrobat Reader download page
- Google 1.0 returned the set of pages containing all your keywords in decreasing rank, now all search engines use link analysis along with many other factors (rank actually getting less important over time)






# Hidden Markov Models

---




# Pacman – Sonar




# Hidden Markov Models

- Markov chains not so useful for most agents
  - Need observations to update your beliefs
- Hidden Markov models (HMMs)
  - Underlying Markov chain over states  $X$
  - You observe outputs (effects) at each time step



# Example: Weather HMM



○ An HMM is defined by:

○ Initial distribution:

$$P(X_1)$$

○ Transitions:

$$P(X_t | X_{t-1})$$

○ Emissions:


$$P(E_t | X_t)$$

$R_{t-1}$	$R_t$	$P(R_t   R_{t-1})$
+r	+r	0.7
+r	-r	0.3
-r	+r	0.3
-r	-r	0.7

$R_t$	$U_t$	$P(U_t   R_t)$
+r	+u	0.9
+r	-u	0.1
-r	+u	0.2
-r	-u	0.8


# Example: Ghostbusters HMM

- $P(X_1) = \text{uniform}$
- $P(X|X') = \text{usually move clockwise, but sometimes move in a random direction or stay in place}$
- $P(R_{ij}|X) = \text{same sensor model as before: red means close, green means far away.}$




1/9	1/9	1/9
1/9	1/9	1/9
1/9	1/9	1/9

$P(X_1)$



1/6	1/6	1/2
0	1/6	0
0	0	0

$P(X|X'=\langle 1,2 \rangle)$




# Video of Demo Ghostbusters – Circular Dynamics -- HMM

---



# Conditional Independence

- HMMs have two important independence properties:
  - Markov hidden process: future depends on past via the present
  - Current observation independent of all else given current state



- Does this mean that evidence variables are guaranteed to be independent?
  - [No, they tend to be correlated by the hidden state]

# Real HMM Examples

---

- Robot tracking:
  - Observations are range readings (continuous)
  - States are positions on a map (continuous)
- Speech recognition HMMs:
  - Observations are acoustic signals (continuous valued)
  - States are specific positions in specific words (so, tens of thousands)
- Machine translation HMMs:
  - Observations are words (tens of thousands)
  - States are translation options





# Filtering / Monitoring

---

- Filtering, or monitoring, is the task of tracking the distribution  $B_t(X) = P_t(X_t \mid e_1, \dots, e_t)$  (the belief state) over time
- We start with  $B_1(X)$  in an initial setting, usually uniform
- As time passes, or we get observations, we update  $B(X)$
- The Kalman filter was invented in the 60's and first implemented as a method of trajectory estimation for the Apollo program

# Example: Robot Localization



Example from  
Michael Pfeiffer



Sensor model: can read in which directions there is a wall, never more than 1 mistake


Motion model: may not execute action with small prob.

# Example: Robot Localization




Lighter grey: was possible to get the reading, but less likely b/c required 1 mistake


# Example: Robot Localization




Prob




t=2




# Example: Robot Localization




Prob



t=3



# Example: Robot Localization




Prob



0

1


$t=4$



# Example: Robot Localization



t=5



# Inference: Find State Given Evidence

---

- We are given evidence at each time and want to know

$$B_t(X) = P(X_t | e_{1:t})$$

- Idea: start with  $P(X_1)$  and derive  $B_t$  in terms of  $B_{t-1}$ 
  - equivalently, derive  $B_{t+1}$  in terms of  $B_t$



# Background:

## Probabilistic Inference

---

- Probabilistic inference: compute a desired probability from other known probabilities (e.g. conditional from joint)
- We generally compute conditional probabilities
  - $P(\text{on time} \mid \text{no reported accidents}) = 0.90$
  - These represent the agent's *beliefs* given the evidence
- Probabilities change with new evidence:
  - $P(\text{on time} \mid \text{no accidents, 5 a.m.}) = 0.95$
  - $P(\text{on time} \mid \text{no accidents, 5 a.m., raining}) = 0.80$
  - Observing new evidence causes *beliefs to be updated*



# Inference by Enumeration

---

- $P(W)$ ?

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

---

- $P(W)$ ?

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

○ P(W)?

$$P(\text{sun}) = .3 + .1 + .1 + .15 = .65$$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

○  $P(W)$ ?

$$P(\text{sun}) = .3 + .1 + .1 + .15 = .65$$

$$P(\text{rain}) = 1 - .65 = .35$$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

- General case:


- Evidence variables:  $E_1 \dots E_k = e_1 \dots e_k$
  - Query\* variable:  $Q$
  - Hidden variables:  $H_1 \dots H_r$
- }  $X_1, X_2, \dots, X_n$   
All variables

- We want:


$$P(Q|e_1 \dots e_k)$$

*\* Works fine with multiple query variables, too*

- Step 1: Select the entries consistent with the evidence



- Step 2: Sum out H to get joint of Query and evidence



- Step 3: Normalize

$$\times \frac{1}{Z}$$

$$Z = \sum_q P(Q, e_1 \dots e_k)$$

$$P(Q|e_1 \dots e_k) = \frac{1}{Z} P(Q, e_1 \dots e_k)$$

$$P(Q, e_1 \dots e_k) = \sum_{h_1 \dots h_r} P(Q, \underbrace{h_1 \dots h_r}_{X_1, X_2, \dots, X_n}, e_1 \dots e_k)$$

# Inference by Enumeration

---

- $P(W \mid \text{winter})?$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

- $P(W \mid \text{winter})?$

$$P(\text{sun}|\text{winter}) \sim .1 + .15 = .25$$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20



# Inference by Enumeration

---

- $P(W \mid \text{winter})?$

$$P(\text{rain} \mid \text{winter}) \sim .05 + .2 = .25$$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

- $P(W \mid \text{winter})?$

$P(\text{sun} \mid \text{winter}) \sim .25$

$P(\text{rain} \mid \text{winter}) \sim .25$

$P(\text{sun} \mid \text{winter}) = .5$

$P(\text{rain} \mid \text{winter}) = .5$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

---

- $P(W \mid \text{winter, hot})?$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

---

- $P(W \mid \text{winter, hot})?$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

- $P(W \mid \text{winter, hot})?$

$P(\text{sun} \mid \text{winter, hot}) \sim .1$

$P(\text{rain} \mid \text{winter, hot}) \sim .05$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

- $P(W \mid \text{winter, hot})?$

$P(\text{sun} \mid \text{winter, hot}) \sim .1$

$P(\text{rain} \mid \text{winter, hot}) \sim .05$

$P(\text{sun} \mid \text{winter, hot}) = 2/3$

$P(\text{rain} \mid \text{winter, hot}) = 1/3$

S	T	W	P
summer	hot	sun	0.30
summer	hot	rain	0.05
summer	cold	sun	0.10
summer	cold	rain	0.05
winter	hot	sun	0.10
winter	hot	rain	0.05
winter	cold	sun	0.15
winter	cold	rain	0.20

# Inference by Enumeration

---

- Obvious problems:
  - Worst-case time complexity  $O(d^n)$
  - Space complexity  $O(d^n)$  to store the joint distribution

# Next Topic

---

- Inference in HMMs