

# CSE 493G1/599G1: Deep Learning

---

## Section 1: Fundamentals

Welcome to section, we're happy you're here!

### Reference Material

Rules of Broadcasting from Jake VanderPlas' *Python Data Science Handbook*:

- (1) If the two arrays differ in their number of dimensions, the shape of the one with fewer dimensions is padded with ones on its leading (left) side.
- (2) If the shape of the two arrays does not match in any dimension, the array with shape equal to 1 in that dimension is stretched to match the other shape.
- (3) If in any dimension the sizes disagree and neither is equal to 1, an error is raised.

Chain Rule for One Independent Variable:

Let  $z = f(x, y)$  be a differentiable function. Further suppose that  $x$  and  $y$  are themselves differentiable functions of  $t$ , in other words  $x = x(t)$  and  $y = y(t)$ . Then,

$$\frac{\partial z}{\partial t} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial t}$$

Chain Rule for Two Independent Variables:

Let  $z = f(x, y)$  be a differentiable function, where  $x$  and  $y$  are themselves differentiable functions of  $a$  and  $b$ . In other words,  $x = x(a, b)$  and  $y = y(a, b)$ . Then,

$$\frac{\partial z}{\partial a} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial a} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial a}$$

and

$$\frac{\partial z}{\partial b} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial b} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial b}$$

Generalized Chain Rule from Gilbert Strang's *Calculus*:

Let  $w = f(x_1, x_2, \dots, x_m)$  be a differentiable function of  $m$  independent variables, and for each  $i \in 1, \dots, m$ , let  $x_i = x_i(t_1, t_2, \dots, t_n)$  be a differentiable function of  $n$  independent variables. Then,

$$\frac{\partial w}{\partial t_j} = \frac{\partial w}{\partial x_1} \frac{\partial x_1}{\partial t_j} + \frac{\partial w}{\partial x_2} \frac{\partial x_2}{\partial t_j} + \dots + \frac{\partial w}{\partial x_m} \frac{\partial x_m}{\partial t_j}$$

for any  $j \in 1, 2, \dots, n$ .

## 1. Broadcasting

Determine if NumPy allows the addition of the following pairs of arrays, and if applicable determine what the result's dimensions will be.

(a) Where `x.shape` is `(2,)` and `y.shape` is `(2, 1)`

(b) Where `x.shape` is `(4,)` and `y.shape` is `(4, 1, 1)`

(c) Where `x.shape` is `(4, 2)` and `y.shape` is `(2, 4, 1)`

(d) Where `x.shape` is `(8, 3)` and `y.shape` is `(2, 8, 1)`

(e) Where `x.shape` is `(6, 5, 3)` and `y.shape` is `(6, 5)`

## 2. Multivariable Chain Rule

(a) Let  $z = x^2y - y^2$  where  $x = t^2$  and  $y = 2t$ . Find  $\frac{dz}{dt}$ .

(b) Let  $z = 3x^2 - 2xy + y^2$ . Also let  $x = 3a + 2b$  and  $y = 4a - b$ . Find  $\frac{\partial z}{\partial a}$  and  $\frac{\partial z}{\partial b}$ .

(c) Let  $w = f(x, y, z)$ ,  $x = x(t, u, v)$ ,  $y = y(t, u, v)$  and  $z = z(t, u, v)$ . Find the formula for  $\frac{\partial w}{\partial t}$ .