

Name: _____

**CSE 505, Fall 2005, Midterm Examination
8 November 2005**

Please do not turn the page until everyone is ready.

Rules:

- The exam is closed-book, closed-note, except for one side of one 8.5x11in piece of paper.
- **Please stop promptly at 1:20.**
- You can rip apart the pages, but please write your name on each page.
- There are **140 points** total, distributed **unevenly** among 6 questions (which have multiple parts).

Advice:

- Read questions carefully. Understand a question before you start writing.
- Write down thoughts and intermediate steps so you can get partial credit.
- The questions are not necessarily in order of difficulty. **Skip around.** In particular, do not spend so much time on a proof that you do not get to all the problems.
- If you have questions, ask.
- Relax. You are here to learn.

Name: _____

For your reference:

$s ::= \text{skip} \mid x := e \mid s; s \mid \text{if } e \text{ s } s \mid \text{while } e \text{ s}$
 $e ::= c \mid x \mid e + e \mid e * e$
 $(c \in \{\dots, -2, -1, 0, 1, 2, \dots\})$
 $(x \in \{x_1, x_2, \dots, y_1, y_2, \dots, z_1, z_2, \dots, \dots\})$

$H; e \Downarrow c$

$\frac{\text{CONST}}{H; c \Downarrow c}$ $\frac{\text{VAR}}{H; x \Downarrow H(x)}$ $\frac{\text{ADD}}{H; e_1 \Downarrow c_1 \quad H; e_2 \Downarrow c_2}{H; e_1 + e_2 \Downarrow c_1 + c_2}$ $\frac{\text{MULT}}{H; e_1 \Downarrow c_1 \quad H; e_2 \Downarrow c_2}{H; e_1 * e_2 \Downarrow c_1 * c_2}$

$H_1; s_1 \rightarrow H_2; s_2$

$\frac{\text{ASSIGN}}{H; x := e \rightarrow H, x \mapsto c; \text{skip}}$ $\frac{\text{SEQ1}}{H; \text{skip}; s \rightarrow H; s}$ $\frac{\text{SEQ2}}{H; s_1 \rightarrow H'; s'_1}{H; s_1; s_2 \rightarrow H'; s'_1; s_2}$
 $\frac{\text{IF1}}{H; e \Downarrow c \quad c > 0}{H; \text{if } e \text{ s}_1 \text{ s}_2 \rightarrow H; s_1}$ $\frac{\text{IF2}}{H; e \Downarrow c \quad c \leq 0}{H; \text{if } e \text{ s}_1 \text{ s}_2 \rightarrow H; s_2}$

$e ::= \lambda x. e \mid x \mid e e \mid c$
 $v ::= \lambda x. e \mid c$
 $\tau ::= \text{int} \mid \tau \rightarrow \tau$

$e \rightarrow e'$

$\frac{}{(\lambda x. e) v \rightarrow e[v/x]}$ $\frac{e_1 \rightarrow e'_1}{e_1 e_2 \rightarrow e'_1 e_2}$ $\frac{e_2 \rightarrow e'_2}{v e_2 \rightarrow v e'_2}$

$e[e'/x] = e''$

$\frac{}{x[e/x] = e}$ $\frac{e_1[e/x] = e'_1 \quad y \neq x \quad y \notin FV(e)}{(\lambda y. e_1)[e/x] = \lambda y. e'_1}$
 $\frac{y \neq x}{y[e/x] = y}$ $\frac{e_1[e/x] = e'_1 \quad e_2[e/x] = e'_2}{(e_1 e_2)[e/x] = e'_1 e'_2}$

$\Gamma \vdash e : \tau$

$\frac{}{\Gamma \vdash c : \text{int}}$ $\frac{}{\Gamma \vdash x : \Gamma(x)}$ $\frac{\Gamma, x : \tau_1 \vdash e : \tau_2}{\Gamma \vdash \lambda x. e : \tau_1 \rightarrow \tau_2}$ $\frac{\Gamma \vdash e_1 : \tau_2 \rightarrow \tau_1 \quad \Gamma \vdash e_2 : \tau_2}{\Gamma \vdash e_1 e_2 : \tau_1}$

Name: _____

1. (IMP with choice)

- (a) (**10** points) Let “?” be a choice operator for IMP expressions: $e_1?e_2$ chooses either e_1 or e_2 and evaluates its choice to produce an answer. Give semantic rules for this extension.
- (b) (**20** points) Theorem: If e_1 is equivalent to e_2 , then e_1 is equivalent to $e_1?e_2$.
- Restate this theorem formally.
 - Prove this theorem formally.

Name: _____

2. (Bad statement rules)

(a) (10 points) Why do we not have this rule in our IMP statement semantics?

$$\frac{H ; s_1 \rightarrow H' ; s'_1}{H ; s_1 ; (s_2 ; s_3) \rightarrow H' ; s'_1 ; (s_2 ; s_3)}$$

(b) (10 points) Why do we not have this rule in our IMP statement semantics?

$$\frac{H ; s_1 \rightarrow H' ; s'_1}{H ; s_2 ; s_1 \rightarrow H' ; s_2 ; s'_1}$$

Name: _____

3. (Functional programming)

(a) (10 points) Consider this Caml code:

```
type t = A of int | B of (int->int)
let x = 2
let f y = x + y
let ans1 = (let x = 3 in
            let a = A (f 4) in
            let x = 5 in
            match a with A x -> x | B x -> x 6)
let ans2 = (let x = 3 in
            let b = B f in
            let x = 5 in
            match b with A x -> x | B x -> x 6)
```

After evaluating this code, what values are `ans1` and `ans2` bound to?

(b) (10 points) Consider this Caml code:

```
let rec g x =
  match x with
  [] -> []
  | hd::tl -> (fun y -> hd + y)::(g tl)
```

- i. What does this function do?
- ii. What is this function's type?
- iii. Write a function `h` that is the *inverse* of `g`. That is, `fun x -> h (g x)` would return a value equivalent to its input.

Name: _____

4. (λ encodings) Recall this encoding of booleans in the λ -calculus:

“true” $\lambda x. \lambda y. x$

“false” $\lambda x. \lambda y. y$

“if” $\lambda b. \lambda t. \lambda f. b t f$

- (a) (10 points) Extend this encoding with a λ term that encodes (*inclusive*) *or*.
- (b) (10 points) Extend this encoding with a λ term that encodes *not*.

Name: _____

5. (Simply-Typed λ calculus)

For all subproblems, assume the simply-typed λ calculus.

- (a) (**6** points) Give a Γ , e_1 , e_2 , and τ such that $\Gamma \vdash e_1 : \tau$ and $\Gamma \vdash e_2 : \tau$ and $e_1 \neq e_2$.
- (b) (**6** points) Give a Γ_1 , Γ_2 , e , and τ such that $\Gamma_1 \vdash e : \tau$ and $\Gamma_2 \vdash e : \tau$ and $\Gamma_1 \neq \Gamma_2$.
- (c) (**8** points) Give a Γ , e , τ_1 , and τ_2 such that $\Gamma \vdash e : \tau_1$ and $\Gamma \vdash e : \tau_2$ and $\tau_1 \neq \tau_2$.

Name: _____

6. (Type-Safety)

We add an explicit infinite-loop to the simply-typed λ -calculus: The term ∞ simply “reduces to itself”.

- (a) (5 points) Extend the semantics of the call-by-value λ -calculus to include ∞ .
- (b) (10 points) Extend the type system of the simply-typed λ -calculus to include ∞ . Be as permissive as possible considering the next problem.
- (c) (15 points) Prove that your extensions maintain type safety. Do *not* repeat the entire type-safety proof. Rather, for each of these lemmas, remind us the structure of the proof (i.e., the induction hypothesis) and then prove any new cases introduced by your extensions.
 - Preservation: If $\cdot \vdash e : \tau$ and $e \rightarrow e'$, then $\cdot \vdash e' : \tau$.
 - Progress: If $\cdot \vdash e : \tau$, then e is a value or there exists an e' such that $e \rightarrow e'$.
 - Substitution: If $\Gamma, x:\tau' \vdash e_1 : \tau$ and $\Gamma \vdash e_2 : \tau'$, then $\Gamma \vdash e_1[e_2/x] : \tau$.