CSE 531
Automata, Computability, Complexity

Myhill-Nerode Theorem

Paul Beame 6 October 1999

Let A be any language over Σ^*. We say that strings x and y in Σ^* are *indistinguishable by A* iff for every string $z \in \Sigma^*$ either both xz and yz are in A or both xz and yz are not in A. We write $x \equiv_A y$ in this case. Note that $\equiv_A$ is an equivalence relation. (Check this yourself.)

Given a DFA $M = (Q, \Sigma, \delta, s, F)$ we say that two strings x and y in Σ^* are *indistinguishable by M* iff $\delta^*(s,x) = \delta^*(s,y)$, i.e. the state reached by M on input x is the same as the state reached by M on input y. We write $x \equiv_M y$ in this case. Note that $\equiv_M$ is an equivalence relation and that it only has a finite number of equivalence classes, one per state.

Lemma 1: If $A = L(M)$ for a DFA M then for any $x, y \in \Sigma^*$ if $x \equiv_M y$ then $x \equiv_A y$.

Proof Suppose that $A = L(M)$. Therefore $w \in A \Leftrightarrow \delta^*(s,w) \in F$. Suppose also that $x \equiv_M y$. Then $\delta^*(s,x) = \delta^*(s,y)$.

Let $z \in \Sigma^*$. Clearly $\delta^*(s,xz) = \delta^*(s,yz)$. Therefore

$$\begin{aligned} xz \in A &\Leftrightarrow \delta^*(s,xz) \in F \\ &\Leftrightarrow \delta^*(s,yz) \in F \\ &\Leftrightarrow yz \in A \end{aligned}$$

It follows that $x \equiv_A y$. $\square$

This lemma says that whenever two elements arrive at the same state of M they are in the same equivalence class of $\equiv_A$. This means that each equivalence class of $\equiv_A$ is a union of equivalence classes of $\equiv_M$.

Corollary 2: If A is regular then $\equiv_A$ has a finite number of equivalence classes.

Proof Let M be a DFA such that $A = L(M)$. The Lemma shows that $\equiv_A$ has at most as many equivalence classes as $\equiv_M$, which has a finite number of equivalence classes (equal to the number of states of M). $\square$

We now get another way of proving that languages are not regular, namely given A find an infinite sequence of strings $x_1, x_2, \ldots$ and prove that they are not equivalent to each other with respect to $\equiv_A$.

Claim: $A = \{0^n 1^n \| n \geq 0\}$ is not regular.

Proof Consider the sequence of strings $x_1, x_2, \ldots$ where $x_i = 0^i$ for $i \geq 1$. We now see that no two of these are equivalent to each other with respect to $\equiv_A$: Consider $x_i = 0^i$ and $x_j = 0^j$ for $i \neq j$. Let $z = 1^i$ and notice that $x_i z = 0^i 1^i \in A$ but $x_j z = 0^j 1^i \notin A$. Therefore no two of these strings are equivalent to each other and thus A cannot be regular. $\square$

One nice thing about this method for proving things nonregular is that, unlike the pumping lemma, it is always guaranteed to be work because the corollary above is a precise characterization of the regular languages. The statement of this fact is known as the Myhill-Nerode Theorem.

Theorem 3: A is regular if and only if $\equiv_A$ has a finite number of equivalences classes. Furthermore there is a DFA M with $L(M) = A$ having precisely one state for each equivalence class of $\equiv_A$.

Proof The corollary above already gives one direction of this statement. All we now need to show is that if $\equiv_A$ has a finite number of equivalence classes then we can build a DFA $M = (Q, \Sigma, \delta, s, F)$ accepting A where there is one state in Q for each equivalence class of $\equiv_A$. Here is how it goes:

Let $A_1, \ldots, A_r$ be the equivalence classes of $\equiv_A$. Remember that the A_i are disjoint and their union is all of Σ^*. Define $Q = \{1, \ldots, r\}$.

Let $s \in Q$ be the one i such that $\epsilon \in A_i$.

Note that for any A_j and any $a \in \Sigma$, for every $x, y \in A_j$, xa and ya will both be contained in the same equivalence class of $\equiv_A$. (For any $z \in \Sigma^*$, $xaz \in A \Leftrightarrow yaz \in A$ since x and y are in the same equivalence class of $\equiv_A$.)

To figure out what $\delta(j, a)$ should be, all we do is pick some $x \in A_j$, find the one k such that $xa \in A_k$ and set $\delta(j, a) = k$. The answer will be the same no matter which x we choose.

To pick the final states, note that for each j, either $A_j \subset A$ or $A_j \cap A = \emptyset$. Therefore we let $F = \{j \mid A_j \subseteq A\}$.

It is easy to argue by induction that $\delta^*(s, x) = j \Leftrightarrow x \in A_j$. This, together with the choice of F ensures that $L(M) = A$. $\square$

By the proof of the corollary above we know that the number of states of M constructed above is the smallest possible. (In fact, if one looks at things carefully one can show that all DFA's of that size for A basically have to look the same.)

However, in general, even though A is a regular language we may not have a nice description of $\equiv_A$ at our disposal in order to build M. What happens if all we have is some DFA accepting A?