

Shading

1

Reading

Required:

- ◆ Shirley, Chapter 9

2

Introduction

Affine transformations help us to place objects into a scene.

Before creating images of these objects, we'll look at models for how light interacts with their surfaces.

Such a model is called a **shading model**.

Other names:

- ◆ Lighting model
- ◆ Light reflection model
- ◆ Local illumination model
- ◆ Reflectance model
- ◆ BRDF

3

An abundance of photons

Properly determining the right color is *really hard*.

Look around the room. Each light source has different characteristics. Trillions of photons are pouring out every second.

These photons can:

- ◆ interact with the atmosphere, or with things in the atmosphere
- ◆ strike a surface and
 - be absorbed
 - be reflected (scattered)
 - cause fluorescence or phosphorescence.
- ◆ interact in a wavelength-dependent manner
- ◆ generally bounce around and around

4

Our problem

We're going to build up to an *approximation* of reality called the **Phong illumination model**.

It has the following characteristics:


- ♦ *not* physically based
- ♦ gives a "first-order" *approximation* to physical light reflection
- ♦ very fast
- ♦ widely used

In addition, we will assume **local illumination**, i.e., light goes: light source → surface → viewer.

No interreflections, no shadows.

5

Setup...



Given:

- ♦ a point **P** on a surface visible through pixel *p*
- ♦ The normal **N** at **P**
- ♦ The lighting direction, **L**, and intensity, *L*, at **P**
- ♦ The viewing direction, **V**, at **P**
- ♦ The shading coefficients at **P**

Compute the color, *I*, of pixel *p*.

Assume that the direction vectors are normalized:

$$\|\mathbf{N}\| = \|\mathbf{L}\| = \|\mathbf{V}\| = 1$$

6

"Iteration zero"

The simplest thing you can do is...

Assign each polygon a single color:

$$I = k_e$$

where

- ♦ *I* is the resulting intensity
- ♦ k_e is the **emissivity** or intrinsic shade associated with the object

This has some special-purpose uses, but not really good for drawing a scene.

[Note: k_e is omitted in Shirley.]

7

"Iteration one"

Let's make the color at least dependent on the overall quantity of light available in the scene:

$$I = k_e + k_a L_a$$

- ♦ k_a is the **ambient reflection coefficient**.
 - really the reflectance of ambient light
 - "ambient" light is assumed to be equal in all directions
- ♦ L_a is the **ambient light intensity**.

Physically, what is "ambient" light?

[Note: Shirley uses c_r and c_a instead of k_a and L_a .

8

Wavelength dependence

Really, k_e , k_a , and L_a are functions over all wavelengths λ .

Ideally, we would do the calculation on these functions. For the ambient shading equation, we would start with:

$$I(\lambda) = k_a(\lambda)L_a(\lambda)$$

then we would find good RGB values to represent the spectrum $I(\lambda)$.

Traditionally, though, k_a and L_a are represented as RGB triples, and the computation is performed on each color channel separately:

$$I_R = k_{a,R} L_{a,R}$$

$$I_G = k_{a,G} L_{a,G}$$

$$I_B = k_{a,B} L_{a,B}$$

9

Diffuse reflection

Let's examine the ambient shading model:

- ♦ objects have different colors
- ♦ we can control the overall light intensity
 - what happens when we turn off the lights?
 - what happens as the light intensity increases?
 - what happens if we change the color of the lights?

So far, objects are uniformly lit.

- ♦ not the way things really appear
- ♦ in reality, light sources are localized in position or direction

Diffuse, or **Lambertian** reflection will allow reflected intensity to vary with the direction of the light.


10

Diffuse reflectors

Diffuse reflection occurs from dull, matte surfaces, like latex paint, or chalk.

These **diffuse** or **Lambertian** reflectors reradiate light equally in all directions.


Picture a rough surface with lots of tiny **microfacets**.



11

Diffuse reflectors

...or picture a surface with little pigment particles embedded beneath the surface (neglect reflection at the surface for the moment):



The microfacets and pigments distribute light rays in all directions.


Embedded pigments are responsible for the coloration of diffusely reflected light in plastics and paints.

Note: the figures above are intuitive, but not strictly (physically) correct.

12

Diffuse reflectors, cont.

The reflected intensity from a diffuse surface does not depend on the direction of the viewer. The incoming light, though, does depend on the direction of the light source:



13

“Iteration two”

The incoming energy is proportional to _____, giving the diffuse reflection equations:

$$I = k_e + k_a L_a + k_d L \cos \theta$$

$$= k_e + k_a I_a + k_d L (\cos \theta)$$

where:

- ♦ k_d is the **diffuse reflection coefficient**
- ♦ L is the intensity of the light source
- ♦ \mathbf{N} is the normal to the surface (unit vector)
- ♦ \mathbf{L} is the direction to the light source (unit vector)
- ♦ $(x)_+$ means $\max\{0, x\}$

[Note: Shirley uses c_r and c_l instead of k_d and L .]

14

Specular reflection

Specular reflection accounts for the highlight that you see on some objects.

It is particularly important for *smooth, shiny* surfaces, such as:


- ♦ metal
- ♦ polished stone
- ♦ plastics
- ♦ apples
- ♦ skin

Properties:

- ♦ Specular reflection depends on the viewing direction \mathbf{V} .
- ♦ For non-metals, the color is determined solely by the color of the light.
- ♦ For metals, the color may be altered (e.g., brass)

15

Specular reflection “derivation”



For a perfect mirror reflector, light is reflected about \mathbf{N} , so

$$I = \begin{cases} L & \text{if } \mathbf{V} = \mathbf{R} \\ 0 & \text{otherwise} \end{cases}$$


For a near-perfect reflector, you might expect the highlight to fall off quickly with increasing angle ϕ .

Also known as:

- ♦ “**rough specular**” reflection
- ♦ “**directional diffuse**” reflection
- ♦ “**glossy**” reflection

16

Derivation, cont.



One way to get this effect is to take $(\mathbf{R} \cdot \mathbf{V})$, raised to a power n_s .

As n_s gets larger,

- ♦ the dropoff becomes {more,less} gradual
- ♦ gives a {larger,smaller} highlight
- ♦ simulates a {more,less} mirror-like surface

17

“Iteration three”

The next update to the Phong shading model is then:

$$I = k_e + k_a I_a + k_d L(\mathbf{N} \cdot \mathbf{L})_+ + k_s L(\mathbf{V} \cdot \mathbf{R})_+^{n_s}$$

where:

- ♦ k_s is the **specular reflection coefficient**
- ♦ n_s is the **specular exponent** or **shininess**
- ♦ \mathbf{R} is the reflection of the light about the normal (unit vector)
- ♦ \mathbf{V} is viewing direction (unit vector)

[Note: Shirley uses \mathbf{e} , \mathbf{r} , and p instead of \mathbf{V} , \mathbf{R} , and n_s . Also, OpenGL uses a separate L_d and L_s , instead of a single L .]


18

Lights

OpenGL supports three different kinds of lights: ambient, directional, and point. Spot lights are also supported as a special form of point light.

We’ve seen ambient light sources, which are not really geometric.

Directional light sources have a single direction and intensity associated with them.




[Note: Lights not discussed in Shirley.]

19

Point lights

The direction of a **point light** sources is determined by the vector from the light position to the surface point.



$$\mathbf{L} = \frac{\mathbf{E} - \mathbf{P}}{\|\mathbf{E} - \mathbf{P}\|}$$

$$d = \|\mathbf{E} - \mathbf{P}\|$$

Physics tells us the intensity must drop off inversely with the square of the distance:

$$f_{\text{atten}} = \frac{1}{d^2}$$

Sometimes, this distance-squared dropoff is considered too “harsh.” A common alternative is:


$$f_{\text{atten}} = \frac{1}{a + bd + cd^2}$$

with user-supplied constants for a , b , and c .

20

Spotlights

OpenGL also allows one to apply a *directional attenuation* of a point light source, giving a **spotlight** effect.



The spotlight intensity factor is computed in OpenGL as:

$$f_{\text{spot}} = (\mathbf{L} \cdot \mathbf{S})_{\beta}^e$$

where

- ♦ \mathbf{L} is the direction to the point light.
- ♦ \mathbf{S} is the center direction of the spotlight.
- ♦ β is the cutoff angle for the spotlight
- ♦ e is the angular falloff coefficient
- ♦ $(x)_{\beta}^e = [\max\{\cos(x) - \beta, 0\}]^e$

21

“Iteration four”

Since light is additive, we can handle multiple lights by taking the sum over every light.

Our equation is now:

$$I = k_e + k_a L_a + \sum_j \frac{(\mathbf{L}_j \cdot \mathbf{S}_j)_{\beta_j}^{e_j}}{a_j + b_j d_j + c_j d_j^2} L_j \left[k_d (\mathbf{N} \cdot \mathbf{L}_j)_+ + k_s (\mathbf{V} \cdot \mathbf{R}_j)_+^{n_s} \right]$$

This is the Phong illumination model.

Which quantities are spatial vectors?

Which are RGB triples?

Which are scalars?

22

Choosing the parameters

Experiment with different parameter settings. To get you started, here are a few suggestions:

- ♦ Try n_s in the range [0,100]
- ♦ Try $k_a + k_d + k_s < 1$
- ♦ Use a small k_a (~0.1)

	n_s	k_d	k_s
Metal	large	Small, color of metal	Large, color of metal
Plastic	medium	Medium, color of plastic	Medium, white
Planet	0	varying	0

23

Materials in OpenGL

The OpenGL code to specify the surface shading properties is fairly straightforward. For example:

```
GLfloat ke[] = { 0.1, 0.15, 0.05, 1.0 };
GLfloat ka[] = { 0.1, 0.15, 0.1, 1.0 };
GLfloat kd[] = { 0.3, 0.3, 0.2, 1.0 };
GLfloat ks[] = { 0.2, 0.2, 0.2, 1.0 };
GLfloat ns[] = { 50.0 };
glMaterialfv(GL_FRONT, GL_EMISSION, ke);
glMaterialfv(GL_FRONT, GL_AMBIENT, ka);
glMaterialfv(GL_FRONT, GL_DIFFUSE, kd);
glMaterialfv(GL_FRONT, GL_SPECULAR, ks);
glMaterialfv(GL_FRONT, GL_SHININESS, ns);
```

Notes:

- ♦ The `GL_FRONT` parameter tells OpenGL that we are specifying the materials for the front of the surface.
- ♦ Only the alpha value of the diffuse color is used for blending. It's usually set to 1.

24

Shading in OpenGL

The OpenGL lighting model allows you to associate different lighting colors according to material properties they will influence.

Thus, our original shading equation:

$$I = k_e + k_a L_a + \sum_j \frac{(\mathbf{L}_j \cdot \mathbf{S}_j)^{\beta_j}}{a_j + b_j d_j + c_j d_j^2} L_j \left[k_d (\mathbf{N} \cdot \mathbf{L}_j)_+ + k_s (\mathbf{V} \cdot \mathbf{R}_j)_+^{n_s} \right]$$

becomes:

$$I = k_e + k_a L_a + \sum_j \frac{(\mathbf{L}_j \cdot \mathbf{S}_j)^{\beta_j}}{a_j + b_j d_j + c_j d_j^2} \left[k_a L_{a_j} + k_d L_{d_j} (\mathbf{N} \cdot \mathbf{L}_j)_+ + k_s L_{s_j} (\mathbf{V} \cdot \mathbf{R}_j)_+^{n_s} \right]$$

where you can have a global ambient light with intensity L_a in addition to have an ambient light intensity L_{a_j} associated with each individual light.

25

Shading in OpenGL, cont'd

In OpenGL this equation, for one light source (the 0th) is specified something like:

```
GLfloat La[] = { 0.2, 0.2, 0.2, 1.0 };
GLfloat La0[] = { 0.1, 0.1, 0.1, 1.0 };
GLfloat Ld0[] = { 1.0, 1.0, 1.0, 1.0 };
GLfloat Ls0[] = { 1.0, 1.0, 1.0, 1.0 };
GLfloat pos0[] = { 1.0, 1.0, 1.0, 0.0 };
GLfloat a0[] = { 1.0 };
GLfloat b0[] = { 0.5 };
GLfloat c0[] = { 0.25 };
GLfloat S0[] = { -1.0, -1.0, 0.0 };
GLfloat beta0[] = { 45 };
GLfloat e0[] = { 2 };
```

```
glLightModelfv(GL_LIGHT_MODEL_AMBIENT, La);
glLightfv(GL_LIGHT0, GL_AMBIENT, La0);
glLightfv(GL_LIGHT0, GL_DIFFUSE, Ld0);
glLightfv(GL_LIGHT0, GL_SPECULAR, Ls0);
glLightfv(GL_LIGHT0, GL_POSITION, pos0);
glLightfv(GL_LIGHT0, GL_CONSTANT_ATTENUATION, a0);
glLightfv(GL_LIGHT0, GL_LINEAR_ATTENUATION, b0);
glLightfv(GL_LIGHT0, GL_QUADRATIC_ATTENUATION, c0);
glLightfv(GL_LIGHT0, GL_SPOT_DIRECTION, S0);
glLightf(GL_LIGHT0, GL_SPOT_CUTOFF, beta0);
glLightf(GL_LIGHT0, GL_SPOT_EXPONENT, e0);
```

26

Shading in OpenGL, cont'd

Notes:

You can have as many as `GL_MAX_LIGHTS` lights in a scene. This number is system-dependent.

For directional lights, you specify a light direction, not position, and the attenuation and spotlight terms are ignored.

The directions of directional lights and spotlights are specified in the coordinate systems of the lights, not the surface points as we've been doing in lecture.

27

BRDF

The Phong illumination model is really a function that maps light from incoming (light) directions ω_{in} to outgoing (viewing) directions ω_{out} :

$$f_r(\omega_{in}, \omega_{out})$$

This function is called the **Bi-directional Reflectance Distribution Function (BRDF)**.

Here's a plot with ω_{in} held constant:





BRDF's can be quite sophisticated...

28

More sophisticated BRDF's

Cook and
Torrance, 1982



Westin, Arvo, Torrance 1992

