

# **Surfaces of Revolution**


**Brian Curless  
CSE 557  
Autumn 2017**

# Surfaces of revolution

Idea: rotate a 2D **profile curve** around an axis.

What kinds of shapes can you model this way?

## Constructing surfaces of revolution



**Given:** A curve  $C(v)$  in the  $xy$ -plane:

$$C(v) = \begin{bmatrix} C_x(v) \\ C_y(v) \\ 0 \\ 1 \end{bmatrix}$$


Let  $R_y(\theta)$  be a rotation about the  $y$ -axis.

**Find:** A surface  $S(u,v)$  which is  $C(v)$  rotated about the  $y$ -axis, where  $u, v \in [0, 1]$ .

**Solution:**

## Constructing surfaces of revolution

We can sample in  $u$  and  $v$  to get a grid of points over the surface.



Suppose we sample:

- ♦ in  $v$ , to give  $C[j]$  where  $j \in [0..M-1]$
- ♦ in  $u$ , to give rotation angle  $\theta[i] = 2\pi i / N$  where  $i \in [0..N]$


We can now write the surface as:

How would we turn this into a mesh of triangles?  
How do we assign per-vertex normals?

## Surface normals



Now that we describe the surface as a triangle mesh, we need to provide surface normals. As we'll see later, these normals are important for drawing and shading the surface (i.e., for "rendering").

One approach is to compute the normal to each triangle. How do we compute these normals?




For surfaces of revolution, we can get better-looking results by analytically computing the normal at each vertex...

## Tangent vectors, tangent planes, and normals



## Normals on a surface of revolution



We can compute tangents to the curve points in the  $xy$ -plane:

$$\mathbf{T}_1[0, j] \approx$$


$$\mathbf{T}_2[0, j] =$$

to get the normal in that plane:

$$\mathbf{N}[0, j] =$$

and then rotate it around:

## Texture coordinates on a surface of revolution



For our surface of revolution, we now have:

**Profile curve:**  $C[j]$  where  $j \in [0..M-1]$


**Rotation angles:**  $\theta[i] = 2\pi i / N$  where  $i \in [0..N]$

The simplest assignment of texture coordinates would be:

Note that you should include the rotation angles for  $i = 0$  and  $i = N$ , even though they produce the same points (after rotating by 0 and  $2\pi$ ). Why do this??



## Texture coordinates on a surface of revolution



We can get distortion in  $v$  if the samples are not evenly spaced along the profile curve.

We can reduce this distortion in  $v$ . Define:

$$d[j] = \begin{cases} \|C[j] - C[j-1]\|, & \text{if } j \neq 0 \\ 0, & \text{if } j = 0 \end{cases}$$

and set  $v$  to fractional distance along the curve:

**You must do this for  $v$  for the programming assignment!**

## **Triangle meshes**

How should we generally represent triangle meshes?